

Cancer Mortality by Race, Illinois, 1986-2006										
Lymphoma, Both Sexes										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	21,186	8.7	0.06	8.6	8.8	19,287	9.1	0.07	9.0	9.2
1986	838	7.9	0.28	7.3	8.4	774	8.2	0.30	7.6	8.8
1987	836	7.8	0.28	7.3	8.4	759	8.1	0.30	7.5	8.7
1988	893	8.4	0.28	7.8	8.9	817	8.7	0.31	8.1	9.3
1989	937	8.6	0.29	8.1	9.2	860	9.0	0.31	8.4	9.6
1990	935	8.6	0.28	8.1	9.2	851	8.9	0.31	8.3	9.5
1991	1,002	9.1	0.29	8.5	9.6	918	9.5	0.31	8.9	10.1
1992	1,032	9.2	0.29	8.7	9.8	945	9.6	0.32	9.0	10.3
1993	1,041	9.2	0.29	8.7	9.8	945	9.6	0.31	9.0	10.2
1994	1,092	9.6	0.29	9.0	10.2	981	9.9	0.32	9.3	10.5
1995	1,138	9.9	0.29	9.3	10.5	1,030	10.2	0.32	9.6	10.9
1996	1,118	9.6	0.29	9.1	10.2	1,009	10.0	0.31	9.4	10.6
1997	1,099	9.4	0.28	8.8	10.0	1,001	9.8	0.31	9.2	10.5
1998	1,080	9.1	0.28	8.6	9.7	970	9.4	0.30	8.8	10.0
1999	1,118	9.4	0.28	8.8	10.0	1,026	9.9	0.31	9.3	10.5
2000	1,065	8.9	0.27	8.3	9.4	977	9.4	0.30	8.8	10.0
2001	1,041	8.6	0.27	8.0	9.1	950	9.0	0.29	8.5	9.6
2002	1,008	8.2	0.26	7.7	8.7	911	8.6	0.28	8.0	9.1
2003	979	7.8	0.25	7.4	8.3	895	8.3	0.28	7.8	8.9
2004	991	7.9	0.25	7.4	8.4	908	8.5	0.28	7.9	9.0
2005	1,000	7.9	0.25	7.4	8.4	905	8.3	0.28	7.8	8.8
2006	943	7.4	0.24	6.9	7.9	855	7.8	0.27	7.3	8.3
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	1,694	6.0	0.15	5.7	6.3	205	4.2	0.32	3.6	4.8
1986	60	5.3	0.71	4.0	6.9	4	3.0	1.78	0.6	8.3
1987	72	6.0	0.74	4.6	7.6	5	4.0	1.89	1.2	9.3
1988	65	5.1	0.65	3.9	6.6	11	8.7	3.06	3.8	16.4
1989	68	5.3	0.67	4.1	6.8	9	5.6	2.00	2.4	10.9
1990	81	6.8	0.79	5.4	8.5	3	2.2	1.42	0.4	6.5
1991	79	6.0	0.71	4.7	7.6	5	3.3	1.63	1.0	7.9
1992	85	6.6	0.75	5.2	8.2	2	1.7	1.19	0.2	5.3
1993	91	7.0	0.76	5.6	8.6	5	3.7	1.76	1.1	8.4
1994	101	7.7	0.79	6.2	9.4	10	5.5	1.90	2.4	10.3
1995	98	7.4	0.78	6.0	9.1	10	5.5	1.94	2.4	10.3
1996	102	7.6	0.78	6.2	9.3	7	2.6	1.08	0.9	5.6
1997	83	6.0	0.68	4.8	7.5	15	5.7	1.61	3.0	9.7
1998	101	7.4	0.76	6.0	9.1	9	4.6	1.70	1.9	8.9
1999	86	6.2	0.68	4.9	7.7	6	1.6	0.66	0.6	3.6
2000	75	5.3	0.63	4.1	6.7	13	3.6	1.16	1.7	6.6
2001	80	5.5	0.62	4.3	6.8	11	2.6	0.84	1.2	4.9
2002	82	5.6	0.63	4.4	6.9	15	4.2	1.18	2.3	7.2
2003	65	4.4	0.56	3.4	5.7	19	6.6	1.62	3.8	10.4
2004	67	4.5	0.55	3.4	5.7	16	4.1	1.09	2.3	6.8
2005	77	5.1	0.59	4.0	6.4	18	5.3	1.30	3.0	8.3
2006	76	5.0	0.58	3.9	6.2	12	3.0	0.91	1.5	5.3
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Lymphoma, Males										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	11,063	11.1	0.11	10.8	11.3	9,990	11.5	0.12	11.3	11.7
1986	422	9.4	0.48	8.5	10.4	382	9.7	0.51	8.7	10.7
1987	431	9.6	0.48	8.7	10.6	380	9.7	0.51	8.7	10.7
1988	457	10.4	0.51	9.4	11.5	417	10.9	0.55	9.8	12.0
1989	488	10.8	0.51	9.8	11.8	442	11.2	0.55	10.1	12.3
1990	527	11.8	0.53	10.7	12.9	481	12.2	0.58	11.1	13.4
1991	520	11.3	0.51	10.3	12.3	472	11.7	0.56	10.7	12.9
1992	550	12.0	0.53	10.9	13.0	495	12.3	0.57	11.2	13.5
1993	513	11.0	0.50	10.1	12.1	461	11.4	0.54	10.3	12.5
1994	571	12.2	0.53	11.2	13.3	508	12.5	0.57	11.4	13.7
1995	585	12.4	0.52	11.4	13.4	529	12.8	0.57	11.7	14.0
1996	581	12.1	0.52	11.2	13.2	515	12.3	0.55	11.3	13.5
1997	560	11.5	0.50	10.6	12.5	496	11.8	0.54	10.7	12.9
1998	584	12.2	0.51	11.2	13.2	519	12.5	0.56	11.4	13.6
1999	603	12.4	0.51	11.4	13.5	551	13.1	0.56	12.0	14.2
2000	570	11.4	0.49	10.5	12.4	520	12.1	0.53	11.0	13.2
2001	547	10.9	0.47	10.0	11.9	502	11.6	0.52	10.6	12.7
2002	513	10.3	0.46	9.4	11.2	468	10.9	0.51	9.9	11.9
2003	494	9.7	0.44	8.9	10.6	446	10.2	0.48	9.2	11.2
2004	515	10.0	0.45	9.2	11.0	474	10.7	0.50	9.8	11.8
2005	517	10.1	0.45	9.3	11.1	467	10.6	0.49	9.6	11.6
2006	515	9.7	0.43	8.9	10.6	465	10.3	0.48	9.3	11.2
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	948	7.8	0.27	7.3	8.3	125	5.7	0.58	4.7	7.0
1986	37	7.7	1.30	5.3	10.6	3	6.3	4.25	0.9	19.2
1987	48	9.2	1.43	6.6	12.4	3	3.7	2.14	0.8	11.8
1988	34	6.0	1.07	4.1	8.5	6	10.3	4.84	3.1	23.6
1989	40	7.0	1.19	4.9	9.8	6	9.8	4.36	3.2	21.8
1990	44	8.7	1.44	6.1	11.9	2	2.4	1.69	0.3	9.2
1991	44	7.8	1.28	5.5	10.7	4	6.8	3.77	1.6	17.5
1992	55	10.3	1.52	7.5	13.7	0				
1993	49	8.6	1.32	6.2	11.5	3	4.7	3.09	0.7	13.7
1994	56	9.7	1.38	7.2	12.8	7	7.0	3.01	2.4	15.3
1995	52	9.0	1.35	6.5	12.0	4	5.7	3.10	1.3	14.4
1996	62	11.8	1.65	8.8	15.4	4	3.2	1.83	0.7	9.0
1997	51	8.2	1.21	6.0	11.0	13	11.4	3.56	5.6	20.4
1998	58	10.0	1.38	7.4	13.0	7	8.2	3.62	2.7	17.6
1999	48	8.2	1.26	5.9	11.0	4	2.1	1.10	0.6	6.1
2000	41	6.8	1.14	4.7	9.3	9	6.8	2.84	2.5	14.2
2001	39	6.2	1.04	4.3	8.5	6	3.1	1.32	1.1	7.1
2002	35	5.7	1.00	3.9	7.9	10	5.4	1.95	2.3	10.6
2003	38	6.2	1.06	4.3	8.6	10	8.0	2.78	3.6	15.0
2004	34	5.5	0.99	3.7	7.7	7	4.6	1.89	1.7	9.6
2005	42	6.8	1.09	4.8	9.2	8	6.6	2.45	2.7	12.7
2006	41	6.3	1.03	4.5	8.6	9	4.6	1.62	2.0	8.8
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Lymphoma, Females										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	10,123	7.0	0.07	6.9	7.1	9,297	7.4	0.08	7.2	7.5
1986	416	6.6	0.33	6.0	7.3	392	7.1	0.36	6.4	7.9
1987	405	6.5	0.33	5.8	7.1	379	6.8	0.36	6.1	7.6
1988	436	6.9	0.33	6.3	7.6	400	7.1	0.36	6.5	7.9
1989	449	7.1	0.34	6.4	7.8	418	7.4	0.37	6.7	8.2
1990	408	6.2	0.31	5.6	6.9	370	6.4	0.33	5.7	7.0
1991	482	7.3	0.34	6.7	8.0	446	7.6	0.37	6.9	8.4
1992	482	7.3	0.33	6.6	7.9	450	7.7	0.37	7.0	8.5
1993	528	7.8	0.34	7.2	8.5	484	8.2	0.38	7.4	8.9
1994	521	7.8	0.34	7.1	8.5	473	8.1	0.38	7.4	8.9
1995	553	8.1	0.34	7.4	8.8	501	8.3	0.38	7.6	9.1
1996	537	7.8	0.34	7.2	8.5	494	8.2	0.37	7.5	9.0
1997	539	7.7	0.34	7.1	8.4	505	8.3	0.37	7.6	9.1
1998	496	7.0	0.32	6.4	7.7	451	7.4	0.35	6.7	8.1
1999	515	7.3	0.32	6.6	7.9	475	7.7	0.36	7.0	8.4
2000	495	7.0	0.32	6.4	7.6	457	7.4	0.35	6.8	8.2
2001	494	6.9	0.31	6.3	7.5	448	7.2	0.34	6.5	7.9
2002	495	6.8	0.31	6.2	7.4	443	7.0	0.34	6.4	7.7
2003	485	6.4	0.30	5.9	7.1	449	6.9	0.33	6.3	7.6
2004	476	6.5	0.30	5.9	7.1	434	6.9	0.34	6.2	7.6
2005	483	6.4	0.30	5.8	7.0	438	6.7	0.33	6.1	7.4
2006	428	5.7	0.28	5.1	6.2	390	6.0	0.31	5.4	6.7
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	746	4.6	0.17	4.3	5.0	80	3.0	0.35	2.3	3.7
1986	23	3.5	0.75	2.2	5.3	1	0.7	0.68	0.0	5.9
1987	24	3.6	0.76	2.3	5.4	2	3.9	2.79	0.5	12.6
1988	31	4.4	0.80	2.9	6.2	5	7.5	3.90	1.9	18.2
1989	28	4.0	0.77	2.6	5.8	3	2.8	1.62	0.6	8.6
1990	37	5.4	0.91	3.8	7.5	1	2.1	2.12	0.1	9.2
1991	35	4.7	0.82	3.3	6.6	1	0.9	0.87	0.0	5.2
1992	30	4.1	0.77	2.7	5.9	2	2.9	2.01	0.3	9.1
1993	42	5.7	0.89	4.1	7.7	2	2.9	2.08	0.4	9.2
1994	45	6.0	0.92	4.4	8.1	3	4.1	2.39	0.8	10.9
1995	46	6.2	0.92	4.5	8.2	6	5.3	2.43	1.7	11.9
1996	40	5.0	0.81	3.6	6.9	3	2.0	1.28	0.3	6.2
1997	32	4.2	0.75	2.8	5.9	2	1.3	1.04	0.1	5.0
1998	43	5.5	0.85	4.0	7.4	2	2.1	1.56	0.2	6.8
1999	38	4.7	0.78	3.3	6.5	2	1.1	0.76	0.1	3.9
2000	34	4.2	0.72	2.9	5.8	4	1.5	0.80	0.4	4.3
2001	41	4.9	0.77	3.5	6.6	5	2.2	1.05	0.6	5.4
2002	47	5.5	0.81	4.0	7.3	5	3.2	1.44	1.0	7.2
2003	27	3.2	0.62	2.1	4.7	9	5.4	1.90	2.4	10.2
2004	33	3.8	0.66	2.6	5.3	9	3.9	1.35	1.7	7.5
2005	35	3.9	0.67	2.7	5.4	10	4.7	1.54	2.2	8.6
2006	35	3.9	0.67	2.7	5.4	3	1.7	0.97	0.3	4.5
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Hodgkin Lymphoma, Both Sexes										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	1,371	0.6	0.01	0.5	0.6	1,172	0.6	0.02	0.5	0.6
1986	103	0.9	0.09	0.7	1.1	94	1.0	0.10	0.8	1.2
1987	90	0.8	0.09	0.6	1.0	77	0.8	0.09	0.6	1.0
1988	73	0.7	0.08	0.5	0.8	62	0.7	0.09	0.5	0.9
1989	70	0.6	0.07	0.5	0.8	60	0.6	0.08	0.5	0.8
1990	71	0.6	0.07	0.5	0.8	59	0.6	0.08	0.4	0.8
1991	72	0.6	0.07	0.5	0.8	59	0.6	0.08	0.5	0.8
1992	67	0.6	0.07	0.4	0.7	56	0.6	0.08	0.4	0.7
1993	65	0.6	0.07	0.4	0.7	56	0.6	0.08	0.4	0.7
1994	68	0.6	0.07	0.5	0.7	56	0.6	0.07	0.4	0.7
1995	58	0.5	0.07	0.4	0.6	48	0.5	0.07	0.4	0.6
1996	60	0.5	0.07	0.4	0.7	51	0.5	0.07	0.4	0.7
1997	60	0.5	0.07	0.4	0.7	53	0.5	0.07	0.4	0.7
1998	61	0.5	0.07	0.4	0.7	49	0.5	0.07	0.4	0.6
1999	57	0.5	0.06	0.4	0.6	51	0.5	0.07	0.4	0.7
2000	61	0.5	0.06	0.4	0.6	53	0.5	0.07	0.4	0.7
2001	54	0.4	0.06	0.3	0.6	48	0.5	0.07	0.3	0.6
2002	65	0.5	0.07	0.4	0.7	57	0.6	0.07	0.4	0.7
2003	52	0.4	0.06	0.3	0.6	46	0.4	0.07	0.3	0.6
2004	56	0.4	0.06	0.3	0.6	49	0.5	0.07	0.3	0.6
2005	50	0.4	0.06	0.3	0.5	41	0.4	0.06	0.3	0.5
2006	58	0.5	0.06	0.4	0.6	47	0.5	0.07	0.3	0.6
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	181	0.5	0.04	0.5	0.6	18	0.3	0.06	0.1	0.4
1986	9	0.6	0.23	0.3	1.3	0				
1987	12	0.7	0.22	0.4	1.3	1	0.6	0.56	0.0	3.6
1988	10	0.6	0.21	0.3	1.2	1	0.4	0.41	0.0	3.1
1989	9	0.6	0.20	0.3	1.1	1	0.5	0.50	0.0	3.2
1990	12	0.8	0.24	0.4	1.4	0				
1991	12	0.7	0.21	0.4	1.3	1	0.5	0.48	0.0	2.9
1992	11	0.6	0.20	0.3	1.2	0				
1993	9	0.6	0.20	0.3	1.1	0				
1994	10	0.6	0.20	0.3	1.2	2	0.7	0.49	0.1	2.8
1995	10	0.6	0.20	0.3	1.1	0				
1996	8	0.5	0.19	0.2	1.0	1	0.6	0.63	0.0	2.9
1997	5	0.3	0.14	0.1	0.8	2	0.5	0.34	0.1	2.1
1998	11	0.8	0.23	0.4	1.4	1	0.2	0.23	0.0	1.6
1999	6	0.4	0.15	0.1	0.8	0				
2000	7	0.5	0.18	0.2	1.0	1	0.1	0.14	0.0	1.3
2001	5	0.3	0.12	0.1	0.7	1	0.2	0.19	0.0	1.3
2002	6	0.3	0.14	0.1	0.7	2	0.4	0.27	0.1	1.5
2003	5	0.3	0.15	0.1	0.8	1	0.1	0.14	0.0	1.1
2004	7	0.4	0.13	0.1	0.7	0				
2005	8	0.5	0.16	0.2	0.9	1	0.2	0.15	0.0	1.0
2006	9	0.6	0.21	0.3	1.2	2	0.4	0.32	0.1	1.6
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Hodgkin Lymphoma, Males										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	773	0.7	0.03	0.6	0.7	655	0.7	0.03	0.6	0.8
1986	47	0.9	0.14	0.7	1.2	43	1.0	0.15	0.7	1.3
1987	56	1.0	0.14	0.8	1.3	45	1.0	0.15	0.7	1.3
1988	40	0.8	0.13	0.6	1.1	35	0.8	0.14	0.6	1.1
1989	40	0.8	0.13	0.6	1.1	35	0.8	0.14	0.6	1.2
1990	46	0.9	0.13	0.6	1.2	38	0.8	0.14	0.6	1.2
1991	40	0.8	0.13	0.5	1.1	33	0.7	0.13	0.5	1.1
1992	36	0.7	0.12	0.5	1.0	29	0.7	0.12	0.4	0.9
1993	25	0.5	0.10	0.3	0.7	19	0.4	0.10	0.3	0.7
1994	37	0.7	0.11	0.5	0.9	31	0.7	0.12	0.4	0.9
1995	35	0.6	0.11	0.4	0.9	30	0.7	0.12	0.4	0.9
1996	35	0.7	0.12	0.5	1.0	31	0.7	0.13	0.5	1.0
1997	30	0.6	0.11	0.4	0.9	24	0.6	0.12	0.4	0.9
1998	43	0.8	0.12	0.6	1.1	33	0.7	0.13	0.5	1.0
1999	36	0.7	0.12	0.5	1.0	33	0.8	0.13	0.5	1.1
2000	34	0.6	0.11	0.4	0.9	29	0.6	0.12	0.4	0.9
2001	31	0.6	0.10	0.4	0.8	30	0.6	0.12	0.4	0.9
2002	38	0.7	0.12	0.5	1.0	32	0.7	0.13	0.5	1.0
2003	28	0.5	0.10	0.3	0.7	24	0.5	0.11	0.3	0.8
2004	37	0.7	0.11	0.5	0.9	35	0.7	0.12	0.5	1.0
2005	24	0.4	0.09	0.3	0.6	18	0.4	0.09	0.2	0.6
2006	35	0.6	0.11	0.4	0.9	28	0.6	0.11	0.4	0.9
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	105	0.7	0.07	0.6	0.9	13	0.4	0.13	0.2	0.8
1986	4	0.6	0.33	0.2	1.8	0				
1987	10	1.3	0.44	0.6	2.6	1	1.2	1.23	0.0	8.2
1988	4	0.5	0.25	0.1	1.4	1	0.8	0.82	0.0	7.0
1989	4	0.5	0.24	0.1	1.4	1	1.1	1.13	0.0	7.4
1990	8	1.2	0.44	0.5	2.5	0				
1991	6	0.9	0.37	0.3	2.0	1	1.1	1.11	0.0	6.9
1992	7	1.0	0.38	0.4	2.1	0				
1993	6	1.0	0.50	0.3	2.4	0				
1994	4	0.6	0.32	0.2	1.7	2	1.5	1.11	0.2	6.4
1995	5	0.6	0.27	0.2	1.5	0				
1996	3	0.4	0.26	0.1	1.4	1	1.6	1.56	0.0	7.2
1997	5	0.7	0.34	0.2	1.8	1	0.6	0.62	0.0	4.3
1998	9	1.5	0.53	0.7	2.9	1	0.5	0.47	0.0	3.8
1999	3	0.4	0.23	0.1	1.2	0				
2000	5	0.9	0.41	0.3	2.0	0				
2001	1	0.1	0.09	0.0	0.7	0				
2002	4	0.5	0.23	0.1	1.3	2	0.8	0.56	0.1	3.5
2003	4	0.5	0.27	0.2	1.4	0				
2004	2	0.2	0.16	0.0	0.9	0				
2005	6	0.8	0.34	0.3	1.8	0				
2006	5	0.8	0.37	0.3	1.9	2	1.0	0.71	0.1	3.5
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Hodgkin Lymphoma, Females										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	598	0.4	0.02	0.4	0.5	517	0.4	0.02	0.4	0.5
1986	56	0.9	0.12	0.7	1.2	51	1.0	0.14	0.7	1.3
1987	34	0.6	0.10	0.4	0.8	32	0.6	0.12	0.4	0.9
1988	33	0.5	0.09	0.4	0.8	27	0.5	0.10	0.3	0.8
1989	30	0.5	0.09	0.3	0.7	25	0.5	0.09	0.3	0.7
1990	25	0.4	0.08	0.2	0.6	21	0.4	0.08	0.2	0.5
1991	32	0.5	0.09	0.3	0.7	26	0.5	0.09	0.3	0.7
1992	31	0.5	0.09	0.3	0.7	27	0.5	0.10	0.3	0.7
1993	40	0.6	0.10	0.4	0.8	37	0.7	0.11	0.5	0.9
1994	31	0.5	0.09	0.3	0.7	25	0.4	0.09	0.3	0.7
1995	23	0.4	0.07	0.2	0.5	18	0.3	0.07	0.2	0.5
1996	25	0.4	0.08	0.3	0.6	20	0.4	0.08	0.2	0.6
1997	30	0.4	0.08	0.3	0.6	29	0.5	0.09	0.3	0.7
1998	18	0.3	0.06	0.2	0.4	16	0.3	0.07	0.2	0.4
1999	21	0.3	0.07	0.2	0.5	18	0.3	0.08	0.2	0.5
2000	27	0.4	0.08	0.3	0.6	24	0.4	0.09	0.3	0.7
2001	23	0.3	0.07	0.2	0.5	18	0.3	0.07	0.2	0.5
2002	27	0.4	0.08	0.3	0.6	25	0.5	0.09	0.3	0.7
2003	24	0.4	0.07	0.2	0.5	22	0.4	0.08	0.3	0.6
2004	19	0.3	0.06	0.2	0.4	14	0.3	0.07	0.1	0.4
2005	26	0.4	0.08	0.3	0.6	23	0.4	0.09	0.3	0.6
2006	23	0.3	0.07	0.2	0.5	19	0.3	0.08	0.2	0.5
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	76	0.4	0.05	0.3	0.5	5	0.1	0.04	0.0	0.2
1986	5	0.6	0.30	0.2	1.5	0				
1987	2	0.2	0.17	0.0	0.9	0				
1988	6	0.7	0.31	0.3	1.6	0				
1989	5	0.6	0.29	0.2	1.5	0				
1990	4	0.5	0.26	0.1	1.3	0				
1991	6	0.6	0.24	0.2	1.3	0				
1992	4	0.4	0.19	0.1	1.0	0				
1993	3	0.4	0.20	0.1	1.0	0				
1994	6	0.6	0.26	0.2	1.4	0				
1995	5	0.6	0.27	0.2	1.4	0				
1996	5	0.6	0.27	0.2	1.4	0				
1997	0					1	0.4	0.35	0.0	2.9
1998	2	0.2	0.15	0.0	0.8	0				
1999	3	0.3	0.19	0.1	1.0	0				
2000	2	0.2	0.15	0.0	0.8	1	0.3	0.29	0.0	2.3
2001	4	0.4	0.20	0.1	1.1	1	0.4	0.36	0.0	2.4
2002	2	0.2	0.15	0.0	0.8	0				
2003	1	0.1	0.13	0.0	0.7	1	0.3	0.30	0.0	2.1
2004	5	0.5	0.21	0.2	1.1	0				
2005	2	0.2	0.13	0.0	0.7	1	0.3	0.30	0.0	2.0
2006	4	0.5	0.22	0.1	1.1	0				
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Non-Hodgkin Lymphoma, Both Sexes										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	19,815	8.2	0.06	8.0	8.3	18,115	8.5	0.06	8.4	8.7
1986	735	7.0	0.26	6.5	7.5	680	7.3	0.28	6.7	7.8
1987	746	7.1	0.26	6.6	7.6	682	7.3	0.28	6.7	7.9
1988	820	7.7	0.27	7.2	8.2	755	8.0	0.29	7.4	8.6
1989	867	8.0	0.28	7.5	8.6	800	8.4	0.30	7.8	9.0
1990	864	8.0	0.27	7.5	8.6	792	8.3	0.30	7.7	8.9
1991	930	8.4	0.28	7.9	9.0	859	8.9	0.30	8.3	9.5
1992	965	8.6	0.28	8.1	9.2	889	9.1	0.31	8.5	9.7
1993	976	8.7	0.28	8.1	9.2	889	9.0	0.30	8.4	9.6
1994	1,024	9.0	0.28	8.5	9.6	925	9.3	0.31	8.7	9.9
1995	1,080	9.4	0.29	8.8	10.0	982	9.8	0.31	9.1	10.4
1996	1,058	9.1	0.28	8.6	9.7	958	9.5	0.31	8.9	10.1
1997	1,039	8.9	0.28	8.4	9.4	948	9.3	0.30	8.7	9.9
1998	1,019	8.6	0.27	8.1	9.2	921	8.9	0.29	8.4	9.5
1999	1,061	8.9	0.27	8.4	9.5	975	9.4	0.30	8.8	10.0
2000	1,004	8.4	0.26	7.8	8.9	924	8.9	0.29	8.3	9.5
2001	987	8.1	0.26	7.6	8.6	902	8.6	0.29	8.0	9.2
2002	943	7.7	0.25	7.2	8.2	854	8.0	0.27	7.5	8.6
2003	927	7.4	0.24	7.0	7.9	849	7.9	0.27	7.3	8.4
2004	935	7.5	0.25	7.0	8.0	859	8.0	0.27	7.5	8.5
2005	950	7.5	0.24	7.0	8.0	864	7.9	0.27	7.4	8.4
2006	885	6.9	0.23	6.5	7.4	808	7.4	0.26	6.9	7.9
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	1,513	5.4	0.14	5.2	5.7	187	3.9	0.31	3.3	4.6
1986	51	4.7	0.67	3.5	6.2	4	3.0	1.78	0.6	8.3
1987	60	5.2	0.70	4.0	6.8	4	3.4	1.81	0.8	8.7
1988	55	4.5	0.62	3.3	5.8	10	8.3	3.03	3.5	15.9
1989	59	4.7	0.64	3.6	6.2	8	5.1	1.94	2.1	10.3
1990	69	6.0	0.75	4.6	7.6	3	2.2	1.42	0.4	6.5
1991	67	5.3	0.68	4.1	6.8	4	2.9	1.55	0.7	7.3
1992	74	6.0	0.72	4.6	7.5	2	1.7	1.19	0.2	5.3
1993	82	6.4	0.73	5.0	8.0	5	3.7	1.76	1.1	8.4
1994	91	7.1	0.76	5.6	8.7	8	4.8	1.84	1.9	9.5
1995	88	6.9	0.75	5.5	8.5	10	5.5	1.94	2.4	10.3
1996	94	7.1	0.75	5.7	8.7	6	1.9	0.88	0.6	4.5
1997	78	5.7	0.66	4.5	7.1	13	5.2	1.57	2.6	9.1
1998	90	6.7	0.72	5.3	8.2	8	4.4	1.68	1.7	8.6
1999	80	5.8	0.67	4.6	7.3	6	1.6	0.66	0.6	3.6
2000	68	4.8	0.60	3.7	6.1	12	3.5	1.15	1.6	6.4
2001	75	5.2	0.61	4.1	6.5	10	2.4	0.82	1.1	4.6
2002	76	5.3	0.61	4.1	6.6	13	3.9	1.15	1.9	6.7
2003	60	4.1	0.54	3.1	5.3	18	6.4	1.61	3.7	10.2
2004	60	4.1	0.54	3.1	5.3	16	4.1	1.09	2.3	6.8
2005	69	4.6	0.57	3.6	5.9	17	5.1	1.29	2.9	8.2
2006	67	4.3	0.54	3.4	5.5	10	2.6	0.86	1.2	4.8
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Non-Hodgkin Lymphoma, Males										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	10,290	10.4	0.10	10.2	10.6	9,335	10.8	0.11	10.6	11.0
1986	375	8.5	0.46	7.6	9.5	339	8.7	0.49	7.8	9.7
1987	375	8.5	0.46	7.7	9.5	335	8.7	0.49	7.7	9.7
1988	417	9.6	0.49	8.7	10.6	382	10.1	0.53	9.0	11.2
1989	448	10.0	0.49	9.1	11.0	407	10.3	0.53	9.3	11.4
1990	481	10.9	0.52	9.9	12.0	443	11.4	0.56	10.3	12.5
1991	480	10.5	0.50	9.6	11.5	439	11.0	0.54	10.0	12.1
1992	514	11.3	0.52	10.3	12.3	466	11.7	0.56	10.6	12.8
1993	488	10.6	0.49	9.6	11.6	442	11.0	0.54	9.9	12.1
1994	534	11.6	0.51	10.6	12.6	477	11.9	0.56	10.8	13.0
1995	550	11.7	0.51	10.7	12.8	499	12.2	0.56	11.1	13.3
1996	546	11.4	0.50	10.5	12.5	484	11.6	0.54	10.6	12.7
1997	530	10.9	0.48	10.0	11.9	472	11.2	0.52	10.2	12.3
1998	541	11.4	0.50	10.4	12.4	486	11.8	0.54	10.7	12.9
1999	567	11.7	0.50	10.8	12.7	518	12.3	0.55	11.3	13.4
2000	536	10.8	0.47	9.9	11.8	491	11.4	0.52	10.4	12.5
2001	516	10.3	0.46	9.5	11.3	472	11.0	0.51	10.0	12.0
2002	475	9.6	0.45	8.7	10.5	436	10.1	0.49	9.2	11.1
2003	466	9.2	0.43	8.4	10.1	422	9.6	0.47	8.7	10.6
2004	478	9.4	0.44	8.6	10.3	439	10.0	0.48	9.1	11.0
2005	493	9.7	0.44	8.9	10.6	449	10.2	0.49	9.3	11.2
2006	480	9.1	0.42	8.3	10.0	437	9.7	0.47	8.8	10.6
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	843	7.1	0.26	6.6	7.6	112	5.3	0.57	4.3	6.5
1986	33	7.0	1.25	4.8	9.9	3	6.3	4.25	0.9	19.2
1987	38	7.8	1.36	5.4	10.9	2	2.5	1.75	0.3	10.0
1988	30	5.5	1.04	3.7	7.9	5	9.5	4.77	2.6	22.7
1989	36	6.6	1.17	4.5	9.3	5	8.7	4.21	2.5	20.4
1990	36	7.5	1.37	5.1	10.6	2	2.4	1.69	0.3	9.2
1991	38	6.9	1.22	4.7	9.7	3	5.7	3.60	1.0	16.2
1992	48	9.3	1.47	6.6	12.6	0				
1993	43	7.6	1.22	5.4	10.3	3	4.7	3.09	0.7	13.7
1994	52	9.1	1.35	6.7	12.1	5	5.5	2.80	1.4	13.5
1995	47	8.4	1.32	6.0	11.3	4	5.7	3.10	1.3	14.4
1996	59	11.4	1.62	8.4	15.0	3	1.6	0.96	0.3	5.8
1997	46	7.5	1.16	5.4	10.1	12	10.8	3.51	5.1	19.7
1998	49	8.4	1.28	6.1	11.3	6	7.7	3.59	2.4	17.1
1999	45	7.8	1.24	5.6	10.6	4	2.1	1.10	0.6	6.1
2000	36	5.9	1.07	4.0	8.3	9	6.8	2.84	2.5	14.2
2001	38	6.1	1.04	4.2	8.4	6	3.1	1.32	1.1	7.1
2002	31	5.2	0.98	3.5	7.4	8	4.6	1.87	1.7	9.7
2003	34	5.6	1.03	3.8	8.0	10	8.0	2.78	3.6	15.0
2004	32	5.2	0.97	3.5	7.5	7	4.6	1.89	1.7	9.6
2005	36	6.0	1.04	4.1	8.3	8	6.6	2.45	2.7	12.7
2006	36	5.5	0.96	3.8	7.7	7	3.6	1.45	1.4	7.6
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										

Cancer Mortality by Race, Illinois, 1986-2006										
Non-Hodgkin Lymphoma, Females										
	All Races					Whites				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	9,525	6.6	0.07	6.4	6.7	8,780	6.9	0.07	6.8	7.1
1986	360	5.7	0.31	5.1	6.4	341	6.1	0.34	5.5	6.8
1987	371	5.9	0.31	5.3	6.5	347	6.2	0.34	5.5	6.9
1988	403	6.4	0.32	5.7	7.0	373	6.6	0.35	6.0	7.3
1989	419	6.6	0.33	6.0	7.3	393	7.0	0.36	6.3	7.7
1990	383	5.9	0.30	5.3	6.5	349	6.0	0.33	5.4	6.7
1991	450	6.8	0.33	6.2	7.5	420	7.2	0.35	6.5	7.9
1992	451	6.8	0.32	6.2	7.4	423	7.2	0.36	6.5	8.0
1993	488	7.2	0.33	6.6	7.9	447	7.5	0.36	6.8	8.2
1994	490	7.3	0.33	6.7	8.0	448	7.6	0.37	6.9	8.4
1995	530	7.7	0.34	7.1	8.4	483	8.0	0.37	7.3	8.8
1996	512	7.4	0.33	6.8	8.1	474	7.9	0.37	7.2	8.6
1997	509	7.3	0.33	6.7	8.0	476	7.8	0.36	7.1	8.5
1998	478	6.8	0.31	6.2	7.4	435	7.1	0.34	6.4	7.8
1999	494	6.9	0.31	6.3	7.6	457	7.3	0.35	6.7	8.1
2000	468	6.5	0.31	6.0	7.2	433	7.0	0.34	6.3	7.7
2001	471	6.5	0.30	5.9	7.1	430	6.9	0.34	6.2	7.6
2002	468	6.4	0.30	5.8	7.0	418	6.6	0.33	5.9	7.2
2003	461	6.1	0.29	5.5	6.7	427	6.5	0.32	5.9	7.2
2004	457	6.2	0.29	5.7	6.8	420	6.6	0.33	6.0	7.3
2005	457	6.0	0.29	5.5	6.6	415	6.3	0.32	5.7	7.0
2006	405	5.3	0.27	4.8	5.9	371	5.7	0.30	5.1	6.3
	Blacks					Asian and Other Races				
Years	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2006	670	4.2	0.16	3.9	4.5	75	2.9	0.35	2.2	3.6
1986	18	2.9	0.69	1.7	4.5	1	0.7	0.68	0.0	5.9
1987	22	3.4	0.74	2.1	5.1	2	3.9	2.79	0.5	12.6
1988	25	3.6	0.74	2.3	5.4	5	7.5	3.90	1.9	18.2
1989	23	3.3	0.71	2.1	5.0	3	2.8	1.62	0.6	8.6
1990	33	4.9	0.87	3.4	6.9	1	2.1	2.12	0.1	9.2
1991	29	4.1	0.78	2.7	6.0	1	0.9	0.87	0.0	5.2
1992	26	3.7	0.74	2.4	5.4	2	2.9	2.01	0.3	9.1
1993	39	5.3	0.87	3.8	7.3	2	2.9	2.08	0.4	9.2
1994	39	5.4	0.88	3.8	7.4	3	4.1	2.39	0.8	10.9
1995	41	5.6	0.88	4.0	7.6	6	5.3	2.43	1.7	11.9
1996	35	4.4	0.76	3.1	6.2	3	2.0	1.28	0.3	6.2
1997	32	4.2	0.75	2.8	5.9	1	1.0	0.98	0.0	4.6
1998	41	5.3	0.84	3.8	7.2	2	2.1	1.56	0.2	6.8
1999	35	4.4	0.75	3.1	6.1	2	1.1	0.76	0.1	3.9
2000	32	4.0	0.71	2.7	5.6	3	1.2	0.75	0.2	3.9
2001	37	4.5	0.74	3.1	6.1	4	1.8	0.98	0.4	4.9
2002	45	5.3	0.80	3.9	7.1	5	3.2	1.44	1.0	7.2
2003	26	3.1	0.61	2.0	4.5	8	5.1	1.87	2.1	9.9
2004	28	3.3	0.63	2.2	4.8	9	3.9	1.35	1.7	7.5
2005	33	3.7	0.65	2.6	5.2	9	4.4	1.51	1.9	8.3
2006	31	3.5	0.63	2.3	4.9	3	1.7	0.97	0.3	4.5
Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.										
Source: Surveillance, Epidemiology and End Results (SEER) Program (www.seer.cancer.gov)SEER*Stat Database: Mortality - All COD, Public-Use with State, Total United States (1969-2006), and Mortality - All COD, Public-Use with State, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released May 2009. Underlying Mortality data provided by NCHS (www.cdc.gov/nchs).										