

Pediatric Cancer Incidence, All Races, Both Sexes, Illinois 1986-2008

VI Renal tumors

Ages 0-14						Ages 0-19				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2008	506	8.1	0.36	7.4	8.9	531	6.4	0.28	5.9	7.0
1986	25	9.5	1.90	6.1	14.0	25	7.1	1.42	4.6	10.5
1987	29	11.1	2.05	7.4	15.9	29	8.3	1.54	5.5	11.9
1988	24	9.2	1.88	5.9	13.7	24	6.9	1.41	4.4	10.3
1989	26	9.9	1.95	6.5	14.6	28	8.0	1.52	5.3	11.6
1990	20	7.7	1.73	4.7	11.9	20	5.8	1.29	3.5	8.9
1991	20	7.6	1.70	4.6	11.7	20	5.7	1.27	3.5	8.8
1992	21	8.0	1.74	4.9	12.2	22	6.3	1.34	3.9	9.5
1993	25	9.1	1.82	5.9	13.5	25	6.8	1.37	4.4	10.1
1994	17	6.2	1.49	3.6	9.9	18	4.9	1.16	2.9	7.8
1995	22	7.9	1.68	4.9	11.9	22	5.9	1.26	3.7	8.9
1996	25	8.9	1.79	5.8	13.2	26	7.0	1.37	4.6	10.3
1997	32	11.5	2.04	7.9	16.3	34	9.2	1.58	6.4	12.9
1998	22	8.0	1.71	5.0	12.2	24	6.6	1.34	4.2	9.8
1999	21	7.6	1.67	4.7	11.7	22	6.0	1.28	3.8	9.1
2000	27	9.9	1.91	6.5	14.4	29	8.0	1.48	5.3	11.5
2001	20	7.3	1.64	4.5	11.3	20	5.5	1.22	3.3	8.5
2002	17	6.3	1.52	3.6	10.0	17	4.7	1.13	2.7	7.5
2003	21	7.6	1.67	4.7	11.7	24	6.6	1.34	4.2	9.8
2004	16	5.8	1.46	3.3	9.5	18	4.9	1.16	2.9	7.8
2005	20	7.4	1.64	4.5	11.4	22	6.1	1.29	3.8	9.2
2006	19	7.0	1.60	4.2	10.9	22	6.1	1.29	3.8	9.2
2007	17	6.2	1.51	3.6	10.0	18	4.9	1.17	2.9	7.8
2008	20	7.4	1.64	4.5	11.4	22	6.1	1.29	3.8	9.2

Rates are per 1 million and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.

Source: Illinois Department of Public Health, Illinois State Cancer Registry, data as of November 2010

Pediatric Cancer Incidence, All Races, Both Sexes, Illinois 1986-2008

VI(a) Nephroblastoma and other nonepithelial renal tumors

Ages 0-14						Ages 0-19				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2008	485	7.8	0.35	7.1	8.5	496	6.0	0.27	5.4	6.5
1986	25	9.5	1.90	6.1	14.0	25	7.1	1.42	4.6	10.5
1987	26	9.8	1.93	6.4	14.4	26	7.4	1.44	4.8	10.8
1988	24	9.2	1.88	5.9	13.7	24	6.9	1.41	4.4	10.3
1989	26	9.9	1.95	6.5	14.6	27	7.7	1.49	5.1	11.3
1990	20	7.7	1.73	4.7	11.9	20	5.8	1.29	3.5	8.9
1991	20	7.6	1.70	4.6	11.7	20	5.7	1.27	3.5	8.8
1992	21	8.0	1.74	4.9	12.2	22	6.3	1.34	3.9	9.5
1993	24	8.8	1.79	5.6	13.1	24	6.6	1.34	4.2	9.8
1994	15	5.4	1.38	3.0	8.9	15	4.0	1.04	2.2	6.6
1995	20	7.1	1.58	4.3	11.0	20	5.3	1.19	3.2	8.2
1996	25	8.9	1.79	5.8	13.2	26	7.0	1.37	4.6	10.3
1997	31	11.1	2.00	7.6	15.8	32	8.6	1.52	5.9	12.2
1998	21	7.7	1.67	4.7	11.7	22	6.0	1.28	3.8	9.1
1999	21	7.6	1.67	4.7	11.7	22	6.0	1.28	3.8	9.1
2000	26	9.5	1.87	6.2	14.0	26	7.1	1.40	4.7	10.5
2001	19	7.0	1.60	4.2	10.9	19	5.2	1.20	3.1	8.1
2002	17	6.3	1.52	3.6	10.0	17	4.7	1.13	2.7	7.5
2003	21	7.6	1.67	4.7	11.7	22	6.0	1.28	3.8	9.1
2004	12	4.4	1.27	2.3	7.7	12	3.3	0.95	1.7	5.7
2005	19	7.0	1.60	4.2	10.9	21	5.8	1.26	3.6	8.8
2006	17	6.2	1.51	3.6	10.0	18	4.9	1.16	2.9	7.8
2007	17	6.2	1.51	3.6	10.0	18	4.9	1.17	2.9	7.8
2008	18	6.6	1.55	3.9	10.4	18	4.9	1.16	2.9	7.8

Rates are per 1 million and are age-adjusted to the 2000 U.S. standard million population. Standard error (SE). Lower confidence interval (LCI) and upper confidence interval (UCI) are 95 percent for rates.

Source: Illinois Department of Public Health, Illinois State Cancer Registry, data as of November 2010