

SECTION I
Cancer Mortality by Race
Illinois, 1986-2012

Invasive Cancer Mortality by Race, Illinois 1986-2012

All Sites Combined										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	658,746	203.5	0.25	203.0	204.0	554,139	197.8	0.27	197.3	198.4
1986	23,022	218.7	1.46	215.8	221.6	19,728	210.9	1.52	207.9	213.9
1987	23,182	218.6	1.45	215.8	221.5	19,852	210.9	1.52	208.0	213.9
1988	23,495	220.0	1.45	217.1	222.8	19,995	211.3	1.51	208.4	214.3
1989	23,932	222.8	1.45	220.0	225.7	20,287	213.6	1.52	210.6	216.6
1990	24,440	225.5	1.46	222.7	228.4	20,762	216.6	1.52	213.6	219.6
1991	24,416	222.4	1.43	219.6	225.2	20,836	214.8	1.50	211.9	217.8
1992	24,622	221.0	1.42	218.2	223.8	20,914	213.2	1.48	210.3	216.1
1993	25,269	224.4	1.42	221.7	227.2	21,365	216.0	1.48	213.1	218.9
1994	25,144	221.1	1.40	218.4	223.9	21,334	213.9	1.47	211.0	216.8
1995	25,101	218.6	1.38	215.9	221.3	21,292	211.9	1.46	209.0	214.8
1996	24,785	214.6	1.37	211.9	217.3	20,816	206.3	1.43	203.5	209.2
1997	24,479	210.0	1.34	207.4	212.6	20,609	202.9	1.42	200.1	205.7
1998	24,638	209.1	1.33	206.4	211.7	20,695	201.9	1.41	199.1	204.7
1999	25,024	210.3	1.33	207.7	212.9	21,085	204.0	1.41	201.3	206.8
2000	25,365	211.4	1.33	208.8	214.0	21,284	205.0	1.41	202.2	207.7
2001	24,778	204.8	1.30	202.3	207.4	20,782	199.0	1.38	196.3	201.7
2002	24,737	202.7	1.29	200.2	205.3	20,702	197.0	1.37	194.3	199.7
2003	24,464	198.2	1.27	195.7	200.7	20,437	192.5	1.35	189.9	195.2
2004	24,289	194.9	1.25	192.5	197.4	20,299	190.1	1.34	187.5	192.7
2005	24,250	192.4	1.24	190.0	194.8	20,226	187.6	1.33	185.0	190.2
2006	24,083	188.9	1.22	186.5	191.3	20,070	184.6	1.31	182.0	187.1
2007	24,113	186.6	1.21	184.2	189.0	20,020	182.0	1.30	179.5	184.5
2008	24,298	185.0	1.20	182.7	187.4	20,305	182.0	1.29	179.5	184.6
2009	24,182	181.7	1.18	179.4	184.0	20,205	179.4	1.28	177.0	182.0
2010	24,070	177.9	1.16	175.6	180.2	19,894	174.1	1.25	171.7	176.6
2011	24,006	174.7	1.14	172.5	177.0	19,944	172.3	1.24	169.9	174.8
2012	24,562	175.5	1.14	173.3	177.8	20,401	173.5	1.24	171.1	176.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

All Sites Combined										
Both Sexes										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	97,168	267.7	0.88	266.0	269.4	7,433	94.7	1.20	92.4	97.1
1986	3,189	298.1	5.48	287.4	309.0	104	85.6	9.75	67.6	106.4
1987	3,231	297.8	5.42	287.2	308.6	98	78.1	9.13	61.3	97.7
1988	3,368	301.8	5.38	291.4	312.6	132	95.1	9.54	77.4	115.3
1989	3,491	311.6	5.45	301.0	322.4	152	104.7	9.79	86.4	125.3
1990	3,539	313.3	5.44	302.7	324.1	138	98.9	9.83	80.6	119.6
1991	3,416	298.2	5.26	288.0	308.6	163	103.2	9.30	85.8	122.7
1992	3,536	299.6	5.19	289.5	309.9	172	101.4	8.80	84.9	119.8
1993	3,739	311.9	5.24	301.7	322.3	165	93.1	8.22	77.7	110.3
1994	3,606	295.9	5.06	286.1	306.0	204	105.0	8.35	89.3	122.4
1995	3,599	289.4	4.95	279.8	299.3	210	100.7	7.88	85.8	117.1
1996	3,753	297.9	4.98	288.2	307.8	216	97.0	7.35	83.1	112.3
1997	3,635	282.4	4.79	273.1	291.9	235	100.0	7.28	86.2	115.1
1998	3,694	280.6	4.72	271.4	290.0	249	97.5	6.95	84.4	111.9
1999	3,681	277.9	4.68	268.8	287.2	258	93.5	6.45	81.2	106.8
2000	3,764	277.8	4.62	268.9	287.0	317	110.3	6.82	97.3	124.3
2001	3,697	271.1	4.55	262.3	280.2	299	94.2	5.99	82.9	106.6
2002	3,713	266.9	4.47	258.2	275.8	322	98.5	6.00	87.1	110.9
2003	3,670	259.0	4.37	250.5	267.7	357	106.7	6.13	95.0	119.3
2004	3,672	253.9	4.28	245.5	262.4	318	89.0	5.40	78.7	100.1
2005	3,661	249.3	4.21	241.2	257.7	363	97.8	5.60	87.2	109.3
2006	3,655	244.7	4.14	236.6	252.9	358	91.7	5.23	81.8	102.5
2007	3,729	243.9	4.08	235.9	252.0	364	86.4	4.90	77.0	96.4
2008	3,614	233.5	3.97	225.7	241.4	379	83.6	4.65	74.7	93.1
2009	3,549	222.0	3.82	214.6	229.6	428	92.8	4.84	83.5	102.7
2010	3,712	228.4	3.84	220.9	236.1	464	94.9	4.73	85.9	104.6
2011	3,585	215.8	3.69	208.6	223.1	477	91.4	4.46	82.9	100.5
2012	3,670	216.4	3.66	209.3	223.7	491	88.0	4.23	79.9	96.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

All Sites Combined										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	338,281	256.0	0.45	255.1	256.9	283,508	247.4	0.47	246.4	248.3
1986	12,128	285.6	2.72	280.3	290.9	10,304	273.6	2.82	268.1	279.2
1987	12,064	282.3	2.69	277.1	287.7	10,164	268.6	2.78	263.2	274.1
1988	12,457	290.8	2.72	285.5	296.2	10,478	276.9	2.82	271.4	282.5
1989	12,547	290.3	2.70	285.0	295.6	10,527	275.9	2.79	270.4	281.4
1990	12,716	292.8	2.70	287.6	298.2	10,678	278.6	2.80	273.1	284.1
1991	12,775	290.6	2.67	285.4	295.9	10,794	278.1	2.77	272.7	283.6
1992	12,802	286.8	2.62	281.7	292.0	10,750	273.1	2.71	267.8	278.5
1993	13,069	290.1	2.62	285.0	295.2	10,944	276.5	2.72	271.2	281.9
1994	13,037	285.7	2.57	280.7	290.8	10,987	274.3	2.68	269.1	279.6
1995	12,985	281.9	2.54	277.0	286.9	10,921	270.7	2.65	265.6	276.0
1996	12,735	274.7	2.49	269.9	279.6	10,598	261.4	2.59	256.4	266.6
1997	12,559	267.2	2.43	262.4	272.0	10,512	256.4	2.54	251.5	261.5
1998	12,616	265.3	2.41	260.6	270.0	10,558	255.0	2.52	250.1	260.0
1999	12,697	264.7	2.39	260.0	269.4	10,675	255.7	2.51	250.8	260.6
2000	12,816	263.7	2.36	259.1	268.3	10,711	253.9	2.48	249.0	258.8
2001	12,730	259.3	2.33	254.8	263.9	10,676	250.9	2.45	246.1	255.8
2002	12,577	252.9	2.29	248.5	257.5	10,583	246.2	2.42	241.5	251.0
2003	12,451	247.2	2.25	242.8	251.6	10,379	238.9	2.37	234.3	243.6
2004	12,344	240.9	2.20	236.6	245.2	10,349	234.6	2.33	230.0	239.2
2005	12,241	236.4	2.17	232.2	240.7	10,213	229.3	2.30	224.8	233.8
2006	12,181	231.1	2.13	227.0	235.3	10,181	225.1	2.26	220.7	229.5
2007	12,119	225.9	2.09	221.9	230.1	10,050	218.7	2.21	214.4	223.1
2008	12,331	225.2	2.06	221.2	229.3	10,339	220.7	2.20	216.4	225.0
2009	12,344	220.4	2.02	216.4	224.4	10,348	216.6	2.16	212.3	220.8
2010	12,219	214.5	1.98	210.7	218.4	10,142	209.1	2.11	205.0	213.3
2011	12,163	209.1	1.94	205.3	212.9	10,123	205.0	2.07	200.9	209.1
2012	12,578	211.1	1.93	207.4	214.9	10,524	208.2	2.07	204.2	212.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

All Sites Combined

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	50,842	357.8	1.69	354.5	361.1	3,930	117.6	2.14	113.4	121.9
1986	1,761	407.9	10.48	387.6	428.9	63	119.6	17.97	87.0	158.9
1987	1,844	424.3	10.63	403.7	445.6	56	99.3	16.16	70.2	134.9
1988	1,898	425.8	10.49	405.5	446.8	81	138.5	18.16	105.2	177.7
1989	1,935	435.9	10.64	415.3	457.2	84	112.5	14.15	86.4	143.2
1990	1,949	437.0	10.66	416.4	458.3	89	144.5	18.28	110.9	183.8
1991	1,892	422.2	10.42	402.0	443.0	89	145.0	18.00	111.9	183.5
1992	1,958	431.5	10.52	411.1	452.5	94	132.5	15.89	103.2	166.5
1993	2,033	435.0	10.36	414.9	455.6	92	126.4	15.36	98.1	159.2
1994	1,945	410.8	9.98	391.5	430.7	105	117.0	13.33	92.3	145.5
1995	1,953	402.5	9.76	383.6	422.0	111	123.4	13.90	97.6	153.0
1996	2,024	415.5	9.88	396.3	435.2	113	113.9	12.49	90.7	140.5
1997	1,904	377.6	9.21	359.7	396.0	143	147.5	14.33	120.8	177.7
1998	1,941	374.9	9.05	357.4	393.0	117	110.3	11.93	88.2	135.7
1999	1,891	369.2	9.02	351.7	387.2	131	120.2	12.24	97.4	146.1
2000	1,923	367.3	8.91	350.1	385.2	182	148.8	12.74	124.8	175.4
2001	1,890	357.4	8.76	340.4	374.9	164	121.6	11.08	100.9	144.9
2002	1,818	334.6	8.33	318.5	351.3	176	124.7	10.76	104.5	147.2
2003	1,876	337.9	8.32	321.8	354.5	196	138.5	11.19	117.5	161.8
2004	1,837	327.4	8.16	311.6	343.7	158	102.1	9.22	84.9	121.5
2005	1,851	326.3	8.11	310.6	342.5	177	114.9	9.89	96.4	135.6
2006	1,819	314.0	7.87	298.8	329.7	181	112.7	9.42	95.0	132.4
2007	1,888	320.5	7.86	305.3	336.3	181	99.3	8.27	83.7	116.6
2008	1,808	302.2	7.55	287.5	317.3	184	94.6	7.88	79.8	111.1
2009	1,761	282.1	7.14	268.3	296.4	235	121.9	8.94	105.0	140.4
2010	1,839	290.6	7.19	276.7	305.0	238	114.5	8.30	98.8	131.7
2011	1,810	275.8	6.85	262.6	289.5	230	102.9	7.46	88.8	118.4
2012	1,794	268.8	6.69	255.8	282.2	260	110.7	7.59	96.3	126.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

All Sites Combined										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	320,465	170.7	0.30	170.1	171.3	270,631	167.1	0.32	166.5	167.7
1986	10,894	178.4	1.74	175.1	181.9	9,424	174.0	1.83	170.5	177.7
1987	11,118	180.9	1.74	177.5	184.3	9,688	177.6	1.84	174.1	181.3
1988	11,038	177.7	1.72	174.3	181.1	9,517	173.3	1.81	169.8	176.9
1989	11,385	182.5	1.73	179.1	185.9	9,760	177.3	1.83	173.7	180.9
1990	11,724	185.6	1.74	182.2	189.0	10,084	180.6	1.83	177.0	184.2
1991	11,641	181.5	1.70	178.1	184.8	10,042	177.5	1.80	174.0	181.1
1992	11,820	181.9	1.69	178.6	185.3	10,164	177.8	1.79	174.3	181.3
1993	12,200	185.3	1.69	182.0	188.6	10,421	180.6	1.80	177.1	184.1
1994	12,107	182.2	1.67	179.0	185.6	10,347	177.8	1.77	174.4	181.4
1995	12,116	180.5	1.65	177.2	183.7	10,371	177.0	1.76	173.5	180.5
1996	12,050	178.4	1.64	175.2	181.7	10,218	173.8	1.75	170.4	177.2
1997	11,920	175.2	1.62	172.0	178.4	10,097	170.7	1.72	167.4	174.2
1998	12,022	174.8	1.61	171.7	178.0	10,137	170.0	1.71	166.7	173.4
1999	12,327	177.0	1.61	173.8	180.2	10,410	172.4	1.71	169.1	175.8
2000	12,549	179.3	1.61	176.1	182.5	10,573	175.0	1.73	171.6	178.4
2001	12,048	171.1	1.57	168.0	174.2	10,106	167.1	1.69	163.8	170.4
2002	12,160	171.3	1.57	168.3	174.4	10,119	166.2	1.68	162.9	169.5
2003	12,013	167.3	1.54	164.3	170.3	10,058	163.4	1.65	160.2	166.7
2004	11,945	165.6	1.53	162.6	168.7	9,950	161.5	1.64	158.3	164.7
2005	12,009	165.0	1.52	162.0	168.0	10,013	161.5	1.64	158.3	164.8
2006	11,902	162.1	1.50	159.2	165.1	9,889	158.6	1.62	155.5	161.9
2007	11,994	161.7	1.50	158.8	164.7	9,970	158.7	1.62	155.5	161.9
2008	11,967	158.9	1.47	156.0	161.8	9,966	156.6	1.60	153.5	159.8
2009	11,838	156.3	1.46	153.5	159.2	9,857	154.8	1.59	151.7	157.9
2010	11,851	153.9	1.44	151.1	156.8	9,752	150.9	1.56	147.8	154.0
2011	11,843	151.7	1.42	148.9	154.5	9,821	150.3	1.55	147.3	153.4
2012	11,984	151.5	1.41	148.8	154.3	9,877	149.7	1.55	146.7	152.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

All Sites Combined										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	46,326	213.1	1.00	211.2	215.1	3,503	78.6	1.42	75.9	81.4
1986	1,428	226.9	6.15	215.0	239.2	41	60.1	10.73	40.9	84.2
1987	1,387	217.2	5.95	205.7	229.2	42	61.6	10.58	42.6	85.2
1988	1,470	222.1	5.91	210.6	233.9	51	62.7	9.97	44.7	84.9
1989	1,556	233.3	6.03	221.6	245.4	68	97.4	13.31	73.0	126.2
1990	1,590	236.6	6.04	224.9	248.7	49	64.9	10.67	45.7	88.5
1991	1,524	222.0	5.78	210.8	233.6	74	74.5	9.75	56.6	95.7
1992	1,578	222.0	5.67	211.0	233.4	78	79.4	10.06	60.9	101.2
1993	1,706	236.6	5.81	225.4	248.3	73	69.9	9.07	53.3	89.6
1994	1,661	226.9	5.63	215.9	238.1	99	95.3	10.63	75.6	117.9
1995	1,646	221.0	5.51	210.3	232.0	99	84.7	9.36	67.3	104.7
1996	1,729	227.3	5.53	216.6	238.4	103	83.9	8.95	67.2	103.0
1997	1,731	224.0	5.44	213.5	234.9	92	67.2	7.64	53.0	83.6
1998	1,753	222.0	5.35	211.6	232.7	132	89.2	8.52	73.3	107.3
1999	1,790	223.5	5.33	213.2	234.2	127	76.8	7.34	63.1	92.4
2000	1,841	225.3	5.29	215.0	235.8	135	82.8	7.65	68.5	99.0
2001	1,807	219.6	5.21	209.5	230.1	135	75.2	6.90	62.2	89.8
2002	1,895	225.9	5.23	215.8	236.4	146	79.7	7.00	66.6	94.5
2003	1,794	210.9	5.02	201.2	221.0	161	84.0	7.03	70.8	98.8
2004	1,835	211.0	4.97	201.4	221.0	160	79.5	6.64	67.0	93.5
2005	1,810	205.0	4.86	195.6	214.8	186	87.0	6.76	74.3	101.1
2006	1,836	204.6	4.82	195.3	214.3	177	78.2	6.19	66.6	91.2
2007	1,841	200.5	4.72	191.3	209.9	183	76.7	6.01	65.4	89.3
2008	1,806	193.6	4.60	184.7	202.8	195	76.1	5.77	65.3	88.2
2009	1,788	187.2	4.48	178.5	196.2	193	72.9	5.53	62.5	84.5
2010	1,873	192.7	4.51	183.9	201.7	226	81.3	5.68	70.5	93.1
2011	1,775	179.0	4.31	170.7	187.7	247	83.4	5.55	72.9	95.0
2012	1,876	185.5	4.35	177.0	194.2	231	72.5	5.00	63.0	82.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Oral Cavity and Pharynx

Both Sexes

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	9,379	2.9	0.03	2.8	2.9	7,381	2.6	0.03	2.6	2.7
1986	384	3.6	0.19	3.3	4.0	307	3.2	0.19	2.9	3.6
1987	384	3.6	0.19	3.3	4.0	314	3.3	0.19	3.0	3.7
1988	378	3.6	0.19	3.2	3.9	298	3.2	0.18	2.8	3.5
1989	380	3.6	0.18	3.2	3.9	286	3.0	0.18	2.7	3.4
1990	427	4.0	0.20	3.6	4.4	325	3.5	0.19	3.1	3.9
1991	421	3.9	0.19	3.5	4.2	315	3.3	0.19	2.9	3.6
1992	365	3.3	0.18	3.0	3.7	281	2.9	0.18	2.6	3.3
1993	363	3.3	0.17	2.9	3.6	282	2.9	0.17	2.6	3.3
1994	354	3.1	0.17	2.8	3.5	271	2.8	0.17	2.4	3.1
1995	370	3.2	0.17	2.9	3.6	284	2.8	0.17	2.5	3.2
1996	368	3.2	0.17	2.9	3.6	307	3.1	0.18	2.8	3.5
1997	339	2.9	0.16	2.6	3.3	257	2.6	0.16	2.3	2.9
1998	341	2.9	0.16	2.6	3.2	268	2.7	0.16	2.3	3.0
1999	301	2.5	0.15	2.3	2.8	231	2.3	0.15	2.0	2.6
2000	329	2.8	0.15	2.5	3.1	258	2.5	0.16	2.2	2.8
2001	302	2.5	0.14	2.2	2.8	233	2.3	0.15	2.0	2.6
2002	310	2.5	0.14	2.3	2.8	242	2.3	0.15	2.0	2.6
2003	334	2.7	0.15	2.4	3.0	253	2.4	0.15	2.1	2.7
2004	336	2.7	0.15	2.4	3.0	267	2.5	0.15	2.2	2.8
2005	294	2.3	0.14	2.1	2.6	227	2.1	0.14	1.8	2.4
2006	309	2.4	0.14	2.1	2.7	250	2.3	0.15	2.0	2.6
2007	319	2.4	0.14	2.2	2.7	261	2.4	0.15	2.1	2.7
2008	358	2.7	0.14	2.4	3.0	295	2.6	0.15	2.3	2.9
2009	299	2.2	0.13	2.0	2.5	252	2.2	0.14	1.9	2.5
2010	303	2.2	0.13	2.0	2.5	247	2.2	0.14	1.9	2.4
2011	344	2.4	0.13	2.2	2.7	273	2.3	0.14	2.1	2.6
2012	367	2.5	0.13	2.3	2.8	297	2.5	0.15	2.2	2.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Oral Cavity and Pharynx

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,829	4.6	0.11	4.4	4.8	169	1.9	0.16	1.6	2.2
1986	75	6.7	0.80	5.2	8.4	*	*	*	*	*
1987	68	6.0	0.74	4.6	7.6	*	*	*	*	*
1988	76	6.3	0.74	4.9	7.9	*	*	*	*	*
1989	90	7.3	0.77	5.8	8.9	*	*	*	*	*
1990	101	8.1	0.82	6.6	9.9	*	*	*	*	*
1991	100	8.0	0.81	6.4	9.7	*	*	*	*	*
1992	78	5.9	0.68	4.6	7.4	*	*	*	*	*
1993	77	5.8	0.67	4.6	7.2	*	*	*	*	*
1994	80	5.9	0.66	4.6	7.3	*	*	*	*	*
1995	82	6.2	0.70	4.9	7.7	*	*	*	*	*
1996	56	4.2	0.57	3.2	5.5	*	*	*	*	*
1997	74	5.2	0.61	4.0	6.5	*	*	*	*	*
1998	67	4.8	0.59	3.7	6.1	*	*	*	*	*
1999	63	4.4	0.57	3.4	5.7	*	*	*	*	*
2000	66	4.4	0.55	3.4	5.6	*	*	*	*	*
2001	60	4.0	0.53	3.1	5.2	*	*	*	*	*
2002	63	4.0	0.51	3.1	5.1	*	*	*	*	*
2003	74	4.7	0.56	3.7	6.0	*	*	*	*	*
2004	64	4.0	0.50	3.1	5.1	*	*	*	*	*
2005	57	3.6	0.48	2.7	4.6	10	2.1	0.71	0.9	4.0
2006	53	3.2	0.45	2.4	4.2	*	*	*	*	*
2007	48	2.9	0.42	2.1	3.8	10	2.5	0.86	1.1	4.7
2008	54	3.2	0.44	2.4	4.1	*	*	*	*	*
2009	42	2.3	0.37	1.7	3.2	*	*	*	*	*
2010	47	2.6	0.39	1.9	3.5	*	*	*	*	*
2011	57	3.0	0.41	2.3	3.9	14	2.6	0.75	1.3	4.4
2012	57	3.0	0.40	2.3	3.9	13	1.9	0.54	1.0	3.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Oral Cavity and Pharynx

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	6,397	4.5	0.06	4.4	4.6	4,918	4.1	0.06	4.0	4.2
1986	262	5.8	0.37	5.1	6.6	207	5.2	0.37	4.5	5.9
1987	256	5.7	0.37	5.0	6.5	209	5.3	0.38	4.6	6.1
1988	273	6.0	0.38	5.3	6.8	212	5.3	0.38	4.6	6.1
1989	251	5.4	0.35	4.8	6.2	184	4.5	0.35	3.9	5.3
1990	300	6.6	0.40	5.9	7.5	221	5.7	0.39	4.9	6.5
1991	282	6.1	0.37	5.4	6.9	203	5.0	0.36	4.3	5.8
1992	237	5.1	0.34	4.4	5.8	179	4.5	0.34	3.8	5.2
1993	261	5.4	0.34	4.8	6.1	199	4.8	0.34	4.1	5.5
1994	243	5.0	0.33	4.4	5.7	175	4.3	0.33	3.6	4.9
1995	248	5.1	0.33	4.4	5.8	187	4.4	0.33	3.8	5.1
1996	261	5.4	0.34	4.7	6.1	213	5.1	0.35	4.4	5.8
1997	236	4.7	0.31	4.1	5.3	176	4.1	0.31	3.5	4.8
1998	242	4.9	0.32	4.3	5.5	187	4.4	0.32	3.8	5.1
1999	216	4.2	0.29	3.6	4.8	167	3.8	0.30	3.2	4.4
2000	210	4.0	0.28	3.5	4.6	156	3.5	0.28	3.0	4.1
2001	222	4.2	0.28	3.7	4.8	171	3.8	0.29	3.2	4.4
2002	202	3.8	0.27	3.2	4.3	156	3.4	0.28	2.9	4.0
2003	233	4.3	0.29	3.8	4.9	166	3.7	0.29	3.1	4.3
2004	239	4.3	0.28	3.8	4.9	184	3.9	0.29	3.4	4.5
2005	194	3.5	0.25	3.0	4.0	137	2.9	0.25	2.4	3.4
2006	187	3.3	0.24	2.8	3.8	149	3.1	0.26	2.6	3.6
2007	215	3.7	0.26	3.2	4.2	171	3.5	0.27	3.0	4.0
2008	233	3.9	0.26	3.4	4.5	193	3.9	0.28	3.4	4.5
2009	195	3.1	0.23	2.7	3.6	158	3.0	0.24	2.6	3.5
2010	205	3.5	0.25	3.0	4.0	165	3.3	0.26	2.8	3.9
2011	240	3.8	0.25	3.3	4.3	190	3.6	0.27	3.1	4.2
2012	254	3.9	0.25	3.4	4.4	203	3.8	0.27	3.2	4.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Oral Cavity and Pharynx

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,362	8.0	0.22	7.6	8.4	117	2.7	0.28	2.2	3.3
1986	54	11.2	1.60	8.3	14.7	*	*	*	*	*
1987	46	9.4	1.44	6.8	12.6	*	*	*	*	*
1988	58	11.3	1.54	8.5	14.8	*	*	*	*	*
1989	63	11.8	1.51	9.1	15.2	*	*	*	*	*
1990	78	14.9	1.76	11.6	18.7	*	*	*	*	*
1991	73	14.0	1.73	10.8	17.8	*	*	*	*	*
1992	54	9.6	1.35	7.2	12.6	*	*	*	*	*
1993	59	10.6	1.44	8.0	13.8	*	*	*	*	*
1994	66	11.2	1.38	8.6	14.2	*	*	*	*	*
1995	57	10.6	1.48	7.9	13.8	*	*	*	*	*
1996	43	7.3	1.14	5.3	9.9	*	*	*	*	*
1997	56	9.3	1.31	6.9	12.2	*	*	*	*	*
1998	52	9.1	1.33	6.7	12.1	*	*	*	*	*
1999	44	7.3	1.14	5.3	9.9	*	*	*	*	*
2000	50	7.8	1.15	5.7	10.4	*	*	*	*	*
2001	45	7.2	1.11	5.2	9.7	*	*	*	*	*
2002	43	6.2	0.96	4.5	8.4	*	*	*	*	*
2003	61	9.2	1.24	7.0	12.0	*	*	*	*	*
2004	51	7.3	1.07	5.4	9.7	*	*	*	*	*
2005	50	7.2	1.05	5.3	9.6	*	*	*	*	*
2006	35	5.0	0.87	3.4	7.0	*	*	*	*	*
2007	39	5.6	0.94	3.9	7.7	*	*	*	*	*
2008	35	4.5	0.80	3.1	6.4	*	*	*	*	*
2009	33	4.2	0.76	2.8	5.9	*	*	*	*	*
2010	33	4.4	0.79	3.0	6.2	*	*	*	*	*
2011	41	4.9	0.79	3.5	6.7	*	*	*	*	*
2012	43	5.4	0.85	3.8	7.3	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Oral Cavity and Pharynx

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,982	1.6	0.03	1.5	1.7	2,463	1.5	0.03	1.5	1.6
1986	122	2.0	0.18	1.6	2.4	100	1.8	0.18	1.4	2.1
1987	128	2.1	0.19	1.8	2.5	105	1.9	0.19	1.6	2.4
1988	105	1.8	0.17	1.4	2.1	86	1.6	0.18	1.3	2.0
1989	129	2.1	0.19	1.8	2.5	102	1.9	0.19	1.5	2.3
1990	127	2.0	0.18	1.7	2.4	104	1.8	0.18	1.5	2.2
1991	139	2.2	0.19	1.9	2.6	112	2.0	0.20	1.7	2.5
1992	128	2.0	0.18	1.7	2.4	102	1.8	0.19	1.5	2.2
1993	102	1.6	0.16	1.3	1.9	83	1.4	0.16	1.1	1.8
1994	111	1.6	0.16	1.4	2.0	96	1.6	0.16	1.3	2.0
1995	122	1.8	0.16	1.5	2.2	97	1.6	0.17	1.3	2.0
1996	107	1.6	0.16	1.3	1.9	94	1.6	0.17	1.3	2.0
1997	103	1.5	0.15	1.3	1.9	81	1.4	0.15	1.1	1.7
1998	99	1.5	0.15	1.2	1.8	81	1.4	0.15	1.1	1.7
1999	85	1.2	0.13	1.0	1.5	64	1.0	0.13	0.8	1.3
2000	119	1.7	0.16	1.4	2.0	102	1.7	0.17	1.4	2.1
2001	80	1.1	0.13	0.9	1.4	62	1.0	0.13	0.8	1.3
2002	108	1.5	0.15	1.3	1.9	86	1.4	0.15	1.1	1.7
2003	101	1.4	0.14	1.1	1.7	87	1.4	0.15	1.1	1.7
2004	97	1.3	0.14	1.1	1.6	83	1.3	0.15	1.1	1.7
2005	100	1.4	0.14	1.1	1.7	90	1.4	0.16	1.2	1.8
2006	122	1.7	0.15	1.4	2.0	101	1.6	0.16	1.3	2.0
2007	104	1.4	0.14	1.2	1.7	90	1.5	0.16	1.2	1.8
2008	125	1.7	0.15	1.4	2.0	102	1.6	0.16	1.3	2.0
2009	104	1.4	0.14	1.1	1.7	94	1.4	0.15	1.2	1.8
2010	98	1.3	0.13	1.0	1.5	82	1.2	0.14	1.0	1.6
2011	104	1.4	0.14	1.1	1.7	83	1.3	0.15	1.0	1.6
2012	113	1.4	0.14	1.2	1.7	94	1.5	0.15	1.2	1.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Oral Cavity and Pharynx

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	467	2.1	0.10	1.9	2.2	52	1.2	0.18	0.9	1.6
1986	21	3.4	0.76	2.1	5.2	*	*	*	*	*
1987	22	3.4	0.75	2.1	5.2	*	*	*	*	*
1988	18	2.6	0.62	1.5	4.1	*	*	*	*	*
1989	27	3.8	0.73	2.5	5.5	*	*	*	*	*
1990	23	3.2	0.68	2.0	4.8	*	*	*	*	*
1991	27	3.7	0.72	2.4	5.4	*	*	*	*	*
1992	24	3.1	0.65	2.0	4.7	*	*	*	*	*
1993	18	2.4	0.56	1.4	3.7	*	*	*	*	*
1994	14	1.8	0.49	1.0	3.1	*	*	*	*	*
1995	25	3.2	0.65	2.1	4.8	*	*	*	*	*
1996	13	1.8	0.50	0.9	3.0	*	*	*	*	*
1997	18	2.2	0.51	1.3	3.4	*	*	*	*	*
1998	15	1.8	0.47	1.0	3.0	*	*	*	*	*
1999	19	2.3	0.54	1.4	3.6	*	*	*	*	*
2000	16	1.9	0.49	1.1	3.1	*	*	*	*	*
2001	15	1.8	0.46	1.0	2.9	*	*	*	*	*
2002	20	2.3	0.51	1.4	3.5	*	*	*	*	*
2003	13	1.5	0.42	0.8	2.6	*	*	*	*	*
2004	13	1.5	0.42	0.8	2.6	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	18	1.9	0.45	1.1	3.0	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	19	2.1	0.47	1.2	3.2	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	14	1.3	0.36	0.7	2.2	*	*	*	*	*
2011	16	1.6	0.40	0.9	2.6	*	*	*	*	*
2012	14	1.3	0.36	0.7	2.2	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Digestive System										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	163,836	50.7	0.13	50.4	50.9	135,198	48.1	0.13	47.9	48.4
1986	6,022	57.7	0.75	56.3	59.2	5,113	54.9	0.78	53.3	56.4
1987	6,144	58.5	0.76	57.1	60.0	5,221	55.8	0.78	54.3	57.4
1988	6,026	56.8	0.74	55.4	58.3	5,037	53.4	0.76	51.9	54.9
1989	5,959	55.9	0.73	54.4	57.3	4,997	52.7	0.75	51.2	54.2
1990	6,179	57.3	0.73	55.9	58.8	5,158	53.8	0.76	52.4	55.3
1991	6,059	55.5	0.72	54.1	56.9	5,096	52.6	0.74	51.1	54.0
1992	6,049	54.5	0.70	53.2	55.9	5,078	51.8	0.73	50.4	53.2
1993	6,114	54.5	0.70	53.1	55.9	5,059	51.1	0.72	49.7	52.5
1994	6,165	54.4	0.69	53.0	55.8	5,168	51.7	0.72	50.3	53.1
1995	6,111	53.3	0.68	52.0	54.6	5,140	51.0	0.71	49.6	52.4
1996	5,981	51.9	0.67	50.6	53.2	4,913	48.6	0.69	47.2	50.0
1997	5,995	51.5	0.67	50.2	52.8	4,955	48.6	0.69	47.3	50.0
1998	6,115	51.9	0.66	50.6	53.2	5,068	49.3	0.69	48.0	50.7
1999	6,108	51.3	0.66	50.0	52.6	5,073	48.9	0.69	47.6	50.3
2000	6,258	52.1	0.66	50.8	53.4	5,157	49.4	0.69	48.1	50.8
2001	6,173	51.0	0.65	49.7	52.3	5,075	48.3	0.68	47.0	49.7
2002	6,105	50.0	0.64	48.7	51.2	5,016	47.5	0.67	46.2	48.8
2003	5,999	48.5	0.63	47.3	49.7	4,881	45.7	0.66	44.4	47.0
2004	6,001	48.1	0.62	46.9	49.3	4,898	45.6	0.65	44.4	46.9
2005	5,980	47.3	0.61	46.1	48.5	4,890	45.1	0.65	43.8	46.4
2006	6,010	47.0	0.61	45.8	48.2	4,883	44.6	0.64	43.3	45.9
2007	5,945	45.8	0.60	44.6	47.0	4,831	43.5	0.63	42.3	44.8
2008	6,086	46.1	0.60	44.9	47.3	4,928	43.9	0.63	42.7	45.1
2009	6,022	44.9	0.58	43.7	46.0	4,903	43.1	0.62	41.9	44.4
2010	6,094	44.7	0.58	43.6	45.9	4,932	42.8	0.62	41.6	44.0
2011	6,022	43.6	0.57	42.4	44.7	4,822	41.4	0.61	40.2	42.6
2012	6,114	43.4	0.56	42.3	44.5	4,906	41.5	0.60	40.3	42.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Digestive System										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	25,931	72.6	0.46	71.7	73.5	2,705	35.3	0.74	33.8	36.8
1986	863	84.6	3.00	78.8	90.6	45	33.0	5.62	22.9	45.6
1987	887	84.1	2.92	78.5	90.0	36	27.3	5.26	18.0	39.2
1988	942	86.3	2.91	80.7	92.2	47	34.8	5.90	24.2	48.0
1989	899	83.5	2.88	78.0	89.3	62	47.7	7.04	34.9	63.0
1990	960	87.3	2.91	81.7	93.2	61	44.7	6.76	32.4	59.4
1991	904	81.3	2.78	75.9	86.9	59	44.0	6.48	32.3	58.2
1992	897	77.9	2.68	72.7	83.3	74	43.3	5.69	32.9	55.6
1993	987	85.1	2.78	79.7	90.7	68	41.0	5.56	30.8	53.0
1994	911	75.9	2.58	70.9	81.1	86	46.2	5.63	35.9	58.3
1995	887	73.6	2.53	68.7	78.7	84	45.9	5.54	35.7	57.8
1996	987	79.7	2.60	74.7	85.0	81	38.6	4.79	29.8	48.9
1997	945	75.4	2.51	70.5	80.4	95	42.0	4.82	33.1	52.3
1998	969	74.9	2.46	70.2	79.8	78	31.9	4.06	24.5	40.7
1999	927	71.8	2.41	67.2	76.7	108	41.3	4.39	33.1	50.6
2000	986	73.7	2.39	69.0	78.5	115	41.8	4.29	33.8	50.9
2001	985	72.4	2.36	67.9	77.2	113	36.2	3.73	29.3	44.2
2002	981	71.4	2.33	66.9	76.1	108	35.9	3.72	29.0	43.8
2003	979	70.2	2.29	65.8	74.8	139	42.8	3.95	35.4	51.1
2004	1,008	70.4	2.27	66.0	75.0	95	27.1	2.98	21.6	33.5
2005	976	67.1	2.19	62.8	71.5	114	32.5	3.28	26.3	39.4
2006	1,000	67.5	2.18	63.3	71.9	127	32.2	3.07	26.5	38.7
2007	990	64.9	2.11	60.9	69.2	124	28.5	2.77	23.3	34.3
2008	1,043	68.1	2.15	63.9	72.4	115	25.2	2.53	20.5	30.5
2009	955	59.5	1.97	55.7	63.5	164	36.3	3.05	30.6	42.7
2010	999	61.5	1.99	57.6	65.5	163	34.3	2.88	28.9	40.3
2011	1,028	61.7	1.97	57.9	65.7	172	33.1	2.70	28.0	38.8
2012	1,036	60.8	1.94	57.1	64.7	172	30.2	2.45	25.6	35.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Digestive System										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	87,251	65.6	0.23	65.2	66.1	71,877	62.5	0.24	62.0	62.9
1986	3,177	75.8	1.41	73.1	78.6	2,685	72.3	1.46	69.5	75.2
1987	3,169	75.5	1.40	72.7	78.3	2,654	71.6	1.45	68.8	74.5
1988	3,208	75.9	1.40	73.1	78.6	2,645	71.1	1.44	68.3	74.0
1989	3,155	74.8	1.39	72.1	77.5	2,638	70.8	1.43	68.0	73.7
1990	3,202	74.2	1.36	71.6	76.9	2,660	69.9	1.40	67.2	72.7
1991	3,225	73.8	1.35	71.2	76.5	2,707	70.2	1.40	67.5	73.0
1992	3,204	72.2	1.32	69.6	74.8	2,662	68.1	1.36	65.4	70.8
1993	3,148	70.2	1.29	67.7	72.8	2,591	65.6	1.32	63.1	68.3
1994	3,235	71.2	1.29	68.7	73.7	2,697	67.6	1.34	65.1	70.3
1995	3,227	70.3	1.27	67.8	72.8	2,689	66.8	1.32	64.2	69.4
1996	3,159	68.4	1.25	66.0	70.9	2,595	64.5	1.29	62.0	67.0
1997	3,180	67.8	1.23	65.4	70.3	2,605	63.9	1.27	61.4	66.4
1998	3,235	67.4	1.21	65.1	69.9	2,682	64.4	1.26	62.0	66.9
1999	3,191	66.2	1.19	63.9	68.6	2,660	63.4	1.24	61.0	65.9
2000	3,253	66.4	1.18	64.1	68.8	2,672	63.0	1.23	60.6	65.5
2001	3,247	65.5	1.17	63.2	67.8	2,679	62.6	1.22	60.2	65.0
2002	3,235	64.5	1.15	62.3	66.8	2,684	62.0	1.21	59.7	64.4
2003	3,309	65.4	1.15	63.1	67.7	2,684	61.7	1.20	59.4	64.1
2004	3,203	61.8	1.11	59.6	64.0	2,616	58.9	1.16	56.6	61.2
2005	3,216	61.1	1.10	59.0	63.3	2,631	58.3	1.15	56.1	60.6
2006	3,163	59.1	1.07	57.0	61.2	2,592	56.6	1.13	54.4	58.8
2007	3,143	57.7	1.05	55.7	59.8	2,546	54.8	1.10	52.7	57.0
2008	3,322	59.0	1.04	57.0	61.1	2,691	56.0	1.10	53.9	58.2
2009	3,360	58.6	1.03	56.6	60.7	2,745	56.4	1.09	54.3	58.6
2010	3,349	57.2	1.01	55.3	59.3	2,720	54.7	1.07	52.6	56.8
2011	3,287	54.9	0.98	53.0	56.9	2,661	52.6	1.04	50.6	54.7
2012	3,449	56.3	0.98	54.4	58.2	2,786	53.7	1.04	51.7	55.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Digestive System										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	13,800	95.2	0.86	93.5	96.9	1,573	45.7	1.31	43.2	48.4
1986	465	109.3	5.48	98.8	120.5	27	38.7	8.31	24.1	58.4
1987	491	112.0	5.39	101.7	123.0	24	33.2	7.65	19.9	51.5
1988	530	117.6	5.47	107.2	128.8	33	54.3	11.15	34.7	79.7
1989	482	111.2	5.46	100.7	122.3	34	43.6	8.48	28.6	63.2
1990	496	109.6	5.27	99.6	120.4	46	77.1	13.90	52.3	107.9
1991	481	105.6	5.15	95.7	116.1	37	62.9	11.82	41.9	89.3
1992	496	110.7	5.34	100.4	121.6	46	60.2	10.12	42.0	82.7
1993	516	111.7	5.28	101.6	122.5	41	62.6	11.16	42.7	87.3
1994	491	102.8	4.96	93.3	112.9	47	53.7	9.15	37.3	74.0
1995	490	102.9	4.98	93.4	113.0	48	59.5	9.94	41.7	81.5
1996	520	103.2	4.85	93.9	113.0	44	45.0	8.05	30.6	63.0
1997	509	98.7	4.63	89.8	108.1	66	67.7	9.81	49.9	89.1
1998	516	97.7	4.55	89.0	107.0	37	32.9	6.10	22.0	46.7
1999	476	92.9	4.52	84.3	102.1	55	55.9	8.77	40.1	75.1
2000	518	96.0	4.46	87.5	105.1	63	51.3	7.60	37.5	68.0
2001	496	89.9	4.31	81.6	98.6	72	50.7	6.88	38.1	65.7
2002	485	88.1	4.23	80.0	96.7	66	51.2	7.01	38.3	66.4
2003	542	94.8	4.33	86.5	103.6	83	56.6	7.22	43.3	72.2
2004	533	90.5	4.18	82.5	99.0	54	31.2	4.71	22.7	41.7
2005	525	89.0	4.15	81.0	97.4	60	36.9	5.42	27.0	48.8
2006	502	83.5	3.95	75.9	91.5	69	41.6	5.55	31.4	53.6
2007	530	86.3	3.99	78.7	94.4	67	31.7	4.35	23.7	41.2
2008	561	91.3	4.09	83.5	99.7	70	35.0	4.71	26.4	45.3
2009	522	79.0	3.67	72.0	86.5	93	47.0	5.41	37.0	58.6
2010	540	81.5	3.72	74.4	89.1	89	41.1	4.86	32.1	51.5
2011	535	78.8	3.60	71.9	86.1	91	39.5	4.61	31.0	49.4
2012	552	80.1	3.59	73.2	87.4	111	44.3	4.63	35.7	54.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Digestive System										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	76,585	39.9	0.15	39.6	40.1	63,321	37.8	0.15	37.5	38.1
1986	2,845	45.6	0.87	44.0	47.4	2,428	43.3	0.89	41.6	45.1
1987	2,975	47.1	0.87	45.4	48.9	2,567	45.4	0.91	43.7	47.2
1988	2,818	44.2	0.84	42.5	45.8	2,392	41.8	0.87	40.1	43.5
1989	2,804	43.5	0.83	41.9	45.1	2,359	40.9	0.85	39.2	42.6
1990	2,977	45.7	0.85	44.1	47.4	2,498	42.9	0.87	41.2	44.7
1991	2,834	42.9	0.81	41.4	44.6	2,389	40.4	0.84	38.8	42.1
1992	2,845	42.4	0.80	40.9	44.0	2,416	40.4	0.83	38.8	42.1
1993	2,966	43.8	0.81	42.3	45.5	2,468	41.2	0.84	39.6	42.9
1994	2,930	42.8	0.80	41.2	44.4	2,471	40.6	0.83	39.0	42.2
1995	2,884	41.6	0.78	40.1	43.2	2,451	40.0	0.82	38.4	41.6
1996	2,822	40.6	0.77	39.1	42.2	2,318	37.8	0.80	36.3	39.4
1997	2,815	40.1	0.76	38.6	41.6	2,350	38.1	0.80	36.5	39.7
1998	2,880	40.7	0.77	39.3	42.3	2,386	38.5	0.80	37.0	40.1
1999	2,917	40.8	0.76	39.3	42.3	2,413	38.5	0.79	36.9	40.1
2000	3,005	41.7	0.77	40.2	43.2	2,485	39.4	0.80	37.9	41.0
2001	2,926	40.4	0.75	38.9	41.9	2,396	37.9	0.79	36.4	39.5
2002	2,870	39.3	0.74	37.9	40.8	2,332	36.7	0.77	35.2	38.2
2003	2,690	36.5	0.71	35.1	37.9	2,197	34.4	0.75	33.0	35.9
2004	2,798	37.8	0.72	36.4	39.3	2,282	35.6	0.76	34.2	37.1
2005	2,764	37.1	0.71	35.7	38.5	2,259	35.2	0.76	33.8	36.8
2006	2,847	37.9	0.72	36.5	39.4	2,291	35.5	0.76	34.0	37.0
2007	2,802	36.8	0.71	35.4	38.2	2,285	34.9	0.75	33.5	36.4
2008	2,764	36.0	0.70	34.7	37.4	2,237	34.2	0.74	32.8	35.7
2009	2,662	34.5	0.68	33.2	35.8	2,158	33.0	0.73	31.6	34.4
2010	2,745	35.1	0.68	33.8	36.5	2,212	33.5	0.73	32.1	35.0
2011	2,735	34.5	0.67	33.2	35.9	2,161	32.3	0.71	30.9	33.7
2012	2,665	33.2	0.66	31.9	34.5	2,120	31.5	0.70	30.2	32.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Digestive System										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	12,131	57.3	0.53	56.3	58.4	1,132	27.2	0.86	25.6	28.9
1986	398	67.6	3.46	60.9	74.6	18	27.6	7.38	15.1	45.3
1987	396	64.5	3.31	58.2	71.3	12	21.6	6.94	10.2	38.5
1988	412	64.7	3.25	58.5	71.4	14	19.9	6.16	9.8	35.0
1989	417	65.3	3.26	59.1	72.0	28	49.1	10.29	31.0	72.3
1990	464	71.6	3.38	65.1	78.5	15	20.2	6.03	10.2	34.7
1991	423	64.6	3.18	58.5	71.1	22	30.1	7.23	17.6	46.8
1992	401	57.8	2.93	52.2	63.8	28	30.5	6.49	19.1	45.4
1993	471	67.8	3.16	61.7	74.2	27	26.0	5.50	16.4	38.7
1994	420	58.7	2.90	53.2	64.6	39	39.9	7.04	27.4	55.7
1995	397	55.1	2.79	49.8	60.8	36	35.9	6.43	24.4	50.2
1996	467	63.7	2.98	58.0	69.8	37	33.3	5.88	22.8	46.4
1997	436	58.6	2.83	53.2	64.4	29	23.6	4.75	15.2	34.4
1998	453	58.9	2.79	53.6	64.6	41	30.5	5.24	21.1	42.2
1999	451	58.1	2.75	52.8	63.7	53	32.7	4.79	24.0	43.3
2000	468	58.6	2.73	53.4	64.2	52	34.7	5.10	25.4	45.9
2001	489	60.6	2.76	55.3	66.2	41	24.8	4.13	17.4	34.1
2002	496	60.2	2.72	55.0	65.8	42	24.9	4.04	17.6	33.9
2003	437	53.0	2.55	48.1	58.2	56	32.2	4.52	24.0	42.2
2004	475	56.3	2.60	51.3	61.6	41	22.9	3.72	16.2	31.2
2005	451	52.6	2.50	47.8	57.7	54	28.5	4.05	21.1	37.4
2006	498	56.4	2.55	51.6	61.7	58	25.6	3.54	19.2	33.4
2007	460	50.9	2.40	46.3	55.8	57	25.2	3.51	18.8	32.9
2008	482	52.6	2.42	48.0	57.6	45	17.8	2.80	12.8	24.1
2009	433	46.0	2.24	41.8	50.6	71	28.3	3.52	21.8	35.9
2010	459	47.8	2.26	43.4	52.4	74	28.5	3.47	22.1	36.0
2011	493	50.0	2.28	45.7	54.7	81	28.5	3.30	22.4	35.6
2012	484	47.8	2.21	43.6	52.3	61	19.2	2.58	14.5	24.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Esophagus										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	15,058	4.6	0.04	4.6	4.7	12,027	4.3	0.04	4.2	4.4
1986	479	4.4	0.21	4.0	4.9	330	3.4	0.19	3.1	3.8
1987	510	4.8	0.22	4.4	5.2	378	4.0	0.21	3.6	4.4
1988	483	4.5	0.21	4.1	4.9	341	3.5	0.19	3.2	3.9
1989	493	4.6	0.21	4.2	5.1	367	3.9	0.21	3.5	4.3
1990	479	4.4	0.20	4.0	4.8	340	3.5	0.19	3.2	3.9
1991	527	4.8	0.21	4.4	5.2	384	3.9	0.20	3.6	4.4
1992	535	4.8	0.21	4.4	5.2	419	4.3	0.21	3.9	4.7
1993	510	4.5	0.20	4.1	4.9	383	3.8	0.20	3.5	4.3
1994	552	4.9	0.21	4.5	5.3	417	4.2	0.21	3.8	4.6
1995	557	4.9	0.21	4.5	5.3	433	4.3	0.21	3.9	4.8
1996	554	4.8	0.20	4.4	5.2	432	4.3	0.21	3.9	4.8
1997	551	4.8	0.20	4.4	5.2	435	4.3	0.21	3.9	4.8
1998	576	4.9	0.20	4.5	5.3	475	4.7	0.22	4.3	5.1
1999	515	4.3	0.19	4.0	4.7	416	4.1	0.20	3.7	4.5
2000	592	5.0	0.20	4.6	5.4	479	4.7	0.21	4.2	5.1
2001	589	4.9	0.20	4.5	5.3	466	4.5	0.21	4.1	4.9
2002	564	4.6	0.19	4.3	5.0	469	4.5	0.21	4.1	4.9
2003	562	4.5	0.19	4.2	4.9	468	4.4	0.21	4.0	4.9
2004	561	4.5	0.19	4.2	4.9	470	4.5	0.21	4.1	4.9
2005	607	4.8	0.20	4.4	5.2	514	4.8	0.21	4.4	5.2
2006	568	4.5	0.19	4.1	4.8	482	4.5	0.20	4.1	4.9
2007	618	4.7	0.19	4.4	5.1	528	4.8	0.21	4.4	5.2
2008	633	4.7	0.19	4.4	5.1	535	4.7	0.21	4.3	5.2
2009	602	4.4	0.18	4.1	4.8	514	4.5	0.20	4.1	4.9
2010	622	4.5	0.18	4.2	4.9	537	4.7	0.20	4.3	5.1
2011	588	4.2	0.18	3.9	4.6	484	4.2	0.19	3.8	4.6
2012	631	4.5	0.18	4.1	4.8	531	4.5	0.20	4.1	4.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Esophagus										
Both Sexes										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,908	7.5	0.14	7.3	7.8	123	1.6	0.15	1.3	1.9
1986	149	12.9	1.08	10.9	15.2	*	*	*	*	*
1987	131	11.6	1.04	9.7	13.8	*	*	*	*	*
1988	141	11.9	1.02	10.0	14.0	*	*	*	*	*
1989	125	10.6	0.97	8.8	12.7	*	*	*	*	*
1990	134	11.3	1.00	9.5	13.5	*	*	*	*	*
1991	139	11.3	0.98	9.5	13.4	*	*	*	*	*
1992	110	8.8	0.85	7.2	10.6	*	*	*	*	*
1993	124	9.9	0.91	8.2	11.8	*	*	*	*	*
1994	132	10.2	0.91	8.5	12.2	*	*	*	*	*
1995	119	9.1	0.85	7.5	10.9	*	*	*	*	*
1996	116	8.5	0.81	7.0	10.3	*	*	*	*	*
1997	110	8.1	0.79	6.7	9.8	*	*	*	*	*
1998	101	7.2	0.72	5.8	8.7	*	*	*	*	*
1999	95	6.9	0.71	5.5	8.4	*	*	*	*	*
2000	111	7.8	0.75	6.4	9.4	*	*	*	*	*
2001	117	8.2	0.77	6.7	9.8	*	*	*	*	*
2002	88	6.1	0.66	4.8	7.5	*	*	*	*	*
2003	89	5.7	0.62	4.6	7.1	*	*	*	*	*
2004	89	5.9	0.64	4.7	7.3	*	*	*	*	*
2005	91	5.8	0.61	4.6	7.1	*	*	*	*	*
2006	79	5.1	0.59	4.0	6.4	*	*	*	*	*
2007	82	5.1	0.58	4.1	6.4	*	*	*	*	*
2008	92	5.5	0.59	4.4	6.8	*	*	*	*	*
2009	80	4.8	0.55	3.8	6.0	*	*	*	*	*
2010	76	4.3	0.50	3.3	5.4	*	*	*	*	*
2011	96	5.4	0.56	4.3	6.6	*	*	*	*	*
2012	92	5.1	0.55	4.1	6.3	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Esophagus										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	11,493	8.3	0.08	8.1	8.4	9,337	7.8	0.08	7.7	8.0
1986	367	8.0	0.43	7.1	8.8	252	6.2	0.41	5.4	7.1
1987	379	8.4	0.44	7.5	9.3	286	7.1	0.43	6.3	8.0
1988	370	8.1	0.44	7.3	9.0	260	6.5	0.42	5.7	7.4
1989	359	8.0	0.43	7.1	8.9	275	7.0	0.43	6.1	7.9
1990	347	7.6	0.42	6.8	8.4	248	6.2	0.41	5.4	7.0
1991	394	8.5	0.44	7.6	9.4	290	7.2	0.43	6.3	8.1
1992	398	8.4	0.43	7.6	9.3	309	7.5	0.43	6.6	8.4
1993	384	8.0	0.42	7.2	8.8	294	7.0	0.41	6.2	7.8
1994	403	8.5	0.43	7.7	9.4	312	7.5	0.44	6.7	8.4
1995	421	8.6	0.43	7.8	9.5	332	7.8	0.43	7.0	8.7
1996	419	8.5	0.42	7.7	9.4	338	7.9	0.44	7.1	8.9
1997	414	8.4	0.42	7.6	9.3	333	7.8	0.43	7.0	8.7
1998	442	8.9	0.43	8.1	9.8	367	8.6	0.45	7.7	9.5
1999	404	8.0	0.40	7.2	8.9	338	7.8	0.42	6.9	8.6
2000	441	8.7	0.42	7.9	9.5	353	8.0	0.43	7.2	8.9
2001	452	8.9	0.42	8.1	9.7	368	8.4	0.44	7.6	9.3
2002	417	8.1	0.40	7.3	8.9	353	8.0	0.43	7.2	8.9
2003	431	8.1	0.40	7.4	9.0	360	8.0	0.42	7.2	8.9
2004	425	8.0	0.39	7.2	8.8	368	8.1	0.42	7.3	8.9
2005	487	8.8	0.41	8.1	9.7	421	9.0	0.44	8.1	9.9
2006	434	7.9	0.39	7.2	8.7	375	8.0	0.42	7.2	8.9
2007	489	8.8	0.40	8.0	9.6	423	8.9	0.44	8.1	9.8
2008	501	8.5	0.39	7.8	9.3	431	8.6	0.42	7.8	9.5
2009	488	8.3	0.38	7.6	9.1	420	8.4	0.42	7.6	9.3
2010	474	7.9	0.37	7.2	8.6	414	8.2	0.41	7.4	9.0
2011	465	7.5	0.36	6.9	8.3	395	7.6	0.39	6.9	8.4
2012	488	7.9	0.37	7.2	8.7	422	8.2	0.40	7.4	9.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Esophagus										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,071	12.9	0.30	12.3	13.5	85	2.3	0.29	1.8	3.0
1986	115	23.6	2.30	19.3	28.5	*	*	*	*	*
1987	93	19.5	2.13	15.6	24.1	*	*	*	*	*
1988	109	21.3	2.10	17.4	25.8	*	*	*	*	*
1989	83	17.1	1.99	13.4	21.4	*	*	*	*	*
1990	94	18.5	1.98	14.8	22.7	*	*	*	*	*
1991	101	19.6	2.02	15.8	23.9	*	*	*	*	*
1992	85	16.7	1.90	13.2	20.8	*	*	*	*	*
1993	88	16.8	1.88	13.4	20.9	*	*	*	*	*
1994	88	16.8	1.89	13.4	20.9	*	*	*	*	*
1995	86	15.8	1.77	12.6	19.7	*	*	*	*	*
1996	79	13.6	1.60	10.7	17.1	*	*	*	*	*
1997	76	13.7	1.63	10.6	17.2	*	*	*	*	*
1998	75	12.8	1.53	10.0	16.2	*	*	*	*	*
1999	62	10.9	1.46	8.3	14.2	*	*	*	*	*
2000	87	15.2	1.72	12.0	18.9	*	*	*	*	*
2001	79	13.1	1.54	10.3	16.4	*	*	*	*	*
2002	61	10.2	1.35	7.7	13.1	*	*	*	*	*
2003	67	10.3	1.30	7.9	13.1	*	*	*	*	*
2004	57	9.6	1.35	7.1	12.5	*	*	*	*	*
2005	65	9.6	1.23	7.3	12.3	*	*	*	*	*
2006	54	8.8	1.28	6.5	11.6	*	*	*	*	*
2007	59	9.1	1.28	6.8	12.0	*	*	*	*	*
2008	66	9.9	1.30	7.6	12.8	*	*	*	*	*
2009	60	9.0	1.22	6.7	11.7	*	*	*	*	*
2010	54	7.2	1.02	5.3	9.4	*	*	*	*	*
2011	66	9.0	1.16	6.9	11.6	*	*	*	*	*
2012	62	8.4	1.13	6.4	10.9	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Esophagus										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,565	1.9	0.03	1.8	2.0	2,690	1.6	0.03	1.6	1.7
1986	112	1.8	0.18	1.5	2.2	78	1.4	0.16	1.1	1.7
1987	131	2.2	0.19	1.8	2.6	92	1.7	0.18	1.3	2.1
1988	113	1.8	0.17	1.5	2.1	81	1.4	0.15	1.1	1.7
1989	134	2.1	0.19	1.8	2.5	92	1.6	0.16	1.3	1.9
1990	132	2.1	0.18	1.7	2.5	92	1.6	0.17	1.3	2.0
1991	133	2.0	0.18	1.7	2.4	94	1.6	0.17	1.3	2.0
1992	137	2.1	0.18	1.7	2.5	110	1.9	0.18	1.5	2.2
1993	126	1.9	0.17	1.6	2.2	89	1.5	0.16	1.2	1.9
1994	149	2.3	0.19	1.9	2.7	105	1.8	0.17	1.4	2.2
1995	136	2.0	0.18	1.7	2.4	101	1.7	0.17	1.4	2.1
1996	135	2.0	0.17	1.7	2.4	94	1.5	0.16	1.2	1.9
1997	137	2.1	0.18	1.7	2.4	102	1.8	0.18	1.4	2.1
1998	134	1.9	0.17	1.6	2.3	108	1.8	0.17	1.5	2.2
1999	111	1.6	0.15	1.3	1.9	78	1.3	0.15	1.0	1.6
2000	151	2.1	0.17	1.8	2.5	126	2.0	0.18	1.7	2.4
2001	137	1.9	0.17	1.6	2.3	98	1.6	0.16	1.3	1.9
2002	147	2.0	0.17	1.7	2.4	116	1.8	0.17	1.5	2.2
2003	131	1.8	0.16	1.5	2.2	108	1.7	0.17	1.4	2.1
2004	136	1.9	0.17	1.6	2.3	102	1.7	0.17	1.3	2.0
2005	120	1.7	0.15	1.4	2.0	93	1.5	0.16	1.2	1.9
2006	134	1.9	0.16	1.6	2.2	107	1.8	0.17	1.4	2.2
2007	129	1.7	0.15	1.4	2.0	105	1.6	0.16	1.3	2.0
2008	132	1.8	0.16	1.5	2.1	104	1.7	0.17	1.4	2.0
2009	114	1.5	0.14	1.2	1.8	94	1.5	0.15	1.2	1.8
2010	148	1.9	0.16	1.6	2.3	123	1.9	0.18	1.6	2.3
2011	123	1.6	0.14	1.3	1.9	89	1.4	0.15	1.1	1.7
2012	143	1.8	0.15	1.5	2.1	109	1.6	0.16	1.3	2.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Esophagus										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	837	3.8	0.13	3.5	4.1	38	1.0	0.16	0.7	1.3
1986	34	5.2	0.91	3.6	7.3	*	*	*	*	*
1987	38	5.9	0.98	4.2	8.1	*	*	*	*	*
1988	32	4.7	0.85	3.2	6.7	*	*	*	*	*
1989	42	6.1	0.96	4.4	8.3	*	*	*	*	*
1990	40	6.0	0.97	4.3	8.2	*	*	*	*	*
1991	38	5.4	0.89	3.8	7.4	*	*	*	*	*
1992	25	3.4	0.69	2.2	5.1	*	*	*	*	*
1993	36	5.0	0.84	3.5	6.9	*	*	*	*	*
1994	44	5.8	0.87	4.2	7.7	*	*	*	*	*
1995	33	4.4	0.77	3.0	6.1	*	*	*	*	*
1996	37	4.9	0.81	3.4	6.7	*	*	*	*	*
1997	34	4.3	0.75	3.0	6.0	*	*	*	*	*
1998	26	3.2	0.63	2.1	4.7	*	*	*	*	*
1999	33	4.1	0.71	2.8	5.7	*	*	*	*	*
2000	24	2.9	0.59	1.8	4.3	*	*	*	*	*
2001	38	4.6	0.75	3.3	6.3	*	*	*	*	*
2002	27	3.2	0.62	2.1	4.6	*	*	*	*	*
2003	22	2.5	0.54	1.6	3.8	*	*	*	*	*
2004	32	3.6	0.64	2.5	5.1	*	*	*	*	*
2005	26	3.0	0.58	1.9	4.3	*	*	*	*	*
2006	25	2.8	0.56	1.8	4.1	*	*	*	*	*
2007	23	2.5	0.53	1.6	3.8	*	*	*	*	*
2008	26	2.7	0.53	1.7	3.9	*	*	*	*	*
2009	20	2.0	0.46	1.2	3.2	*	*	*	*	*
2010	22	2.2	0.47	1.4	3.3	*	*	*	*	*
2011	30	2.8	0.53	1.9	4.1	*	*	*	*	*
2012	30	2.9	0.54	2.0	4.2	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Stomach										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	16,122	5.0	0.04	4.9	5.1	12,270	4.4	0.04	4.3	4.5
1986	715	7.0	0.26	6.5	7.5	583	6.4	0.27	5.9	6.9
1987	710	6.8	0.26	6.3	7.4	555	6.0	0.26	5.5	6.6
1988	656	6.3	0.25	5.8	6.8	499	5.4	0.24	4.9	5.9
1989	730	6.9	0.26	6.4	7.4	585	6.2	0.26	5.7	6.7
1990	688	6.4	0.25	5.9	6.9	546	5.7	0.25	5.3	6.2
1991	697	6.4	0.24	5.9	6.9	560	5.8	0.24	5.3	6.3
1992	622	5.6	0.22	5.2	6.0	477	4.9	0.22	4.4	5.3
1993	684	6.1	0.24	5.7	6.6	517	5.3	0.23	4.8	5.8
1994	645	5.7	0.23	5.3	6.2	513	5.2	0.23	4.7	5.6
1995	659	5.8	0.22	5.3	6.2	520	5.2	0.23	4.7	5.6
1996	602	5.2	0.21	4.8	5.7	452	4.5	0.21	4.1	4.9
1997	579	5.0	0.21	4.6	5.4	448	4.4	0.21	4.0	4.8
1998	642	5.5	0.22	5.0	5.9	485	4.7	0.21	4.3	5.2
1999	601	5.1	0.21	4.7	5.5	461	4.4	0.21	4.1	4.9
2000	556	4.6	0.20	4.2	5.0	412	3.9	0.19	3.6	4.3
2001	599	4.9	0.20	4.5	5.3	453	4.3	0.20	3.9	4.7
2002	544	4.5	0.19	4.1	4.8	414	3.9	0.19	3.6	4.3
2003	571	4.6	0.19	4.2	5.0	417	3.9	0.19	3.5	4.3
2004	578	4.6	0.19	4.3	5.0	431	4.0	0.19	3.7	4.4
2005	525	4.2	0.18	3.8	4.6	393	3.6	0.18	3.3	4.0
2006	505	4.0	0.18	3.6	4.3	376	3.5	0.18	3.1	3.8
2007	485	3.8	0.17	3.4	4.1	337	3.1	0.17	2.7	3.4
2008	508	3.9	0.17	3.5	4.2	371	3.3	0.17	3.0	3.7
2009	541	4.1	0.18	3.7	4.4	408	3.6	0.18	3.3	4.0
2010	460	3.4	0.16	3.1	3.7	327	2.9	0.16	2.6	3.2
2011	507	3.7	0.17	3.4	4.0	371	3.2	0.17	2.9	3.6
2012	513	3.7	0.17	3.4	4.0	359	3.1	0.16	2.8	3.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Stomach										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,340	9.5	0.17	9.2	9.9	512	6.6	0.32	6.0	7.3
1986	117	12.2	1.18	10.0	14.7	15	12.6	3.92	6.1	22.2
1987	148	14.0	1.19	11.8	16.5	*	*	*	*	*
1988	143	13.4	1.17	11.2	15.9	14	10.8	3.46	5.1	19.3
1989	132	12.5	1.13	10.4	14.9	13	10.1	3.16	4.9	17.9
1990	128	11.9	1.09	9.9	14.2	14	9.5	3.15	4.4	17.2
1991	127	11.8	1.08	9.8	14.1	10	8.0	2.91	3.3	15.2
1992	129	11.1	1.01	9.2	13.3	16	8.2	2.33	4.3	13.9
1993	150	13.1	1.10	11.0	15.4	17	11.4	2.98	6.3	18.4
1994	111	9.4	0.92	7.7	11.4	21	10.1	2.47	5.9	16.0
1995	118	9.7	0.92	8.0	11.7	21	10.7	2.56	6.3	16.7
1996	137	11.1	0.97	9.2	13.1	13	5.3	1.58	2.7	9.3
1997	109	9.0	0.88	7.3	10.9	22	8.8	2.13	5.2	13.8
1998	141	11.1	0.95	9.3	13.1	16	5.7	1.55	3.1	9.5
1999	118	9.2	0.87	7.6	11.1	22	6.8	1.56	4.1	10.6
2000	119	9.2	0.86	7.6	11.0	25	8.8	1.97	5.4	13.4
2001	121	8.9	0.83	7.4	10.7	25	7.8	1.72	4.8	11.8
2002	110	8.0	0.78	6.5	9.7	20	7.1	1.73	4.1	11.1
2003	128	9.5	0.86	7.9	11.3	26	8.9	1.90	5.6	13.3
2004	132	9.2	0.82	7.7	10.9	15	4.8	1.29	2.6	7.9
2005	111	7.7	0.75	6.3	9.3	21	5.9	1.40	3.5	9.2
2006	110	7.5	0.73	6.2	9.1	19	4.9	1.21	2.9	7.8
2007	124	8.2	0.75	6.8	9.8	24	5.5	1.22	3.4	8.4
2008	123	8.5	0.78	7.0	10.1	14	2.7	0.77	1.4	4.6
2009	107	6.7	0.66	5.5	8.1	26	5.6	1.17	3.5	8.3
2010	113	7.0	0.67	5.7	8.4	20	5.0	1.18	3.0	7.8
2011	109	6.9	0.67	5.6	8.3	27	5.6	1.16	3.6	8.3
2012	125	7.8	0.71	6.5	9.3	29	4.9	0.98	3.2	7.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Stomach										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	9,438	7.1	0.07	7.0	7.3	7,155	6.3	0.08	6.1	6.4
1986	410	10.1	0.52	9.1	11.1	346	9.5	0.54	8.5	10.6
1987	399	9.6	0.50	8.6	10.6	298	8.2	0.50	7.3	9.3
1988	401	9.6	0.50	8.6	10.6	293	8.1	0.49	7.2	9.1
1989	448	10.8	0.53	9.8	11.9	353	9.7	0.54	8.7	10.8
1990	390	9.2	0.48	8.2	10.2	310	8.3	0.49	7.4	9.3
1991	428	9.7	0.48	8.7	10.7	337	8.7	0.49	7.7	9.7
1992	371	8.3	0.44	7.4	9.2	284	7.3	0.44	6.4	8.2
1993	394	8.8	0.46	8.0	9.8	284	7.3	0.44	6.5	8.2
1994	342	7.5	0.42	6.7	8.4	265	6.7	0.42	5.9	7.6
1995	393	8.6	0.45	7.8	9.6	316	8.0	0.46	7.1	8.9
1996	358	8.0	0.44	7.2	8.9	272	7.0	0.44	6.2	7.9
1997	357	7.7	0.42	6.9	8.5	274	6.8	0.42	6.0	7.6
1998	376	7.9	0.41	7.1	8.7	282	6.8	0.41	6.0	7.7
1999	337	7.0	0.39	6.3	7.8	258	6.2	0.39	5.4	7.0
2000	331	6.8	0.38	6.1	7.6	250	6.0	0.38	5.3	6.8
2001	333	6.6	0.37	5.9	7.4	255	5.9	0.37	5.2	6.6
2002	321	6.5	0.37	5.8	7.2	254	5.9	0.38	5.2	6.7
2003	338	6.7	0.37	6.0	7.4	256	5.9	0.37	5.2	6.7
2004	348	6.7	0.37	6.0	7.5	257	5.8	0.37	5.1	6.6
2005	285	5.6	0.34	5.0	6.3	212	4.9	0.34	4.2	5.6
2006	287	5.3	0.32	4.7	6.0	219	4.8	0.32	4.1	5.4
2007	290	5.4	0.32	4.8	6.0	198	4.3	0.31	3.7	5.0
2008	293	5.2	0.31	4.6	5.8	207	4.3	0.30	3.7	5.0
2009	313	5.4	0.31	4.8	6.1	229	4.7	0.31	4.1	5.3
2010	280	4.9	0.30	4.3	5.5	202	4.1	0.30	3.6	4.7
2011	301	5.1	0.30	4.6	5.8	221	4.4	0.30	3.8	5.1
2012	314	5.2	0.30	4.6	5.8	223	4.3	0.29	3.7	4.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Stomach										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,995	14.2	0.34	13.6	14.9	288	8.5	0.57	7.4	9.7
1986	57	15.0	2.18	11.0	19.7	*	*	*	*	*
1987	96	21.6	2.36	17.2	26.6	*	*	*	*	*
1988	97	21.9	2.40	17.5	27.1	11	18.1	6.62	7.6	34.9
1989	87	20.3	2.35	16.0	25.4	*	*	*	*	*
1990	69	15.3	1.94	11.7	19.5	11	15.8	6.02	6.3	31.1
1991	83	18.3	2.15	14.3	22.9	*	*	*	*	*
1992	78	17.0	2.07	13.2	21.5	*	*	*	*	*
1993	100	21.9	2.36	17.5	26.9	10	16.3	5.50	7.3	29.9
1994	63	12.9	1.72	9.7	16.6	14	16.2	5.14	7.7	28.8
1995	66	14.0	1.86	10.6	18.1	11	14.5	4.97	6.4	26.7
1996	80	16.5	1.97	12.8	20.7	*	*	*	*	*
1997	71	15.1	1.92	11.6	19.2	12	10.0	3.60	4.2	19.2
1998	85	17.0	1.96	13.3	21.2	*	*	*	*	*
1999	70	14.4	1.83	11.0	18.3	*	*	*	*	*
2000	66	12.5	1.65	9.5	16.1	15	12.0	3.71	5.9	21.1
2001	67	12.5	1.65	9.5	16.1	11	7.6	2.78	3.1	14.7
2002	55	9.9	1.41	7.4	13.0	12	10.9	3.61	5.0	19.7
2003	69	12.2	1.55	9.3	15.6	13	8.5	2.70	4.1	15.2
2004	81	13.4	1.59	10.5	16.8	10	7.0	2.36	3.2	13.0
2005	61	11.2	1.53	8.4	14.5	12	7.1	2.32	3.3	12.9
2006	59	9.9	1.36	7.4	12.9	*	*	*	*	*
2007	82	13.8	1.63	10.8	17.4	10	5.0	1.78	2.1	9.6
2008	75	13.0	1.60	10.1	16.5	11	4.2	1.37	1.9	7.8
2009	74	11.4	1.39	8.8	14.4	10	4.7	1.62	2.1	8.9
2010	67	10.9	1.41	8.3	14.0	11	7.6	2.41	3.6	13.4
2011	68	10.9	1.39	8.4	13.9	12	5.7	1.85	2.7	10.3
2012	69	11.3	1.43	8.7	14.4	22	8.9	2.09	5.3	13.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Stomach										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	6,684	3.5	0.04	3.4	3.6	5,115	3.1	0.04	3.0	3.2
1986	305	4.9	0.29	4.4	5.5	237	4.3	0.28	3.7	4.9
1987	311	5.0	0.28	4.4	5.6	257	4.6	0.29	4.1	5.2
1988	255	4.0	0.25	3.5	4.6	206	3.6	0.26	3.1	4.2
1989	282	4.3	0.26	3.9	4.9	232	4.0	0.26	3.5	4.6
1990	298	4.6	0.27	4.1	5.2	236	4.1	0.27	3.6	4.7
1991	269	4.0	0.24	3.5	4.5	223	3.7	0.25	3.2	4.2
1992	251	3.7	0.24	3.3	4.2	193	3.2	0.23	2.8	3.7
1993	290	4.3	0.25	3.8	4.8	233	3.9	0.26	3.4	4.5
1994	303	4.4	0.26	4.0	5.0	248	4.1	0.27	3.6	4.7
1995	266	3.9	0.24	3.4	4.4	204	3.4	0.24	2.9	3.9
1996	244	3.5	0.23	3.1	4.0	180	3.0	0.22	2.5	3.4
1997	222	3.1	0.21	2.7	3.6	174	2.8	0.21	2.4	3.2
1998	266	3.7	0.23	3.3	4.2	203	3.2	0.23	2.8	3.7
1999	264	3.7	0.23	3.3	4.2	203	3.2	0.23	2.8	3.7
2000	225	3.1	0.21	2.7	3.5	162	2.6	0.20	2.2	3.0
2001	266	3.7	0.23	3.2	4.1	198	3.1	0.22	2.7	3.6
2002	223	3.1	0.21	2.7	3.5	160	2.5	0.20	2.1	2.9
2003	233	3.1	0.20	2.7	3.5	161	2.5	0.20	2.1	2.9
2004	230	3.1	0.21	2.8	3.6	174	2.8	0.21	2.4	3.2
2005	240	3.3	0.21	2.9	3.7	181	2.9	0.22	2.5	3.3
2006	218	3.0	0.20	2.6	3.4	157	2.5	0.21	2.1	3.0
2007	195	2.6	0.19	2.3	3.0	139	2.2	0.19	1.8	2.6
2008	215	2.9	0.20	2.5	3.3	164	2.6	0.21	2.2	3.0
2009	228	3.0	0.20	2.6	3.4	179	2.8	0.21	2.4	3.2
2010	180	2.3	0.18	2.0	2.7	125	2.0	0.18	1.6	2.4
2011	206	2.6	0.19	2.3	3.0	150	2.3	0.19	1.9	2.7
2012	199	2.6	0.18	2.2	2.9	136	2.1	0.18	1.7	2.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Stomach										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,345	6.5	0.18	6.1	6.8	224	5.2	0.37	4.5	6.0
1986	60	10.5	1.38	7.9	13.5	*	*	*	*	*
1987	52	8.7	1.23	6.4	11.4	*	*	*	*	*
1988	46	7.7	1.16	5.6	10.3	*	*	*	*	*
1989	45	7.4	1.13	5.4	9.9	*	*	*	*	*
1990	59	9.4	1.24	7.1	12.1	*	*	*	*	*
1991	44	7.2	1.10	5.2	9.7	*	*	*	*	*
1992	51	7.4	1.05	5.5	9.7	*	*	*	*	*
1993	50	7.4	1.06	5.5	9.8	*	*	*	*	*
1994	48	7.0	1.01	5.1	9.2	*	*	*	*	*
1995	52	7.1	0.99	5.3	9.3	10	8.4	2.82	3.8	15.6
1996	57	7.6	1.02	5.7	9.9	*	*	*	*	*
1997	38	5.3	0.86	3.7	7.2	10	7.8	2.64	3.5	14.5
1998	56	7.3	0.98	5.5	9.5	*	*	*	*	*
1999	48	6.2	0.90	4.6	8.2	13	6.3	1.81	3.3	11.0
2000	53	6.8	0.95	5.1	8.9	10	6.5	2.21	2.9	12.1
2001	54	6.7	0.91	5.0	8.7	14	7.9	2.26	4.1	13.5
2002	55	6.7	0.90	5.0	8.7	*	*	*	*	*
2003	59	7.4	0.97	5.6	9.5	13	8.6	2.49	4.4	14.6
2004	51	6.2	0.87	4.6	8.1	*	*	*	*	*
2005	50	5.8	0.82	4.3	7.6	*	*	*	*	*
2006	51	5.9	0.83	4.4	7.7	10	4.4	1.47	2.1	8.2
2007	42	4.7	0.73	3.4	6.4	14	5.9	1.66	3.1	10.0
2008	48	5.6	0.81	4.1	7.4	*	*	*	*	*
2009	33	3.5	0.62	2.4	5.0	16	6.3	1.64	3.5	10.2
2010	46	4.6	0.69	3.4	6.2	*	*	*	*	*
2011	41	4.3	0.68	3.1	5.8	15	5.5	1.49	3.0	9.2
2012	56	5.7	0.78	4.3	7.5	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Colon and Rectum										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	72,266	22.4	0.08	22.2	22.5	60,819	21.6	0.09	21.4	21.8
1986	2,965	28.6	0.53	27.6	29.7	2,614	28.2	0.56	27.1	29.3
1987	2,987	28.6	0.53	27.6	29.7	2,638	28.3	0.56	27.2	29.4
1988	3,057	28.9	0.53	27.9	30.0	2,642	28.1	0.55	27.0	29.2
1989	2,891	27.2	0.51	26.2	28.2	2,514	26.5	0.53	25.4	27.5
1990	3,014	28.1	0.52	27.1	29.1	2,577	27.0	0.54	25.9	28.0
1991	2,853	26.2	0.49	25.3	27.2	2,479	25.6	0.52	24.6	26.6
1992	2,903	26.2	0.49	25.3	27.2	2,501	25.5	0.51	24.5	26.5
1993	2,938	26.2	0.48	25.3	27.2	2,518	25.4	0.51	24.4	26.4
1994	2,863	25.3	0.47	24.4	26.3	2,475	24.8	0.50	23.8	25.8
1995	2,847	24.9	0.47	24.0	25.8	2,454	24.3	0.49	23.3	25.3
1996	2,670	23.1	0.45	22.3	24.0	2,252	22.2	0.47	21.2	23.1
1997	2,751	23.6	0.45	22.8	24.5	2,325	22.7	0.47	21.8	23.7
1998	2,708	23.0	0.44	22.1	23.8	2,270	22.0	0.46	21.1	22.9
1999	2,760	23.2	0.44	22.3	24.0	2,327	22.3	0.46	21.4	23.3
2000	2,802	23.3	0.44	22.4	24.2	2,352	22.4	0.46	21.5	23.4
2001	2,722	22.4	0.43	21.6	23.3	2,280	21.6	0.45	20.7	22.5
2002	2,705	22.1	0.43	21.3	23.0	2,257	21.2	0.45	20.4	22.1
2003	2,618	21.1	0.41	20.3	22.0	2,141	19.9	0.43	19.1	20.8
2004	2,512	20.1	0.40	19.3	20.9	2,074	19.2	0.42	18.4	20.0
2005	2,363	18.7	0.39	17.9	19.5	1,935	17.8	0.41	17.0	18.6
2006	2,502	19.5	0.39	18.7	20.2	2,028	18.3	0.41	17.5	19.1
2007	2,468	19.0	0.38	18.2	19.7	2,010	18.0	0.40	17.2	18.8
2008	2,392	18.2	0.38	17.5	18.9	1,925	17.2	0.40	16.4	18.0
2009	2,283	17.0	0.36	16.3	17.7	1,863	16.3	0.38	15.6	17.1
2010	2,298	16.8	0.35	16.1	17.5	1,861	16.0	0.38	15.3	16.8
2011	2,178	15.7	0.34	15.1	16.4	1,734	14.8	0.36	14.1	15.5
2012	2,216	15.8	0.34	15.1	16.5	1,773	15.0	0.36	14.3	15.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Colon and Rectum										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	10,647	30.4	0.30	29.8	31.0	800	10.8	0.42	10.0	11.7
1986	341	34.7	1.95	31.0	38.7	10	7.1	2.26	3.4	13.1
1987	340	33.2	1.87	29.6	37.0	*	*	*	*	*
1988	400	37.2	1.93	33.5	41.2	15	10.9	3.15	5.6	18.6
1989	356	34.3	1.89	30.7	38.2	21	15.3	3.81	8.7	24.3
1990	420	39.5	1.99	35.6	43.5	17	14.9	4.20	7.8	24.8
1991	357	33.0	1.80	29.5	36.6	17	14.3	3.90	7.7	23.4
1992	386	34.4	1.81	31.0	38.1	16	10.3	2.97	5.4	17.5
1993	403	35.8	1.83	32.3	39.6	17	9.2	2.38	5.1	14.9
1994	363	30.8	1.66	27.6	34.2	25	15.5	3.49	9.4	23.5
1995	369	31.3	1.67	28.1	34.7	24	15.2	3.47	9.2	23.1
1996	398	32.7	1.68	29.5	36.1	20	8.6	2.18	4.9	13.8
1997	404	33.5	1.70	30.2	37.0	22	9.7	2.33	5.7	15.1
1998	408	32.3	1.63	29.1	35.6	30	13.8	2.83	8.8	20.2
1999	407	32.3	1.64	29.2	35.7	26	11.9	2.58	7.4	17.8
2000	418	31.8	1.59	28.8	35.1	32	11.8	2.31	7.7	17.0
2001	399	29.9	1.53	27.0	33.0	43	14.7	2.44	10.3	20.1
2002	415	30.7	1.54	27.8	33.9	33	11.3	2.10	7.6	16.1
2003	443	32.4	1.57	29.4	35.6	34	10.0	1.88	6.6	14.3
2004	419	29.8	1.49	27.0	32.9	19	5.6	1.40	3.2	9.0
2005	391	27.4	1.42	24.7	30.3	37	10.8	1.90	7.4	15.1
2006	422	29.0	1.44	26.2	31.9	52	13.2	1.98	9.6	17.6
2007	423	27.8	1.38	25.1	30.6	35	8.3	1.48	5.6	11.6
2008	423	27.8	1.38	25.2	30.7	44	9.6	1.55	6.8	13.1
2009	368	23.4	1.25	21.0	25.9	52	12.0	1.78	8.7	15.9
2010	387	24.0	1.25	21.6	26.6	50	10.9	1.65	7.9	14.5
2011	395	24.2	1.25	21.9	26.8	49	9.7	1.48	7.0	13.0
2012	392	23.1	1.20	20.9	25.6	51	9.5	1.41	6.9	12.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Colon and Rectum

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	35,886	27.7	0.15	27.4	28.0	30,323	26.9	0.16	26.6	27.2
1986	1,473	36.2	0.99	34.3	38.2	1,311	36.1	1.05	34.1	38.2
1987	1,452	35.6	0.98	33.7	37.6	1,286	35.5	1.03	33.5	37.6
1988	1,499	36.3	0.98	34.4	38.3	1,297	35.5	1.03	33.5	37.6
1989	1,466	35.7	0.98	33.8	37.7	1,276	35.0	1.02	33.0	37.0
1990	1,501	35.7	0.96	33.8	37.6	1,289	34.5	1.00	32.6	36.5
1991	1,419	33.4	0.92	31.6	35.3	1,248	33.1	0.97	31.3	35.1
1992	1,469	33.9	0.92	32.1	35.7	1,269	33.1	0.96	31.2	35.0
1993	1,414	32.2	0.88	30.4	33.9	1,223	31.4	0.92	29.6	33.2
1994	1,443	32.5	0.88	30.8	34.3	1,257	32.1	0.93	30.3	33.9
1995	1,389	31.2	0.86	29.5	32.9	1,197	30.5	0.91	28.8	32.4
1996	1,303	28.9	0.82	27.3	30.5	1,111	28.1	0.86	26.4	29.8
1997	1,353	29.6	0.82	28.0	31.3	1,144	28.6	0.86	27.0	30.4
1998	1,348	28.7	0.80	27.2	30.3	1,133	27.7	0.84	26.1	29.4
1999	1,358	28.9	0.80	27.4	30.5	1,149	28.0	0.84	26.4	29.7
2000	1,349	28.2	0.78	26.7	29.7	1,127	27.1	0.82	25.5	28.7
2001	1,352	28.0	0.77	26.5	29.5	1,137	27.1	0.81	25.5	28.7
2002	1,326	26.9	0.75	25.5	28.5	1,115	26.2	0.79	24.6	27.8
2003	1,358	27.7	0.76	26.2	29.2	1,115	26.3	0.80	24.7	27.9
2004	1,235	24.5	0.71	23.2	26.0	1,021	23.5	0.75	22.1	25.0
2005	1,190	23.2	0.68	21.9	24.6	973	22.1	0.72	20.7	23.5
2006	1,211	23.2	0.68	21.9	24.6	992	22.1	0.71	20.8	23.6
2007	1,170	22.0	0.66	20.8	23.4	952	21.0	0.69	19.7	22.4
2008	1,211	22.2	0.65	20.9	23.5	979	20.9	0.68	19.6	22.3
2009	1,173	21.0	0.63	19.8	22.3	965	20.3	0.66	19.0	21.7
2010	1,159	20.1	0.60	19.0	21.3	936	19.1	0.64	17.9	20.4
2011	1,079	18.3	0.57	17.2	19.5	865	17.3	0.60	16.2	18.5
2012	1,186	19.9	0.59	18.8	21.1	956	19.0	0.63	17.8	20.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Colon and Rectum

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	5,136	37.2	0.55	36.1	38.3	427	13.4	0.74	12.0	14.9
1986	154	39.0	3.40	32.6	46.1	*	*	*	*	*
1987	160	38.2	3.23	32.1	45.0	*	*	*	*	*
1988	193	45.4	3.53	38.7	52.8	*	*	*	*	*
1989	178	44.3	3.60	37.5	51.8	12	14.2	4.32	7.1	25.6
1990	201	48.3	3.66	41.4	55.9	11	25.3	8.67	11.2	46.3
1991	164	38.2	3.23	32.1	45.0	*	*	*	*	*
1992	191	43.7	3.41	37.3	50.8	*	*	*	*	*
1993	179	41.6	3.37	35.3	48.6	12	16.6	5.22	8.0	29.5
1994	176	39.4	3.20	33.4	46.1	10	12.2	4.09	5.6	22.7
1995	182	40.4	3.22	34.4	47.2	10	12.4	4.60	5.1	24.0
1996	181	37.6	3.01	31.9	43.9	11	11.6	4.23	4.8	22.2
1997	193	38.9	2.97	33.3	45.1	16	18.2	5.48	9.1	31.3
1998	202	39.7	2.97	34.1	45.9	13	12.0	3.88	5.6	21.6
1999	198	40.6	3.07	34.8	47.0	11	15.7	5.36	7.0	28.6
2000	204	39.7	2.95	34.1	45.8	18	16.3	4.40	8.9	26.7
2001	189	35.8	2.79	30.6	41.7	26	20.6	4.58	12.7	31.2
2002	190	35.1	2.69	30.0	40.7	21	16.2	3.95	9.4	25.4
2003	225	42.4	3.01	36.7	48.6	18	12.7	3.50	6.8	21.0
2004	203	36.6	2.76	31.4	42.4	11	6.9	2.37	3.1	12.9
2005	201	35.6	2.70	30.5	41.2	16	11.4	3.22	6.0	19.1
2006	195	34.1	2.58	29.3	39.5	24	15.2	3.47	9.2	23.2
2007	204	33.6	2.51	28.8	38.8	14	6.9	2.06	3.5	12.0
2008	210	35.4	2.60	30.5	40.9	22	11.6	2.82	6.8	18.3
2009	176	27.4	2.20	23.3	32.0	32	16.1	3.15	10.6	23.3
2010	202	31.7	2.38	27.2	36.6	21	9.5	2.35	5.5	15.1
2011	190	29.3	2.25	25.1	34.0	24	11.5	2.60	7.0	17.5
2012	195	29.2	2.20	25.0	33.7	35	15.4	2.85	10.3	21.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Colon and Rectum

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	36,380	18.8	0.10	18.6	19.0	30,496	18.0	0.10	17.8	18.2
1986	1,492	24.0	0.63	22.7	25.2	1,303	23.2	0.65	22.0	24.5
1987	1,535	24.3	0.63	23.1	25.6	1,352	23.9	0.66	22.7	25.3
1988	1,558	24.4	0.62	23.2	25.7	1,345	23.5	0.65	22.2	24.8
1989	1,425	21.9	0.59	20.8	23.1	1,238	21.3	0.61	20.1	22.5
1990	1,513	23.1	0.60	22.0	24.3	1,288	22.0	0.62	20.8	23.2
1991	1,434	21.7	0.58	20.5	22.8	1,231	20.7	0.60	19.6	21.9
1992	1,434	21.3	0.57	20.2	22.5	1,232	20.6	0.59	19.4	21.8
1993	1,524	22.4	0.58	21.2	23.5	1,295	21.4	0.60	20.2	22.6
1994	1,420	20.7	0.55	19.6	21.8	1,218	19.9	0.58	18.8	21.1
1995	1,458	20.9	0.55	19.8	22.0	1,257	20.3	0.58	19.2	21.5
1996	1,367	19.5	0.53	18.5	20.6	1,141	18.4	0.55	17.3	19.5
1997	1,398	19.7	0.53	18.7	20.8	1,181	19.0	0.56	17.9	20.1
1998	1,360	19.0	0.52	18.0	20.1	1,137	18.2	0.55	17.1	19.3
1999	1,402	19.4	0.52	18.4	20.4	1,178	18.6	0.55	17.5	19.7
2000	1,453	19.9	0.53	18.9	21.0	1,225	19.2	0.56	18.1	20.3
2001	1,370	18.7	0.51	17.7	19.7	1,143	17.8	0.54	16.8	18.9
2002	1,379	18.6	0.51	17.7	19.7	1,142	17.6	0.53	16.6	18.7
2003	1,260	17.0	0.48	16.0	17.9	1,026	15.9	0.50	14.9	16.9
2004	1,277	17.1	0.48	16.1	18.0	1,053	16.2	0.51	15.2	17.2
2005	1,173	15.5	0.46	14.6	16.4	962	14.7	0.49	13.8	15.7
2006	1,291	16.8	0.47	15.9	17.7	1,036	15.5	0.49	14.6	16.5
2007	1,298	16.8	0.47	15.8	17.7	1,058	15.8	0.50	14.9	16.8
2008	1,181	15.3	0.45	14.4	16.2	946	14.3	0.48	13.4	15.3
2009	1,110	14.2	0.44	13.4	15.1	898	13.5	0.46	12.6	14.5
2010	1,139	14.3	0.43	13.4	15.2	925	13.6	0.46	12.8	14.6
2011	1,099	13.7	0.42	12.9	14.6	869	12.8	0.45	12.0	13.8
2012	1,030	12.6	0.40	11.8	13.4	817	11.9	0.43	11.1	12.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Colon and Rectum										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	5,511	26.1	0.36	25.5	26.8	373	9.0	0.49	8.0	10.0
1986	187	32.1	2.40	27.6	37.1	*	*	*	*	*
1987	180	29.7	2.26	25.4	34.4	*	*	*	*	*
1988	207	32.3	2.29	28.0	37.1	*	*	*	*	*
1989	178	28.2	2.16	24.2	32.8	*	*	*	*	*
1990	219	34.0	2.34	29.6	38.8	*	*	*	*	*
1991	193	29.7	2.17	25.6	34.3	10	14.2	5.08	6.0	26.8
1992	195	28.6	2.08	24.7	33.0	*	*	*	*	*
1993	224	32.5	2.20	28.4	37.1	*	*	*	*	*
1994	187	26.2	1.94	22.5	30.3	15	17.4	5.00	9.0	29.3
1995	187	26.1	1.93	22.4	30.1	14	16.4	4.75	8.5	27.7
1996	217	29.7	2.03	25.8	33.9	*	*	*	*	*
1997	211	29.0	2.01	25.2	33.1	*	*	*	*	*
1998	206	27.1	1.90	23.5	31.1	17	14.6	3.83	8.1	23.7
1999	209	27.2	1.90	23.6	31.2	15	10.2	2.83	5.4	17.1
2000	214	26.9	1.85	23.4	30.7	14	8.6	2.52	4.4	14.8
2001	210	26.1	1.81	22.6	29.8	17	10.3	2.67	5.8	16.7
2002	225	27.6	1.85	24.1	31.4	12	7.7	2.30	3.9	13.4
2003	218	26.2	1.79	22.8	29.9	16	8.2	2.17	4.5	13.4
2004	216	25.5	1.75	22.2	29.2	*	*	*	*	*
2005	190	22.3	1.63	19.2	25.7	21	10.6	2.40	6.4	16.2
2006	227	25.5	1.71	22.3	29.1	28	12.2	2.43	7.9	17.8
2007	219	23.9	1.64	20.8	27.4	21	9.3	2.10	5.6	14.2
2008	213	23.1	1.60	20.1	26.5	22	8.4	1.88	5.2	12.9
2009	192	20.6	1.50	17.7	23.7	20	8.6	1.99	5.1	13.3
2010	185	19.2	1.43	16.5	22.2	29	11.6	2.24	7.6	16.7
2011	205	21.0	1.48	18.2	24.1	25	8.5	1.79	5.4	12.7
2012	197	19.2	1.39	16.6	22.1	16	5.2	1.35	2.9	8.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Liver										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	12,228	3.8	0.03	3.7	3.8	9,352	3.3	0.03	3.3	3.4
1986	297	2.8	0.17	2.5	3.2	233	2.5	0.17	2.2	2.9
1987	333	3.1	0.17	2.8	3.5	257	2.7	0.17	2.4	3.1
1988	305	2.8	0.16	2.5	3.2	246	2.6	0.17	2.3	2.9
1989	359	3.3	0.18	3.0	3.7	278	2.9	0.17	2.6	3.3
1990	391	3.6	0.18	3.2	4.0	307	3.2	0.18	2.8	3.5
1991	365	3.3	0.17	3.0	3.7	289	3.0	0.18	2.6	3.3
1992	381	3.4	0.18	3.1	3.8	295	3.0	0.18	2.7	3.4
1993	409	3.6	0.18	3.3	4.0	322	3.2	0.18	2.9	3.6
1994	424	3.7	0.18	3.4	4.1	323	3.2	0.18	2.9	3.6
1995	473	4.1	0.19	3.7	4.5	359	3.6	0.19	3.2	4.0
1996	465	4.1	0.19	3.7	4.4	358	3.6	0.19	3.2	4.0
1997	469	4.0	0.19	3.7	4.4	351	3.5	0.19	3.1	3.9
1998	452	3.8	0.18	3.5	4.2	344	3.4	0.18	3.0	3.7
1999	464	3.9	0.18	3.6	4.3	344	3.3	0.18	3.0	3.7
2000	453	3.8	0.18	3.5	4.2	347	3.4	0.18	3.0	3.7
2001	444	3.7	0.17	3.3	4.0	322	3.1	0.17	2.8	3.5
2002	432	3.5	0.17	3.2	3.9	324	3.1	0.17	2.8	3.5
2003	477	3.9	0.18	3.5	4.2	365	3.4	0.18	3.1	3.8
2004	490	3.9	0.18	3.6	4.3	358	3.4	0.18	3.0	3.7
2005	507	4.0	0.18	3.6	4.3	402	3.7	0.19	3.4	4.1
2006	499	3.9	0.17	3.5	4.2	393	3.6	0.18	3.3	4.0
2007	495	3.8	0.17	3.5	4.2	371	3.4	0.18	3.0	3.8
2008	521	3.9	0.17	3.6	4.3	388	3.4	0.18	3.1	3.8
2009	539	4.0	0.17	3.6	4.3	415	3.7	0.18	3.3	4.0
2010	585	4.3	0.18	3.9	4.6	448	3.9	0.19	3.5	4.2
2011	592	4.2	0.17	3.8	4.5	441	3.7	0.18	3.4	4.1
2012	607	4.2	0.17	3.9	4.6	472	4.0	0.19	3.6	4.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Liver										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,343	6.3	0.13	6.0	6.6	533	6.4	0.30	5.8	7.0
1986	54	4.7	0.67	3.5	6.2	10	5.4	1.89	2.4	10.7
1987	68	6.2	0.78	4.8	7.9	*	*	*	*	*
1988	51	4.6	0.67	3.4	6.1	*	*	*	*	*
1989	69	5.9	0.73	4.6	7.5	12	6.5	1.97	3.2	11.7
1990	70	6.2	0.76	4.8	7.9	14	7.8	2.49	3.7	14.0
1991	65	5.5	0.70	4.2	7.0	11	6.7	2.16	3.2	12.2
1992	63	5.4	0.70	4.1	6.9	23	12.8	3.01	7.6	19.9
1993	75	6.3	0.74	4.9	7.9	12	6.0	1.86	3.0	10.7
1994	85	6.9	0.77	5.5	8.6	16	6.5	1.72	3.6	10.8
1995	97	8.0	0.83	6.4	9.7	17	9.7	2.59	5.3	15.8
1996	91	7.3	0.78	5.8	8.9	16	8.1	2.25	4.3	13.4
1997	92	7.0	0.75	5.6	8.6	26	10.8	2.36	6.7	16.3
1998	96	7.3	0.76	5.9	9.0	12	4.3	1.39	2.1	7.9
1999	96	7.1	0.74	5.8	8.7	24	7.3	1.59	4.6	11.1
2000	80	5.7	0.65	4.5	7.1	26	8.6	1.82	5.4	12.8
2001	101	7.0	0.71	5.7	8.6	21	5.7	1.36	3.4	9.0
2002	90	6.4	0.69	5.1	7.9	18	6.1	1.51	3.5	9.7
2003	85	5.9	0.65	4.7	7.3	27	8.2	1.70	5.2	12.1
2004	103	6.9	0.69	5.6	8.4	29	6.8	1.34	4.4	9.9
2005	84	5.3	0.59	4.2	6.6	21	5.8	1.37	3.4	9.0
2006	85	5.2	0.58	4.1	6.4	21	4.6	1.06	2.8	7.2
2007	103	6.4	0.64	5.2	7.8	21	4.6	1.11	2.7	7.3
2008	114	7.2	0.69	5.9	8.6	19	3.8	0.94	2.2	6.1
2009	94	5.5	0.58	4.4	6.8	30	6.8	1.34	4.5	9.9
2010	106	6.1	0.61	5.0	7.4	31	6.7	1.27	4.5	9.6
2011	114	6.2	0.60	5.1	7.5	37	6.6	1.14	4.5	9.2
2012	112	6.1	0.59	5.0	7.4	23	4.1	0.89	2.5	6.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Liver										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	8,014	5.8	0.07	5.7	5.9	6,101	5.2	0.07	5.0	5.3
1986	188	4.3	0.33	3.7	5.0	140	3.7	0.33	3.1	4.4
1987	217	5.0	0.35	4.3	5.7	160	4.2	0.34	3.5	4.9
1988	194	4.5	0.33	3.8	5.2	155	4.1	0.34	3.5	4.8
1989	217	4.9	0.34	4.3	5.6	173	4.5	0.35	3.8	5.2
1990	238	5.4	0.36	4.7	6.1	185	4.7	0.35	4.0	5.4
1991	213	4.7	0.33	4.0	5.4	164	4.1	0.33	3.5	4.8
1992	237	5.3	0.35	4.6	6.0	184	4.7	0.35	4.0	5.4
1993	243	5.4	0.35	4.7	6.1	194	4.9	0.36	4.2	5.7
1994	252	5.3	0.34	4.6	6.0	188	4.5	0.34	3.9	5.2
1995	296	6.3	0.38	5.6	7.1	219	5.3	0.37	4.6	6.1
1996	300	6.3	0.37	5.6	7.1	231	5.6	0.37	4.9	6.4
1997	291	6.0	0.36	5.3	6.7	214	5.1	0.35	4.4	5.8
1998	290	5.8	0.35	5.2	6.5	219	5.1	0.35	4.4	5.8
1999	286	5.7	0.34	5.0	6.4	211	4.9	0.34	4.2	5.6
2000	282	5.5	0.33	4.9	6.2	218	5.0	0.34	4.3	5.7
2001	280	5.5	0.33	4.8	6.2	205	4.7	0.33	4.1	5.4
2002	289	5.5	0.33	4.9	6.2	217	4.9	0.33	4.2	5.6
2003	333	6.3	0.35	5.7	7.1	256	5.7	0.36	5.0	6.5
2004	307	5.6	0.32	4.9	6.2	224	4.8	0.32	4.2	5.5
2005	355	6.4	0.35	5.7	7.1	280	6.0	0.36	5.3	6.7
2006	339	6.0	0.33	5.4	6.7	263	5.5	0.35	4.9	6.3
2007	333	5.8	0.32	5.2	6.4	244	5.0	0.33	4.4	5.7
2008	361	6.1	0.33	5.5	6.7	275	5.5	0.33	4.8	6.1
2009	397	6.6	0.34	6.0	7.3	309	6.1	0.36	5.5	6.9
2010	419	6.9	0.34	6.2	7.6	326	6.3	0.36	5.6	7.1
2011	435	6.9	0.34	6.2	7.6	320	6.0	0.35	5.4	6.8
2012	422	6.4	0.32	5.8	7.1	327	5.9	0.34	5.3	6.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Liver										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,519	9.7	0.26	9.2	10.2	394	10.1	0.58	9.0	11.3
1986	41	8.4	1.40	5.9	11.6	*	*	*	*	*
1987	50	11.0	1.67	8.0	14.8	*	*	*	*	*
1988	32	6.7	1.24	4.5	9.6	*	*	*	*	*
1989	35	7.3	1.28	5.0	10.2	*	*	*	*	*
1990	41	9.0	1.51	6.3	12.4	12	14.5	5.49	5.8	28.6
1991	40	8.1	1.36	5.6	11.1	*	*	*	*	*
1992	36	8.0	1.43	5.5	11.2	17	19.8	5.29	10.8	32.7
1993	43	8.3	1.33	5.9	11.3	*	*	*	*	*
1994	52	10.6	1.56	7.8	14.0	12	10.5	3.23	5.1	19.0
1995	64	13.0	1.74	9.8	16.8	13	18.6	6.01	8.7	33.1
1996	57	10.7	1.49	7.9	14.0	12	12.0	4.16	5.3	22.4
1997	58	10.5	1.46	7.8	13.7	19	15.8	3.89	9.1	25.2
1998	63	11.6	1.53	8.8	14.9	*	*	*	*	*
1999	59	10.6	1.43	7.9	13.7	16	11.3	3.10	6.0	19.0
2000	49	8.5	1.26	6.2	11.3	15	9.4	2.98	4.5	16.9
2001	55	9.2	1.32	6.8	12.1	20	12.3	3.24	6.8	20.2
2002	60	10.3	1.41	7.7	13.4	12	8.8	2.74	4.2	15.6
2003	57	9.2	1.28	6.9	12.1	20	12.9	3.33	7.2	20.8
2004	62	9.7	1.29	7.3	12.5	21	10.3	2.45	6.1	16.2
2005	61	8.9	1.19	6.8	11.6	14	6.9	2.06	3.5	12.1
2006	60	8.6	1.19	6.5	11.3	16	7.7	2.06	4.2	12.8
2007	74	10.9	1.33	8.5	13.8	15	6.4	1.93	3.2	11.3
2008	71	10.4	1.30	8.0	13.3	15	7.3	2.04	3.8	12.3
2009	66	8.8	1.16	6.7	11.4	22	12.1	2.88	7.1	18.8
2010	73	10.0	1.24	7.7	12.7	20	8.9	2.19	5.1	14.1
2011	85	10.8	1.23	8.6	13.5	30	11.8	2.34	7.6	17.2
2012	75	9.4	1.13	7.3	11.9	20	8.4	2.08	4.9	13.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Liver										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	4,214	2.2	0.03	2.2	2.3	3,251	2.0	0.03	1.9	2.0
1986	109	1.8	0.17	1.4	2.1	93	1.7	0.17	1.3	2.1
1987	116	1.8	0.17	1.5	2.2	97	1.7	0.17	1.4	2.1
1988	111	1.8	0.17	1.4	2.1	91	1.6	0.17	1.3	2.0
1989	142	2.2	0.19	1.9	2.6	105	1.9	0.19	1.5	2.3
1990	153	2.4	0.19	2.0	2.8	122	2.1	0.19	1.8	2.5
1991	152	2.3	0.19	2.0	2.7	125	2.2	0.20	1.8	2.6
1992	144	2.2	0.18	1.8	2.6	111	1.9	0.18	1.6	2.3
1993	166	2.5	0.19	2.1	2.9	128	2.1	0.19	1.8	2.5
1994	172	2.5	0.20	2.2	3.0	135	2.3	0.20	1.9	2.7
1995	177	2.6	0.20	2.2	3.0	140	2.4	0.20	2.0	2.8
1996	165	2.4	0.19	2.0	2.8	127	2.1	0.19	1.8	2.5
1997	178	2.6	0.20	2.2	3.0	137	2.3	0.20	1.9	2.7
1998	162	2.3	0.18	2.0	2.7	125	2.0	0.18	1.7	2.4
1999	178	2.5	0.19	2.2	2.9	133	2.2	0.19	1.8	2.6
2000	171	2.4	0.19	2.1	2.8	129	2.1	0.19	1.7	2.5
2001	164	2.3	0.18	2.0	2.7	117	1.8	0.17	1.5	2.2
2002	143	2.0	0.16	1.6	2.3	107	1.7	0.16	1.4	2.0
2003	144	2.0	0.16	1.7	2.3	109	1.7	0.17	1.4	2.1
2004	183	2.5	0.19	2.2	2.9	134	2.1	0.18	1.7	2.5
2005	152	2.0	0.16	1.7	2.4	122	1.9	0.17	1.5	2.2
2006	160	2.2	0.17	1.9	2.6	130	2.1	0.19	1.7	2.5
2007	162	2.2	0.18	1.9	2.6	127	2.0	0.18	1.7	2.4
2008	160	2.1	0.17	1.8	2.4	113	1.7	0.16	1.4	2.1
2009	142	1.9	0.16	1.6	2.2	106	1.6	0.16	1.3	2.0
2010	166	2.1	0.17	1.8	2.5	122	1.9	0.17	1.5	2.2
2011	157	2.1	0.17	1.7	2.4	121	1.9	0.17	1.5	2.2
2012	185	2.4	0.18	2.0	2.7	145	2.3	0.19	1.9	2.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Liver										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	824	3.9	0.14	3.6	4.1	139	3.4	0.30	2.8	4.1
1986	13	2.1	0.60	1.1	3.6	*	*	*	*	*
1987	18	2.9	0.68	1.7	4.5	*	*	*	*	*
1988	19	3.1	0.72	1.9	4.8	*	*	*	*	*
1989	34	4.9	0.87	3.4	6.9	*	*	*	*	*
1990	29	4.4	0.82	2.9	6.2	*	*	*	*	*
1991	25	3.6	0.74	2.3	5.4	*	*	*	*	*
1992	27	3.8	0.74	2.5	5.5	*	*	*	*	*
1993	32	4.6	0.83	3.1	6.5	*	*	*	*	*
1994	33	4.5	0.80	3.1	6.3	*	*	*	*	*
1995	33	4.7	0.82	3.2	6.5	*	*	*	*	*
1996	34	4.7	0.81	3.3	6.6	*	*	*	*	*
1997	34	4.5	0.79	3.1	6.3	*	*	*	*	*
1998	33	4.3	0.76	3.0	6.1	*	*	*	*	*
1999	37	4.7	0.78	3.3	6.5	*	*	*	*	*
2000	31	3.8	0.68	2.5	5.3	11	7.8	2.38	3.8	13.7
2001	46	5.5	0.82	4.1	7.4	*	*	*	*	*
2002	30	3.7	0.68	2.5	5.2	*	*	*	*	*
2003	28	3.4	0.64	2.2	4.9	*	*	*	*	*
2004	41	4.8	0.76	3.4	6.5	*	*	*	*	*
2005	23	2.6	0.55	1.6	3.9	*	*	*	*	*
2006	25	2.7	0.55	1.8	4.1	*	*	*	*	*
2007	29	3.2	0.60	2.1	4.6	*	*	*	*	*
2008	43	4.7	0.73	3.4	6.4	*	*	*	*	*
2009	28	3.0	0.58	2.0	4.4	*	*	*	*	*
2010	33	3.4	0.59	2.3	4.7	11	4.9	1.50	2.4	8.6
2011	29	2.9	0.55	1.9	4.2	*	*	*	*	*
2012	37	3.6	0.60	2.5	5.0	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Pancreas										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	36,097	11.2	0.06	11.0	11.3	30,232	10.8	0.06	10.7	10.9
1986	1,183	11.1	0.33	10.5	11.8	1,007	10.7	0.34	10.0	11.3
1987	1,192	11.3	0.33	10.6	11.9	1,016	10.8	0.34	10.1	11.5
1988	1,134	10.6	0.32	10.0	11.2	967	10.2	0.33	9.5	10.8
1989	1,129	10.5	0.32	9.9	11.2	940	9.9	0.32	9.2	10.5
1990	1,237	11.4	0.33	10.8	12.1	1,054	11.0	0.34	10.4	11.7
1991	1,244	11.4	0.33	10.8	12.1	1,055	10.9	0.34	10.3	11.6
1992	1,209	10.9	0.32	10.3	11.6	1,028	10.5	0.33	9.9	11.2
1993	1,209	10.8	0.31	10.2	11.4	997	10.1	0.32	9.5	10.7
1994	1,251	11.0	0.31	10.4	11.6	1,057	10.5	0.32	9.9	11.2
1995	1,183	10.3	0.30	9.7	10.9	1,020	10.1	0.32	9.5	10.8
1996	1,280	11.1	0.31	10.5	11.7	1,061	10.5	0.32	9.9	11.2
1997	1,267	10.9	0.31	10.3	11.5	1,067	10.5	0.32	9.9	11.1
1998	1,298	11.0	0.31	10.4	11.6	1,106	10.8	0.32	10.2	11.4
1999	1,349	11.4	0.31	10.8	12.0	1,154	11.1	0.33	10.5	11.8
2000	1,354	11.3	0.31	10.7	11.9	1,123	10.8	0.32	10.2	11.5
2001	1,377	11.4	0.31	10.8	12.0	1,166	11.2	0.33	10.5	11.8
2002	1,392	11.4	0.31	10.8	12.0	1,158	11.0	0.32	10.4	11.7
2003	1,332	10.8	0.30	10.2	11.4	1,111	10.5	0.31	9.8	11.1
2004	1,424	11.4	0.30	10.8	12.0	1,190	11.1	0.32	10.5	11.8
2005	1,482	11.7	0.31	11.1	12.3	1,212	11.1	0.32	10.5	11.8
2006	1,463	11.5	0.30	10.9	12.1	1,193	10.9	0.32	10.3	11.6
2007	1,424	11.0	0.29	10.5	11.6	1,188	10.8	0.31	10.1	11.4
2008	1,521	11.6	0.30	11.0	12.2	1,268	11.3	0.32	10.7	12.0
2009	1,518	11.4	0.29	10.8	11.9	1,251	11.0	0.32	10.4	11.7
2010	1,544	11.4	0.29	10.9	12.0	1,286	11.2	0.32	10.6	11.9
2011	1,560	11.4	0.29	10.8	12.0	1,296	11.3	0.32	10.6	11.9
2012	1,541	10.9	0.28	10.4	11.5	1,261	10.6	0.30	10.0	11.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Pancreas										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	5,380	15.2	0.21	14.8	15.6	483	6.6	0.33	6.0	7.3
1986	169	16.6	1.33	14.1	19.3	*	*	*	*	*
1987	170	16.1	1.27	13.7	18.7	*	*	*	*	*
1988	161	15.0	1.21	12.7	17.5	*	*	*	*	*
1989	175	16.2	1.27	13.9	18.9	13	12.6	4.11	5.9	22.5
1990	174	15.4	1.21	13.1	18.0	*	*	*	*	*
1991	177	16.0	1.24	13.7	18.6	12	9.2	3.06	4.2	16.7
1992	172	15.1	1.18	12.8	17.5	*	*	*	*	*
1993	200	17.1	1.24	14.7	19.7	12	7.0	2.38	3.1	12.9
1994	181	15.3	1.16	13.1	17.7	13	8.6	2.69	4.2	15.1
1995	150	12.9	1.08	10.9	15.2	13	6.3	1.86	3.2	10.9
1996	201	16.7	1.20	14.4	19.2	18	10.2	2.66	5.7	16.5
1997	188	14.6	1.08	12.5	16.9	12	6.6	2.13	3.1	11.8
1998	180	13.9	1.06	11.9	16.1	12	5.3	1.66	2.6	9.4
1999	172	13.2	1.02	11.3	15.4	23	10.1	2.29	6.2	15.4
2000	209	15.7	1.10	13.6	18.0	22	8.8	2.01	5.3	13.5
2001	198	14.7	1.07	12.7	17.0	13	5.0	1.47	2.5	8.5
2002	220	16.0	1.10	13.9	18.3	14	4.6	1.33	2.4	7.9
2003	194	13.9	1.02	12.0	16.0	27	8.2	1.68	5.2	12.0
2004	213	14.9	1.04	12.9	17.0	21	6.3	1.48	3.8	9.8
2005	248	17.2	1.11	15.1	19.5	22	6.8	1.57	4.1	10.5
2006	251	17.0	1.09	14.9	19.3	19	4.9	1.19	2.8	7.7
2007	212	14.5	1.01	12.6	16.6	24	5.4	1.20	3.3	8.2
2008	230	15.3	1.03	13.3	17.4	23	6.0	1.33	3.7	9.0
2009	236	15.0	1.00	13.1	17.1	31	7.0	1.35	4.6	10.1
2010	225	14.5	0.99	12.6	16.5	33	6.4	1.22	4.3	9.2
2011	233	14.2	0.95	12.4	16.2	31	6.2	1.17	4.1	8.9
2012	241	14.3	0.94	12.5	16.3	39	6.8	1.16	4.7	9.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.
*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*
** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Pancreas										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	17,432	13.0	0.10	12.8	13.2	14,658	12.6	0.11	12.4	12.8
1986	582	13.4	0.58	12.3	14.6	497	12.9	0.61	11.8	14.2
1987	561	13.1	0.58	12.0	14.3	479	12.7	0.60	11.5	13.9
1988	572	13.2	0.57	12.1	14.4	495	12.8	0.60	11.7	14.1
1989	526	12.1	0.55	11.0	13.2	440	11.4	0.57	10.4	12.6
1990	570	13.0	0.56	11.9	14.1	486	12.6	0.59	11.5	13.8
1991	619	14.1	0.58	12.9	15.2	532	13.7	0.61	12.5	14.9
1992	567	12.7	0.55	11.6	13.8	468	11.8	0.56	10.7	12.9
1993	568	12.6	0.55	11.6	13.8	473	11.9	0.56	10.9	13.1
1994	622	13.5	0.55	12.4	14.6	521	12.8	0.57	11.7	14.0
1995	563	11.9	0.51	10.9	12.9	476	11.5	0.53	10.4	12.6
1996	614	13.0	0.54	12.0	14.1	505	12.3	0.56	11.2	13.4
1997	601	12.6	0.52	11.6	13.7	501	12.1	0.55	11.1	13.2
1998	614	12.7	0.52	11.7	13.8	527	12.5	0.55	11.5	13.7
1999	630	12.9	0.52	11.9	13.9	543	12.8	0.55	11.7	13.9
2000	646	13.0	0.52	12.0	14.1	541	12.6	0.55	11.6	13.7
2001	649	12.9	0.51	11.9	14.0	558	12.8	0.55	11.8	14.0
2002	673	13.3	0.52	12.3	14.4	574	13.1	0.55	12.0	14.2
2003	663	12.9	0.51	11.9	13.9	539	12.2	0.53	11.2	13.3
2004	713	13.6	0.52	12.6	14.7	591	13.2	0.55	12.1	14.3
2005	700	13.3	0.51	12.3	14.3	572	12.7	0.54	11.6	13.8
2006	708	13.0	0.50	12.1	14.1	580	12.5	0.53	11.5	13.6
2007	672	12.3	0.48	11.4	13.3	567	12.1	0.51	11.1	13.1
2008	740	13.2	0.50	12.3	14.3	622	13.0	0.53	12.0	14.1
2009	770	13.4	0.49	12.5	14.4	647	13.2	0.53	12.2	14.3
2010	751	13.0	0.48	12.0	13.9	629	12.7	0.52	11.7	13.8
2011	768	13.0	0.48	12.1	14.0	658	13.2	0.53	12.2	14.3
2012	770	12.4	0.46	11.5	13.4	637	12.1	0.49	11.2	13.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Pancreas										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,525	17.4	0.37	16.7	18.1	248	7.7	0.55	6.6	8.8
1986	82	18.9	2.23	14.7	23.7	*	*	*	*	*
1987	79	18.1	2.15	14.2	22.8	*	*	*	*	*
1988	74	17.3	2.15	13.4	22.0	*	*	*	*	*
1989	81	18.0	2.14	14.0	22.6	*	*	*	*	*
1990	78	16.2	1.97	12.6	20.5	*	*	*	*	*
1991	79	17.9	2.12	14.0	22.5	*	*	*	*	*
1992	93	22.5	2.51	17.8	27.8	*	*	*	*	*
1993	89	19.9	2.25	15.8	24.8	*	*	*	*	*
1994	95	19.7	2.15	15.7	24.3	*	*	*	*	*
1995	78	16.8	2.03	13.1	21.2	*	*	*	*	*
1996	101	20.3	2.15	16.3	24.8	*	*	*	*	*
1997	92	17.0	1.84	13.6	21.0	*	*	*	*	*
1998	81	15.1	1.76	11.9	18.9	*	*	*	*	*
1999	76	14.5	1.74	11.3	18.2	11	11.7	4.09	5.1	21.8
2000	96	17.4	1.87	13.9	21.4	*	*	*	*	*
2001	85	15.1	1.74	11.9	18.8	*	*	*	*	*
2002	91	16.9	1.88	13.4	20.9	*	*	*	*	*
2003	108	18.4	1.88	14.9	22.4	16	11.4	3.21	6.0	19.2
2004	112	18.0	1.78	14.7	21.8	10	5.8	2.00	2.6	11.0
2005	115	19.4	1.91	15.9	23.5	13	9.0	2.82	4.3	15.8
2006	118	19.1	1.85	15.7	23.1	10	6.1	2.16	2.6	11.6
2007	93	15.7	1.73	12.5	19.4	12	6.0	1.97	2.8	11.0
2008	104	17.2	1.78	13.9	21.0	14	8.3	2.47	4.2	14.3
2009	107	16.7	1.71	13.5	20.4	16	8.2	2.26	4.4	13.7
2010	103	15.9	1.64	12.9	19.5	19	8.4	2.13	4.7	13.5
2011	99	14.3	1.52	11.5	17.5	11	5.2	1.74	2.4	9.6
2012	116	16.7	1.63	13.7	20.2	17	6.4	1.68	3.5	10.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Pancreas										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	18,665	9.8	0.07	9.6	9.9	15,574	9.4	0.08	9.2	9.5
1986	601	9.5	0.39	8.8	10.4	510	9.1	0.41	8.3	9.9
1987	631	10.0	0.40	9.2	10.8	537	9.5	0.41	8.7	10.3
1988	562	8.7	0.37	8.0	9.5	472	8.2	0.38	7.5	9.0
1989	603	9.4	0.39	8.7	10.2	500	8.8	0.40	8.1	9.6
1990	667	10.3	0.40	9.5	11.1	568	9.8	0.42	9.0	10.7
1991	625	9.6	0.39	8.8	10.4	523	9.0	0.40	8.2	9.8
1992	642	9.6	0.38	8.8	10.4	560	9.4	0.40	8.6	10.2
1993	641	9.6	0.38	8.8	10.3	524	8.8	0.39	8.1	9.6
1994	629	9.2	0.37	8.4	9.9	536	8.8	0.38	8.1	9.6
1995	620	8.9	0.36	8.2	9.7	544	8.9	0.39	8.2	9.7
1996	666	9.6	0.38	8.9	10.4	556	9.2	0.39	8.4	10.0
1997	666	9.5	0.37	8.8	10.2	566	9.2	0.39	8.4	9.9
1998	684	9.8	0.38	9.1	10.5	579	9.5	0.40	8.7	10.3
1999	719	10.1	0.38	9.4	10.9	611	9.8	0.40	9.0	10.6
2000	708	9.9	0.38	9.2	10.7	582	9.4	0.39	8.6	10.2
2001	728	10.2	0.38	9.4	11.0	608	9.8	0.40	9.1	10.7
2002	719	10.0	0.38	9.3	10.8	584	9.4	0.39	8.6	10.2
2003	669	9.2	0.36	8.5	9.9	572	9.1	0.39	8.4	9.9
2004	711	9.6	0.37	8.9	10.4	599	9.4	0.39	8.7	10.2
2005	782	10.5	0.38	9.8	11.3	640	10.0	0.40	9.3	10.9
2006	755	10.2	0.38	9.5	11.0	613	9.7	0.40	8.9	10.5
2007	752	10.0	0.37	9.3	10.8	621	9.6	0.39	8.9	10.5
2008	781	10.1	0.37	9.4	10.9	646	9.9	0.40	9.1	10.7
2009	748	9.7	0.36	9.0	10.5	604	9.3	0.39	8.6	10.1
2010	793	10.2	0.37	9.5	11.0	657	10.1	0.40	9.3	10.9
2011	792	10.0	0.36	9.3	10.8	638	9.5	0.39	8.8	10.3
2012	771	9.7	0.36	9.0	10.4	624	9.4	0.38	8.6	10.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Pancreas										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,855	13.5	0.26	13.1	14.1	235	5.8	0.40	5.1	6.7
1986	87	14.7	1.62	11.7	18.2	*	*	*	*	*
1987	91	14.6	1.56	11.7	17.9	*	*	*	*	*
1988	87	13.5	1.47	10.8	16.7	*	*	*	*	*
1989	94	14.8	1.56	11.9	18.2	*	*	*	*	*
1990	96	14.6	1.51	11.8	17.8	*	*	*	*	*
1991	98	14.8	1.52	12.0	18.1	*	*	*	*	*
1992	79	11.1	1.27	8.8	13.9	*	*	*	*	*
1993	111	15.5	1.49	12.8	18.7	*	*	*	*	*
1994	86	12.2	1.33	9.7	15.1	*	*	*	*	*
1995	72	10.3	1.23	8.1	13.0	*	*	*	*	*
1996	100	14.0	1.41	11.3	17.0	10	10.0	3.35	4.5	18.3
1997	96	12.6	1.29	10.2	15.3	*	*	*	*	*
1998	99	12.9	1.30	10.4	15.7	*	*	*	*	*
1999	96	12.3	1.26	9.9	15.0	12	9.3	2.78	4.6	16.1
2000	113	14.3	1.35	11.8	17.2	13	9.2	2.68	4.7	15.8
2001	113	14.2	1.35	11.7	17.1	*	*	*	*	*
2002	129	15.6	1.38	13.0	18.5	*	*	*	*	*
2003	86	10.6	1.15	8.5	13.1	11	5.8	1.83	2.8	10.5
2004	101	12.1	1.22	9.9	14.7	11	6.4	2.00	3.1	11.3
2005	133	15.6	1.36	13.0	18.5	*	*	*	*	*
2006	133	15.2	1.33	12.7	18.0	*	*	*	*	*
2007	119	13.6	1.26	11.3	16.3	12	4.8	1.49	2.4	8.6
2008	126	13.8	1.24	11.5	16.5	*	*	*	*	*
2009	129	13.7	1.22	11.4	16.4	15	6.1	1.64	3.3	10.1
2010	122	13.1	1.20	10.9	15.7	14	4.8	1.37	2.5	8.2
2011	134	13.9	1.21	11.6	16.4	20	7.2	1.67	4.3	11.2
2012	125	12.5	1.14	10.4	15.0	22	7.0	1.56	4.3	10.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Respiratory System										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	185,905	57.2	0.13	57.0	57.5	156,241	55.7	0.14	55.5	56.0
1986	6,263	58.1	0.74	56.6	59.5	5,351	56.0	0.78	54.5	57.5
1987	6,407	59.1	0.75	57.6	60.6	5,429	56.5	0.78	55.0	58.0
1988	6,510	59.7	0.75	58.2	61.1	5,515	57.1	0.78	55.6	58.7
1989	6,615	60.4	0.75	58.9	61.9	5,557	57.5	0.78	56.0	59.1
1990	6,736	61.1	0.75	59.6	62.5	5,664	58.2	0.78	56.7	59.7
1991	6,780	60.8	0.74	59.4	62.3	5,780	58.8	0.78	57.3	60.4
1992	6,844	60.7	0.74	59.2	62.1	5,758	58.0	0.77	56.5	59.5
1993	7,114	62.7	0.75	61.2	64.1	6,005	60.3	0.78	58.8	61.9
1994	7,168	62.6	0.74	61.1	64.1	6,071	60.6	0.78	59.1	62.2
1995	7,203	62.5	0.74	61.0	63.9	6,037	60.0	0.77	58.5	61.5
1996	7,130	61.6	0.73	60.2	63.1	5,916	58.7	0.77	57.2	60.2
1997	6,891	59.1	0.71	57.7	60.6	5,824	57.5	0.75	56.0	59.0
1998	7,003	59.5	0.71	58.2	61.0	5,853	57.4	0.75	55.9	58.8
1999	6,941	58.6	0.70	57.2	59.9	5,838	56.9	0.75	55.5	58.4
2000	7,045	59.1	0.70	57.7	60.5	5,880	57.2	0.75	55.7	58.6
2001	6,999	58.3	0.70	56.9	59.6	5,913	57.2	0.74	55.7	58.7
2002	7,050	58.2	0.69	56.8	59.6	5,911	56.9	0.74	55.4	58.3
2003	7,076	57.8	0.69	56.5	59.2	5,960	56.9	0.74	55.5	58.4
2004	6,979	56.4	0.68	55.1	57.7	5,861	55.5	0.73	54.1	57.0
2005	6,800	54.5	0.66	53.2	55.8	5,682	53.5	0.71	52.1	54.9
2006	6,873	54.4	0.66	53.1	55.7	5,722	53.4	0.71	52.0	54.8
2007	7,108	55.5	0.66	54.2	56.8	5,905	54.5	0.72	53.1	55.9
2008	6,994	53.6	0.65	52.4	54.9	5,874	53.3	0.70	51.9	54.7
2009	6,858	51.9	0.63	50.7	53.1	5,792	52.0	0.69	50.7	53.4
2010	6,836	51.1	0.63	49.8	52.3	5,696	50.6	0.68	49.3	52.0
2011	6,802	49.9	0.61	48.7	51.1	5,693	49.7	0.67	48.4	51.1
2012	6,880	49.4	0.60	48.2	50.6	5,754	49.3	0.66	48.0	50.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Respiratory System										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	28,040	75.4	0.46	74.5	76.3	1,624	21.6	0.58	20.5	22.7
1986	887	78.9	2.73	73.7	84.4	25	25.4	5.94	15.1	39.0
1987	951	83.2	2.77	77.9	88.8	27	24.7	5.42	15.2	37.1
1988	962	83.0	2.75	77.7	88.6	33	26.2	5.08	17.2	37.7
1989	1,031	88.4	2.82	82.9	94.1	27	17.9	3.84	11.2	26.8
1990	1,035	88.1	2.81	82.7	93.8	37	29.0	5.45	19.3	41.2
1991	968	81.6	2.69	76.5	87.1	32	21.4	4.26	13.8	31.0
1992	1,055	86.7	2.73	81.4	92.2	31	21.0	4.29	13.4	30.7
1993	1,079	87.6	2.73	82.3	93.1	30	16.4	3.46	10.3	24.3
1994	1,055	84.0	2.64	78.9	89.3	42	21.8	3.65	15.3	30.0
1995	1,119	86.8	2.65	81.7	92.1	47	22.7	3.70	16.0	30.9
1996	1,167	90.8	2.71	85.6	96.3	47	23.0	3.60	16.5	31.0
1997	1,030	78.6	2.49	73.8	83.6	37	17.0	3.09	11.5	23.9
1998	1,096	80.9	2.49	76.1	86.0	54	20.9	3.10	15.3	27.7
1999	1,054	77.6	2.43	72.9	82.4	49	18.2	2.77	13.2	24.4
2000	1,086	78.3	2.41	73.6	83.2	79	29.4	3.59	22.8	37.1
2001	1,029	74.8	2.37	70.2	79.5	57	20.4	2.87	15.2	26.7
2002	1,069	75.2	2.34	70.7	79.9	70	24.3	3.15	18.5	31.1
2003	1,046	72.0	2.27	67.6	76.6	70	21.1	2.69	16.2	27.0
2004	1,042	70.4	2.22	66.1	74.9	76	22.7	2.80	17.6	28.8
2005	1,041	70.0	2.21	65.8	74.5	77	20.0	2.48	15.5	25.4
2006	1,050	68.7	2.16	64.6	73.1	101	26.1	2.80	20.9	32.1
2007	1,119	72.1	2.20	67.9	76.5	84	22.0	2.56	17.3	27.5
2008	1,023	65.7	2.10	61.7	70.0	97	21.7	2.39	17.3	26.8
2009	979	60.3	1.97	56.5	64.2	87	19.1	2.21	15.0	23.9
2010	1,033	63.1	2.01	59.2	67.2	107	22.8	2.36	18.4	27.8
2011	1,013	60.4	1.94	56.6	64.3	96	18.4	2.00	14.7	22.7
2012	1,021	59.4	1.91	55.7	63.3	105	19.1	1.98	15.4	23.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Respiratory System										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	110,904	81.4	0.25	81.0	81.9	92,517	78.5	0.26	78.0	79.0
1986	4,310	96.8	1.53	93.8	99.8	3,647	92.4	1.59	89.3	95.6
1987	4,305	95.6	1.51	92.6	98.6	3,597	90.2	1.56	87.2	93.3
1988	4,354	96.7	1.52	93.7	99.7	3,687	92.5	1.58	89.4	95.6
1989	4,304	94.3	1.48	91.5	97.3	3,593	89.2	1.53	86.2	92.2
1990	4,351	95.8	1.50	92.8	98.7	3,623	90.4	1.55	87.4	93.5
1991	4,406	96.1	1.49	93.2	99.1	3,719	91.9	1.55	88.9	95.0
1992	4,306	92.2	1.44	89.4	95.1	3,577	86.9	1.49	84.0	89.9
1993	4,463	94.8	1.45	92.0	97.7	3,721	90.2	1.51	87.3	93.2
1994	4,485	94.5	1.44	91.7	97.3	3,778	90.8	1.50	87.9	93.8
1995	4,415	92.3	1.42	89.5	95.1	3,668	87.7	1.47	84.9	90.7
1996	4,294	89.5	1.39	86.8	92.3	3,509	83.7	1.43	81.0	86.6
1997	4,137	85.6	1.35	82.9	88.3	3,453	82.0	1.41	79.3	84.8
1998	4,129	84.5	1.33	81.9	87.2	3,435	81.0	1.40	78.3	83.8
1999	4,097	83.0	1.31	80.5	85.6	3,396	79.4	1.37	76.7	82.1
2000	4,137	83.1	1.31	80.5	85.6	3,428	79.6	1.37	76.9	82.3
2001	4,120	82.4	1.30	79.8	84.9	3,431	79.3	1.36	76.6	82.0
2002	4,033	79.5	1.27	77.1	82.1	3,392	77.7	1.35	75.1	80.4
2003	4,001	77.5	1.24	75.1	79.9	3,338	75.3	1.31	72.7	77.9
2004	3,972	76.1	1.22	73.7	78.5	3,314	74.0	1.30	71.5	76.6
2005	3,813	72.3	1.19	70.0	74.7	3,181	70.4	1.26	68.0	72.9
2006	3,858	72.1	1.18	69.8	74.4	3,192	69.9	1.25	67.4	72.4
2007	3,885	71.1	1.16	68.8	73.4	3,202	68.7	1.23	66.3	71.2
2008	3,850	69.5	1.14	67.2	71.7	3,223	68.1	1.22	65.8	70.6
2009	3,725	65.6	1.10	63.4	67.8	3,135	64.9	1.18	62.6	67.3
2010	3,709	64.4	1.08	62.3	66.5	3,095	63.4	1.16	61.1	65.7
2011	3,716	63.2	1.06	61.2	65.3	3,069	61.7	1.13	59.5	63.9
2012	3,729	61.9	1.04	59.9	64.0	3,114	61.1	1.12	59.0	63.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Respiratory System										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	17,344	115.5	0.92	113.7	117.4	1,043	32.4	1.13	30.3	34.7
1986	641	138.4	5.79	127.3	150.2	22	47.8	12.19	27.0	76.2
1987	688	147.5	5.95	136.1	159.6	20	42.4	11.17	23.5	68.6
1988	644	136.8	5.74	125.8	148.5	23	43.2	10.55	25.1	67.7
1989	694	147.1	5.91	135.7	159.1	17	22.6	6.22	12.1	37.9
1990	702	148.4	5.94	137.0	160.4	26	44.2	9.87	27.0	66.9
1991	663	139.5	5.74	128.4	151.1	24	38.4	9.16	22.6	59.6
1992	708	145.9	5.82	134.7	157.6	21	34.3	8.73	19.4	54.6
1993	723	144.1	5.64	133.2	155.5	19	25.3	6.85	13.7	41.5
1994	679	134.6	5.44	124.2	145.7	28	29.7	6.11	19.0	43.9
1995	723	138.7	5.45	128.3	149.8	24	27.2	6.51	16.0	42.4
1996	756	147.8	5.67	136.9	159.3	29	33.3	6.69	21.5	48.6
1997	656	124.7	5.10	114.9	135.1	28	30.1	6.42	18.8	44.8
1998	661	121.2	4.96	111.7	131.3	33	28.9	5.84	18.6	42.3
1999	664	122.6	4.98	113.1	132.8	37	31.2	5.43	21.5	43.5
2000	649	115.7	4.76	106.6	125.4	60	51.3	7.46	37.7	67.6
2001	650	118.7	4.89	109.3	128.6	39	35.2	6.31	23.9	49.3
2002	604	103.5	4.41	95.0	112.4	37	29.0	5.64	19.0	41.7
2003	610	103.4	4.40	95.0	112.3	53	36.4	5.49	26.4	48.5
2004	610	102.2	4.34	93.9	111.1	48	33.4	5.39	23.6	45.3
2005	580	98.0	4.30	89.8	106.8	52	31.0	4.88	22.2	41.8
2006	597	96.7	4.18	88.7	105.2	69	45.9	6.20	34.6	59.3
2007	634	102.0	4.28	93.8	110.7	49	32.0	4.99	23.0	43.0
2008	578	93.6	4.11	85.7	102.0	49	25.4	4.13	18.0	34.6
2009	531	82.0	3.76	74.8	89.6	59	30.0	4.40	22.0	39.7
2010	554	84.2	3.78	76.9	91.9	60	30.3	4.36	22.4	39.8
2011	591	87.5	3.79	80.2	95.2	56	24.8	3.61	18.3	32.8
2012	554	80.2	3.59	73.3	87.5	61	26.5	3.75	19.7	34.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Respiratory System										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	75,001	40.4	0.15	40.1	40.7	63,724	39.9	0.16	39.6	40.2
1986	1,953	31.9	0.74	30.5	33.4	1,704	31.7	0.79	30.2	33.3
1987	2,102	34.6	0.77	33.1	36.1	1,832	34.1	0.82	32.5	35.7
1988	2,156	34.8	0.76	33.4	36.4	1,828	33.7	0.80	32.1	35.3
1989	2,311	37.2	0.79	35.7	38.8	1,964	36.1	0.83	34.5	37.8
1990	2,385	37.9	0.79	36.4	39.5	2,041	36.9	0.83	35.3	38.6
1991	2,374	37.3	0.78	35.8	38.8	2,061	37.0	0.83	35.4	38.6
1992	2,538	39.4	0.79	37.9	41.0	2,181	38.6	0.84	36.9	40.3
1993	2,651	40.7	0.80	39.2	42.3	2,284	40.2	0.85	38.6	41.9
1994	2,683	40.9	0.80	39.3	42.5	2,293	40.1	0.85	38.5	41.8
1995	2,788	42.1	0.80	40.5	43.7	2,369	41.1	0.86	39.4	42.8
1996	2,836	42.7	0.81	41.1	44.3	2,407	41.9	0.86	40.2	43.6
1997	2,754	41.1	0.79	39.6	42.7	2,371	40.8	0.85	39.2	42.5
1998	2,874	42.5	0.80	41.0	44.1	2,418	41.3	0.85	39.7	43.0
1999	2,844	41.6	0.79	40.1	43.2	2,442	41.4	0.85	39.8	43.1
2000	2,908	42.5	0.79	41.0	44.1	2,452	41.8	0.85	40.1	43.5
2001	2,879	41.7	0.78	40.1	43.2	2,482	42.0	0.85	40.4	43.7
2002	3,017	43.4	0.80	41.9	45.0	2,519	42.5	0.86	40.9	44.3
2003	3,075	43.8	0.80	42.3	45.4	2,622	43.8	0.87	42.1	45.5
2004	3,007	42.6	0.78	41.0	44.1	2,547	42.5	0.85	40.8	44.2
2005	2,987	42.0	0.78	40.5	43.6	2,501	41.6	0.84	40.0	43.3
2006	3,015	42.0	0.77	40.5	43.6	2,530	41.8	0.84	40.2	43.5
2007	3,223	44.5	0.79	43.0	46.1	2,703	44.4	0.87	42.7	46.1
2008	3,144	42.4	0.77	40.9	44.0	2,651	42.6	0.84	41.0	44.3
2009	3,133	42.2	0.76	40.7	43.7	2,657	42.7	0.84	41.1	44.4
2010	3,127	41.6	0.75	40.1	43.1	2,601	41.4	0.83	39.8	43.1
2011	3,086	40.3	0.74	38.8	41.7	2,624	41.1	0.82	39.5	42.7
2012	3,151	40.4	0.73	38.9	41.8	2,640	40.8	0.81	39.2	42.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Respiratory System										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	10,696	48.9	0.48	48.0	49.9	581	13.7	0.60	12.6	14.9
1986	246	37.7	2.44	33.0	42.7	*	*	*	*	*
1987	263	39.6	2.48	34.9	44.7	*	*	*	*	*
1988	318	46.7	2.66	41.6	52.2	10	13.2	4.40	6.1	24.5
1989	337	49.0	2.71	43.8	54.5	10	14.0	4.81	6.2	26.2
1990	333	48.1	2.67	43.0	53.6	11	17.9	6.09	8.0	32.8
1991	305	43.6	2.53	38.8	48.9	*	*	*	*	*
1992	347	48.4	2.62	43.4	53.8	10	11.5	4.07	4.9	21.7
1993	356	49.4	2.65	44.4	54.8	11	10.1	3.49	4.4	18.9
1994	376	50.5	2.63	45.5	55.9	14	15.3	4.34	8.0	25.8
1995	396	52.3	2.65	47.2	57.7	23	19.5	4.43	11.8	29.8
1996	411	53.7	2.67	48.6	59.2	18	15.5	3.87	8.8	24.7
1997	374	48.3	2.52	43.5	53.5	*	*	*	*	*
1998	435	54.5	2.63	49.5	59.9	21	14.8	3.43	8.9	22.9
1999	390	48.2	2.46	43.5	53.3	12	8.2	2.54	4.0	14.5
2000	437	53.4	2.57	48.5	58.7	19	13.3	3.25	7.7	20.9
2001	379	46.3	2.40	41.8	51.3	18	10.7	2.61	6.2	16.9
2002	465	55.9	2.61	50.9	61.2	33	21.1	3.80	14.3	29.6
2003	436	51.0	2.46	46.3	56.0	17	9.5	2.41	5.4	15.3
2004	432	49.4	2.39	44.8	54.3	28	15.0	2.97	9.7	21.8
2005	461	52.2	2.45	47.5	57.2	25	11.9	2.51	7.5	17.8
2006	453	50.4	2.39	45.8	55.3	32	13.2	2.43	8.8	18.8
2007	485	53.1	2.43	48.4	58.0	35	15.1	2.70	10.3	21.2
2008	445	47.9	2.30	43.5	52.6	48	19.1	2.91	13.9	25.6
2009	448	46.6	2.23	42.4	51.2	28	11.3	2.24	7.3	16.4
2010	479	49.7	2.30	45.3	54.5	47	17.8	2.70	12.9	23.8
2011	422	42.6	2.11	38.6	47.0	40	13.4	2.21	9.4	18.4
2012	467	46.0	2.16	41.9	50.4	44	13.9	2.19	9.9	18.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Larynx										
Both Sexes										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	4,793	1.5	0.02	1.4	1.5	3,566	1.3	0.02	1.2	1.3
1986	192	1.8	0.13	1.5	2.1	155	1.6	0.13	1.4	1.9
1987	173	1.6	0.12	1.4	1.9	135	1.4	0.12	1.2	1.7
1988	194	1.8	0.13	1.5	2.0	148	1.5	0.12	1.3	1.8
1989	168	1.6	0.12	1.3	1.8	133	1.4	0.12	1.2	1.7
1990	180	1.7	0.12	1.4	1.9	138	1.4	0.12	1.2	1.7
1991	203	1.8	0.13	1.6	2.1	140	1.4	0.12	1.2	1.7
1992	195	1.7	0.13	1.5	2.0	136	1.4	0.12	1.2	1.6
1993	222	2.0	0.13	1.7	2.3	167	1.7	0.13	1.4	2.0
1994	185	1.6	0.12	1.4	1.9	132	1.3	0.12	1.1	1.6
1995	198	1.7	0.12	1.5	2.0	150	1.5	0.12	1.3	1.8
1996	209	1.8	0.13	1.6	2.1	146	1.5	0.12	1.3	1.7
1997	170	1.5	0.11	1.3	1.7	122	1.2	0.11	1.0	1.5
1998	191	1.6	0.12	1.4	1.9	138	1.4	0.12	1.1	1.6
1999	180	1.5	0.11	1.3	1.8	136	1.3	0.12	1.1	1.6
2000	178	1.5	0.11	1.3	1.7	129	1.3	0.11	1.1	1.5
2001	168	1.4	0.11	1.2	1.6	120	1.2	0.11	1.0	1.4
2002	166	1.4	0.11	1.2	1.6	135	1.3	0.11	1.1	1.6
2003	185	1.5	0.11	1.3	1.7	142	1.4	0.11	1.2	1.6
2004	145	1.2	0.10	1.0	1.4	106	1.0	0.10	0.8	1.2
2005	151	1.2	0.10	1.0	1.4	116	1.1	0.10	0.9	1.3
2006	168	1.3	0.10	1.1	1.5	122	1.1	0.10	1.0	1.4
2007	171	1.3	0.10	1.1	1.5	127	1.1	0.10	1.0	1.4
2008	155	1.2	0.10	1.0	1.4	116	1.1	0.10	0.9	1.3
2009	182	1.3	0.10	1.1	1.5	136	1.2	0.10	1.0	1.4
2010	137	1.0	0.09	0.8	1.2	105	0.9	0.09	0.7	1.1
2011	162	1.2	0.09	1.0	1.4	117	1.0	0.09	0.8	1.2
2012	165	1.2	0.09	1.0	1.4	119	1.0	0.09	0.8	1.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Larynx										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,202	3.1	0.09	2.9	3.3	25	0.3	0.07	0.2	0.5
1986	37	3.3	0.56	2.3	4.6	*	*	*	*	*
1987	37	3.0	0.51	2.1	4.2	*	*	*	*	*
1988	46	3.8	0.57	2.8	5.1	*	*	*	*	*
1989	35	3.0	0.52	2.1	4.2	*	*	*	*	*
1990	42	3.6	0.57	2.6	4.9	*	*	*	*	*
1991	61	4.9	0.64	3.7	6.3	*	*	*	*	*
1992	58	4.6	0.61	3.4	5.9	*	*	*	*	*
1993	55	4.4	0.61	3.3	5.8	*	*	*	*	*
1994	51	4.0	0.57	3.0	5.3	*	*	*	*	*
1995	46	3.4	0.51	2.5	4.6	*	*	*	*	*
1996	63	4.5	0.57	3.5	5.8	*	*	*	*	*
1997	47	3.5	0.52	2.6	4.7	*	*	*	*	*
1998	52	3.7	0.52	2.7	4.8	*	*	*	*	*
1999	43	3.0	0.46	2.1	4.0	*	*	*	*	*
2000	46	3.2	0.48	2.3	4.3	*	*	*	*	*
2001	47	3.4	0.50	2.5	4.5	*	*	*	*	*
2002	31	2.2	0.40	1.5	3.1	*	*	*	*	*
2003	39	2.7	0.45	1.9	3.7	*	*	*	*	*
2004	39	2.6	0.42	1.8	3.5	*	*	*	*	*
2005	34	2.2	0.38	1.5	3.0	*	*	*	*	*
2006	46	2.8	0.41	2.0	3.7	*	*	*	*	*
2007	44	2.7	0.42	2.0	3.6	*	*	*	*	*
2008	38	2.4	0.39	1.7	3.3	*	*	*	*	*
2009	45	2.6	0.40	1.9	3.5	*	*	*	*	*
2010	31	1.8	0.33	1.2	2.5	*	*	*	*	*
2011	45	2.5	0.39	1.8	3.4	*	*	*	*	*
2012	44	2.4	0.37	1.7	3.2	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Larynx										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,810	2.7	0.04	2.6	2.8	2,844	2.4	0.05	2.3	2.5
1986	163	3.7	0.30	3.1	4.3	134	3.4	0.31	2.8	4.1
1987	136	2.9	0.26	2.4	3.5	104	2.6	0.26	2.1	3.1
1988	157	3.4	0.28	2.9	4.0	120	3.0	0.28	2.4	3.6
1989	134	3.0	0.27	2.5	3.6	108	2.7	0.27	2.2	3.3
1990	142	3.1	0.26	2.6	3.6	107	2.6	0.26	2.1	3.1
1991	161	3.5	0.28	3.0	4.1	115	2.9	0.28	2.4	3.5
1992	156	3.3	0.27	2.8	3.8	110	2.7	0.26	2.2	3.2
1993	173	3.6	0.28	3.1	4.2	128	3.1	0.28	2.6	3.7
1994	162	3.4	0.27	2.9	3.9	115	2.7	0.26	2.2	3.3
1995	161	3.4	0.27	2.8	3.9	122	2.9	0.27	2.4	3.5
1996	160	3.2	0.26	2.8	3.8	114	2.7	0.26	2.2	3.2
1997	140	2.8	0.24	2.4	3.3	99	2.3	0.24	1.9	2.8
1998	148	3.0	0.25	2.5	3.5	106	2.5	0.25	2.0	3.0
1999	138	2.7	0.23	2.3	3.2	108	2.5	0.24	2.0	3.0
2000	140	2.8	0.24	2.3	3.3	106	2.5	0.24	2.0	3.0
2001	136	2.7	0.23	2.2	3.2	95	2.2	0.23	1.8	2.7
2002	136	2.6	0.22	2.2	3.1	110	2.4	0.23	2.0	3.0
2003	134	2.5	0.22	2.1	3.0	103	2.2	0.22	1.8	2.7
2004	112	2.1	0.20	1.7	2.5	79	1.7	0.20	1.4	2.2
2005	123	2.3	0.21	1.9	2.7	97	2.1	0.22	1.7	2.6
2006	129	2.3	0.20	1.9	2.7	91	1.9	0.20	1.6	2.4
2007	139	2.5	0.21	2.1	2.9	101	2.1	0.21	1.7	2.6
2008	121	2.1	0.20	1.7	2.5	90	1.9	0.20	1.5	2.3
2009	143	2.3	0.20	2.0	2.8	104	2.0	0.20	1.7	2.5
2010	112	1.9	0.18	1.5	2.3	90	1.8	0.19	1.4	2.2
2011	124	2.0	0.19	1.7	2.4	92	1.8	0.19	1.4	2.2
2012	130	2.1	0.19	1.8	2.5	96	1.9	0.20	1.5	2.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Larynx										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	946	5.9	0.20	5.6	6.4	20	0.6	0.15	0.4	1.0
1986	29	6.5	1.31	4.2	9.6	*	*	*	*	*
1987	31	6.0	1.10	4.0	8.5	*	*	*	*	*
1988	37	7.3	1.24	5.1	10.1	*	*	*	*	*
1989	26	5.6	1.19	3.5	8.3	*	*	*	*	*
1990	35	7.4	1.30	5.1	10.4	*	*	*	*	*
1991	44	8.8	1.39	6.3	11.9	*	*	*	*	*
1992	46	8.6	1.33	6.2	11.6	*	*	*	*	*
1993	45	8.7	1.37	6.3	11.8	*	*	*	*	*
1994	45	8.8	1.39	6.3	11.9	*	*	*	*	*
1995	37	6.7	1.15	4.7	9.3	*	*	*	*	*
1996	46	7.9	1.20	5.7	10.6	*	*	*	*	*
1997	40	7.2	1.18	5.1	9.9	*	*	*	*	*
1998	41	6.9	1.12	4.9	9.4	*	*	*	*	*
1999	30	5.1	0.99	3.4	7.4	*	*	*	*	*
2000	32	5.5	1.00	3.7	7.8	*	*	*	*	*
2001	40	7.1	1.16	5.0	9.7	*	*	*	*	*
2002	26	4.5	0.93	2.8	6.6	*	*	*	*	*
2003	28	4.9	0.98	3.2	7.2	*	*	*	*	*
2004	33	5.1	0.92	3.4	7.2	*	*	*	*	*
2005	25	3.9	0.82	2.5	5.8	*	*	*	*	*
2006	38	5.3	0.91	3.7	7.4	*	*	*	*	*
2007	38	5.7	0.99	3.9	8.0	*	*	*	*	*
2008	30	4.5	0.86	3.0	6.5	*	*	*	*	*
2009	38	5.6	0.96	3.9	7.8	*	*	*	*	*
2010	21	3.1	0.71	1.8	4.8	*	*	*	*	*
2011	32	4.4	0.83	3.0	6.4	*	*	*	*	*
2012	33	4.3	0.79	2.9	6.2	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Larynx										
Females										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	983	0.5	0.02	0.5	0.6	722	0.5	0.02	0.4	0.5
1986	29	0.5	0.08	0.3	0.7	21	0.4	0.08	0.2	0.6
1987	37	0.6	0.10	0.4	0.9	31	0.6	0.11	0.4	0.8
1988	37	0.6	0.10	0.4	0.8	28	0.5	0.10	0.3	0.7
1989	34	0.6	0.10	0.4	0.8	25	0.5	0.10	0.3	0.7
1990	38	0.6	0.10	0.4	0.8	31	0.6	0.10	0.4	0.8
1991	42	0.7	0.11	0.5	0.9	25	0.5	0.10	0.3	0.7
1992	39	0.6	0.10	0.5	0.9	26	0.5	0.09	0.3	0.7
1993	49	0.7	0.11	0.6	1.0	39	0.7	0.11	0.5	0.9
1994	23	0.4	0.08	0.2	0.5	17	0.3	0.07	0.2	0.5
1995	37	0.6	0.09	0.4	0.8	28	0.5	0.10	0.3	0.8
1996	49	0.7	0.11	0.5	1.0	32	0.6	0.10	0.4	0.8
1997	30	0.5	0.08	0.3	0.7	23	0.4	0.09	0.3	0.6
1998	43	0.6	0.10	0.5	0.9	32	0.6	0.10	0.4	0.8
1999	42	0.7	0.10	0.5	0.9	28	0.5	0.10	0.3	0.7
2000	38	0.6	0.10	0.4	0.8	23	0.4	0.09	0.3	0.6
2001	32	0.5	0.08	0.3	0.7	25	0.4	0.09	0.3	0.6
2002	30	0.4	0.08	0.3	0.6	25	0.4	0.08	0.3	0.6
2003	51	0.7	0.10	0.5	1.0	39	0.6	0.10	0.5	0.9
2004	33	0.5	0.08	0.3	0.7	27	0.5	0.09	0.3	0.7
2005	28	0.4	0.08	0.3	0.6	19	0.3	0.07	0.2	0.5
2006	39	0.6	0.09	0.4	0.8	31	0.5	0.10	0.4	0.8
2007	32	0.4	0.08	0.3	0.6	26	0.4	0.09	0.3	0.6
2008	34	0.5	0.08	0.3	0.7	26	0.4	0.08	0.3	0.6
2009	39	0.5	0.08	0.4	0.7	32	0.5	0.09	0.4	0.7
2010	25	0.3	0.07	0.2	0.5	15	0.2	0.06	0.1	0.4
2011	38	0.5	0.08	0.3	0.7	25	0.4	0.08	0.2	0.6
2012	35	0.4	0.08	0.3	0.6	23	0.4	0.08	0.2	0.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Larynx										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	256	1.1	0.07	1.0	1.3	*	*	*	*	*
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	17	2.3	0.56	1.3	3.7	*	*	*	*	*
1992	12	1.7	0.49	0.9	2.9	*	*	*	*	*
1993	10	1.4	0.46	0.7	2.6	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	17	2.1	0.51	1.2	3.3	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	11	1.4	0.42	0.7	2.5	*	*	*	*	*
1999	13	1.5	0.42	0.8	2.6	*	*	*	*	*
2000	14	1.6	0.44	0.9	2.7	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	11	1.3	0.39	0.6	2.3	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	10	1.0	0.30	0.5	1.8	*	*	*	*	*
2011	13	1.3	0.36	0.7	2.2	*	*	*	*	*
2012	11	1.0	0.31	0.5	1.9	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lung and Bronchus										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	179,992	55.4	0.13	55.2	55.7	151,702	54.1	0.14	53.9	54.4
1986	6,001	55.6	0.73	54.2	57.1	5,137	53.7	0.76	52.2	55.2
1987	6,187	57.0	0.73	55.6	58.5	5,250	54.6	0.76	53.1	56.1
1988	6,251	57.3	0.73	55.9	58.7	5,312	55.1	0.76	53.6	56.6
1989	6,403	58.4	0.74	57.0	59.9	5,389	55.8	0.77	54.3	57.3
1990	6,502	58.9	0.74	57.5	60.4	5,478	56.3	0.77	54.8	57.8
1991	6,531	58.6	0.73	57.2	60.0	5,599	57.0	0.77	55.5	58.5
1992	6,604	58.5	0.72	57.1	60.0	5,581	56.2	0.76	54.7	57.7
1993	6,847	60.3	0.73	58.9	61.7	5,801	58.3	0.77	56.8	59.8
1994	6,925	60.5	0.73	59.0	61.9	5,889	58.8	0.77	57.3	60.3
1995	6,965	60.4	0.73	59.0	61.8	5,854	58.1	0.76	56.7	59.7
1996	6,877	59.4	0.72	58.0	60.8	5,733	56.9	0.75	55.4	58.3
1997	6,680	57.3	0.70	56.0	58.7	5,667	55.9	0.74	54.5	57.4
1998	6,774	57.6	0.70	56.2	59.0	5,680	55.7	0.74	54.2	57.1
1999	6,713	56.6	0.69	55.3	58.0	5,657	55.1	0.73	53.7	56.6
2000	6,828	57.2	0.69	55.9	58.6	5,718	55.6	0.74	54.1	57.0
2001	6,793	56.5	0.69	55.2	57.9	5,762	55.7	0.74	54.3	57.2
2002	6,847	56.5	0.68	55.2	57.9	5,743	55.3	0.73	53.8	56.7
2003	6,854	56.0	0.68	54.7	57.4	5,784	55.2	0.73	53.8	56.7
2004	6,804	55.0	0.67	53.7	56.3	5,729	54.3	0.72	52.9	55.7
2005	6,613	53.0	0.65	51.7	54.3	5,536	52.1	0.70	50.7	53.5
2006	6,671	52.8	0.65	51.5	54.1	5,568	51.9	0.70	50.6	53.3
2007	6,902	54.0	0.65	52.7	55.3	5,749	53.1	0.71	51.7	54.5
2008	6,809	52.2	0.64	51.0	53.5	5,732	52.0	0.69	50.6	53.3
2009	6,646	50.3	0.62	49.1	51.6	5,631	50.6	0.68	49.3	51.9
2010	6,671	49.9	0.62	48.7	51.1	5,567	49.5	0.67	48.2	50.9
2011	6,616	48.6	0.61	47.4	49.8	5,555	48.6	0.66	47.3	49.9
2012	6,678	47.9	0.60	46.8	49.1	5,601	48.0	0.65	46.8	49.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lung and Bronchus										
Both Sexes										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	26,694	71.9	0.45	71.1	72.8	1,596	21.2	0.58	20.1	22.4
1986	839	74.7	2.65	69.6	80.0	25	25.4	5.94	15.1	39.0
1987	911	79.9	2.72	74.7	85.4	26	24.1	5.39	14.7	36.5
1988	906	78.4	2.68	73.2	83.8	33	26.2	5.08	17.2	37.7
1989	987	84.5	2.76	79.2	90.1	27	17.9	3.84	11.2	26.8
1990	987	84.1	2.75	78.8	89.6	37	29.0	5.45	19.3	41.2
1991	902	76.2	2.60	71.2	81.5	30	20.0	4.14	12.7	29.5
1992	993	81.8	2.66	76.7	87.2	30	20.5	4.26	13.0	30.2
1993	1,016	82.5	2.65	77.4	87.8	30	16.4	3.46	10.3	24.3
1994	996	79.3	2.56	74.3	84.4	40	20.5	3.51	14.2	28.3
1995	1,066	82.8	2.59	77.8	88.0	45	21.6	3.62	15.1	29.7
1996	1,097	85.7	2.64	80.6	91.0	47	23.0	3.60	16.5	31.0
1997	978	74.7	2.43	70.0	79.6	35	16.2	3.04	10.8	23.0
1998	1,041	77.1	2.43	72.4	82.0	53	20.5	3.08	14.9	27.3
1999	1,008	74.4	2.38	69.8	79.2	48	17.8	2.74	12.9	23.9
2000	1,034	74.7	2.36	70.2	79.5	76	28.4	3.54	21.8	36.0
2001	975	70.9	2.31	66.4	75.5	56	20.2	2.87	15.0	26.5
2002	1,035	72.9	2.30	68.4	77.5	69	24.1	3.14	18.4	30.9
2003	1,004	69.0	2.22	64.7	73.5	66	19.8	2.59	15.0	25.4
2004	999	67.6	2.18	63.4	71.9	76	22.7	2.80	17.6	28.8
2005	1,001	67.5	2.17	63.3	71.8	76	19.8	2.47	15.3	25.2
2006	1,002	65.9	2.12	61.8	70.2	101	26.1	2.80	20.9	32.1
2007	1,069	69.0	2.15	64.8	73.3	84	22.0	2.56	17.3	27.5
2008	982	63.2	2.06	59.2	67.3	95	21.4	2.38	17.0	26.5
2009	929	57.3	1.92	53.6	61.2	86	19.0	2.20	14.9	23.7
2010	998	61.1	1.98	57.3	65.1	106	22.6	2.35	18.2	27.6
2011	965	57.7	1.90	54.0	61.5	96	18.4	2.00	14.7	22.7
2012	974	56.8	1.87	53.2	60.6	103	18.5	1.94	14.9	22.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lung and Bronchus

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	106,379	78.2	0.24	77.7	78.7	89,051	75.6	0.26	75.1	76.1
1986	4,100	92.0	1.49	89.1	94.9	3,474	87.9	1.55	84.9	91.0
1987	4,138	92.0	1.48	89.1	94.9	3,464	87.0	1.53	84.0	90.0
1988	4,161	92.4	1.48	89.5	95.4	3,537	88.7	1.54	85.7	91.8
1989	4,142	90.8	1.46	87.9	93.7	3,464	86.0	1.50	83.1	89.0
1990	4,175	91.9	1.47	89.1	94.9	3,485	87.0	1.52	84.1	90.1
1991	4,221	92.1	1.46	89.2	95.0	3,582	88.5	1.52	85.5	91.5
1992	4,123	88.3	1.41	85.6	91.1	3,442	83.7	1.46	80.8	86.6
1993	4,265	90.7	1.42	87.9	93.5	3,572	86.6	1.48	83.7	89.6
1994	4,286	90.3	1.41	87.6	93.1	3,633	87.4	1.48	84.6	90.4
1995	4,228	88.4	1.39	85.7	91.1	3,525	84.3	1.44	81.5	87.2
1996	4,104	85.7	1.36	83.0	88.4	3,372	80.5	1.41	77.8	83.3
1997	3,970	82.2	1.32	79.6	84.9	3,331	79.2	1.39	76.5	81.9
1998	3,956	81.1	1.31	78.5	83.7	3,305	77.9	1.37	75.3	80.7
1999	3,936	79.9	1.29	77.3	82.4	3,267	76.4	1.35	73.8	79.1
2000	3,971	79.8	1.28	77.3	82.3	3,298	76.6	1.34	74.0	79.3
2001	3,960	79.2	1.27	76.7	81.8	3,316	76.6	1.34	74.0	79.3
2002	3,864	76.3	1.24	73.9	78.8	3,253	74.6	1.32	72.1	77.3
2003	3,843	74.5	1.22	72.1	76.9	3,212	72.5	1.29	70.0	75.1
2004	3,840	73.6	1.20	71.2	76.0	3,217	71.8	1.28	69.4	74.4
2005	3,664	69.5	1.16	67.2	71.8	3,063	67.8	1.24	65.4	70.3
2006	3,706	69.4	1.16	67.2	71.7	3,080	67.5	1.23	65.1	70.0
2007	3,723	68.2	1.14	66.0	70.5	3,081	66.2	1.21	63.8	68.6
2008	3,708	66.9	1.12	64.8	69.2	3,115	65.9	1.20	63.5	68.3
2009	3,563	62.9	1.07	60.8	65.1	3,015	62.6	1.16	60.3	64.9
2010	3,575	62.1	1.06	60.1	64.2	2,986	61.2	1.14	59.0	63.5
2011	3,581	61.0	1.04	59.0	63.1	2,966	59.7	1.12	57.5	61.9
2012	3,576	59.4	1.02	57.4	61.4	2,996	58.8	1.10	56.7	61.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lung and Bronchus

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	16,307	109.0	0.90	107.2	110.8	1,021	31.8	1.12	29.6	34.1
1986	604	130.4	5.62	119.6	141.8	22	47.8	12.19	27.0	76.2
1987	655	141.2	5.84	129.9	153.0	19	41.2	11.10	22.4	67.3
1988	601	128.2	5.57	117.5	139.5	23	43.2	10.55	25.1	67.7
1989	661	140.1	5.76	129.0	151.8	17	22.6	6.22	12.1	37.9
1990	664	140.3	5.78	129.2	152.1	26	44.2	9.87	27.0	66.9
1991	617	130.4	5.56	119.7	141.7	22	35.3	8.83	20.1	55.8
1992	660	136.9	5.66	126.0	148.4	21	34.3	8.73	19.4	54.6
1993	674	134.3	5.45	123.9	145.4	19	25.3	6.85	13.7	41.5
1994	627	124.2	5.21	114.2	134.8	26	26.3	5.58	16.6	39.5
1995	681	130.9	5.30	120.8	141.7	22	24.8	6.25	14.1	39.5
1996	703	138.4	5.51	127.8	149.6	29	33.3	6.69	21.5	48.6
1997	613	117.1	4.96	107.6	127.1	26	28.2	6.29	17.3	42.7
1998	619	114.2	4.83	104.9	124.0	32	28.1	5.79	17.9	41.4
1999	632	117.2	4.88	107.9	127.2	37	31.2	5.43	21.5	43.5
2000	615	110.1	4.65	101.2	119.5	58	49.7	7.37	36.3	65.8
2001	606	110.9	4.74	101.8	120.5	38	34.8	6.29	23.5	48.8
2002	575	98.6	4.31	90.3	107.4	36	28.7	5.64	18.8	41.4
2003	581	98.3	4.28	90.1	107.0	50	33.6	5.22	24.1	45.1
2004	575	96.9	4.24	88.7	105.5	48	33.4	5.39	23.6	45.3
2005	550	93.3	4.20	85.2	101.8	51	30.5	4.86	21.8	41.3
2006	557	91.1	4.07	83.3	99.4	69	45.9	6.20	34.6	59.3
2007	593	95.8	4.15	87.8	104.2	49	32.0	4.99	23.0	43.0
2008	545	88.6	4.01	80.9	96.8	48	25.1	4.12	17.7	34.3
2009	490	76.0	3.63	69.1	83.4	58	29.7	4.39	21.7	39.3
2010	530	80.8	3.71	73.6	88.3	59	29.9	4.34	22.0	39.4
2011	559	83.0	3.70	75.9	90.6	56	24.8	3.61	18.3	32.8
2012	520	75.7	3.50	69.0	82.9	60	25.7	3.67	19.1	33.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lung and Bronchus										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	73,613	39.6	0.15	39.3	39.9	62,651	39.2	0.16	38.9	39.6
1986	1,901	31.1	0.73	29.7	32.5	1,663	30.9	0.78	29.4	32.5
1987	2,049	33.7	0.76	32.2	35.2	1,786	33.2	0.81	31.7	34.8
1988	2,090	33.8	0.75	32.3	35.3	1,775	32.7	0.79	31.2	34.3
1989	2,261	36.4	0.78	34.9	38.0	1,925	35.4	0.82	33.8	37.0
1990	2,327	37.0	0.78	35.5	38.5	1,993	36.0	0.82	34.4	37.7
1991	2,310	36.3	0.76	34.8	37.8	2,017	36.2	0.82	34.6	37.8
1992	2,481	38.5	0.78	37.0	40.1	2,139	37.8	0.83	36.2	39.5
1993	2,582	39.7	0.79	38.2	41.3	2,229	39.3	0.84	37.7	41.0
1994	2,639	40.2	0.79	38.7	41.8	2,256	39.5	0.84	37.8	41.2
1995	2,737	41.3	0.80	39.8	42.9	2,329	40.4	0.85	38.7	42.1
1996	2,773	41.7	0.80	40.2	43.3	2,361	41.1	0.86	39.4	42.8
1997	2,710	40.4	0.78	38.9	42.0	2,336	40.2	0.84	38.6	41.9
1998	2,818	41.7	0.79	40.2	43.3	2,375	40.6	0.84	39.0	42.3
1999	2,777	40.6	0.78	39.1	42.2	2,390	40.5	0.84	38.9	42.2
2000	2,857	41.7	0.79	40.2	43.3	2,420	41.2	0.85	39.6	42.9
2001	2,833	41.0	0.78	39.5	42.5	2,446	41.4	0.85	39.8	43.1
2002	2,983	42.9	0.79	41.4	44.5	2,490	42.1	0.85	40.4	43.8
2003	3,011	42.9	0.79	41.4	44.5	2,572	43.0	0.86	41.3	44.7
2004	2,964	42.0	0.78	40.4	43.5	2,512	41.9	0.85	40.2	43.6
2005	2,949	41.5	0.77	40.0	43.0	2,473	41.1	0.84	39.5	42.8
2006	2,965	41.3	0.77	39.8	42.9	2,488	41.1	0.84	39.5	42.8
2007	3,179	44.0	0.79	42.4	45.5	2,668	43.9	0.86	42.2	45.6
2008	3,101	41.8	0.76	40.4	43.4	2,617	42.1	0.84	40.4	43.7
2009	3,083	41.5	0.76	40.0	43.0	2,616	42.0	0.83	40.4	43.7
2010	3,096	41.2	0.75	39.7	42.7	2,581	41.1	0.82	39.5	42.8
2011	3,035	39.6	0.73	38.2	41.1	2,589	40.6	0.81	39.0	42.2
2012	3,102	39.8	0.73	38.3	41.2	2,605	40.2	0.80	38.7	41.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lung and Bronchus

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	10,387	47.5	0.47	46.6	48.5	575	13.6	0.60	12.4	14.8
1986	235	36.0	2.39	31.4	40.9	*	*	*	*	*
1987	256	38.6	2.44	33.9	43.7	*	*	*	*	*
1988	305	44.9	2.61	39.9	50.3	10	13.2	4.40	6.1	24.5
1989	326	47.2	2.66	42.2	52.7	10	14.0	4.81	6.2	26.2
1990	323	46.7	2.64	41.7	52.2	11	17.9	6.09	8.0	32.8
1991	285	40.8	2.44	36.1	45.9	*	*	*	*	*
1992	333	46.4	2.57	41.5	51.7	*	*	*	*	*
1993	342	47.5	2.59	42.5	52.8	11	10.1	3.49	4.4	18.9
1994	369	49.5	2.60	44.5	54.9	14	15.3	4.34	8.0	25.8
1995	385	50.9	2.62	45.9	56.2	23	19.5	4.43	11.8	29.8
1996	394	51.7	2.63	46.7	57.1	18	15.5	3.87	8.8	24.7
1997	365	47.1	2.48	42.4	52.2	*	*	*	*	*
1998	422	52.9	2.59	48.0	58.2	21	14.8	3.43	8.9	22.9
1999	376	46.6	2.42	41.9	51.5	11	7.4	2.44	3.5	13.6
2000	419	51.3	2.52	46.5	56.5	18	12.7	3.20	7.2	20.2
2001	369	45.1	2.36	40.6	49.9	18	10.7	2.61	6.2	16.9
2002	460	55.2	2.59	50.3	60.6	33	21.1	3.80	14.3	29.6
2003	423	49.5	2.42	44.8	54.4	16	9.1	2.37	5.0	14.8
2004	424	48.4	2.37	43.8	53.2	28	15.0	2.97	9.7	21.8
2005	451	51.1	2.43	46.4	56.1	25	11.9	2.51	7.5	17.8
2006	445	49.5	2.37	45.0	54.3	32	13.2	2.43	8.8	18.8
2007	476	52.1	2.41	47.5	57.0	35	15.1	2.70	10.3	21.2
2008	437	47.0	2.27	42.7	51.7	47	18.9	2.90	13.6	25.3
2009	439	45.7	2.21	41.5	50.3	28	11.3	2.24	7.3	16.4
2010	468	48.7	2.28	44.3	53.3	47	17.8	2.70	12.9	23.8
2011	406	41.1	2.07	37.1	45.3	40	13.4	2.21	9.4	18.4
2012	454	44.7	2.13	40.7	49.1	43	13.4	2.14	9.5	18.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Skin excluding Basal and Squamous

Both Sexes

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	10,043	3.1	0.03	3.1	3.2	9,613	3.5	0.04	3.4	3.5
1986	298	2.9	0.17	2.6	3.3	283	3.1	0.19	2.8	3.5
1987	304	2.9	0.17	2.6	3.3	286	3.1	0.19	2.8	3.5
1988	330	3.2	0.18	2.8	3.5	311	3.4	0.19	3.0	3.8
1989	306	2.9	0.17	2.6	3.2	286	3.1	0.18	2.7	3.5
1990	341	3.2	0.17	2.9	3.5	317	3.4	0.19	3.0	3.8
1991	329	3.0	0.17	2.7	3.4	311	3.3	0.19	2.9	3.7
1992	336	3.0	0.17	2.7	3.4	320	3.3	0.19	3.0	3.7
1993	370	3.3	0.17	3.0	3.7	356	3.7	0.20	3.3	4.1
1994	346	3.0	0.16	2.7	3.4	331	3.3	0.18	3.0	3.7
1995	350	3.1	0.16	2.8	3.4	337	3.4	0.19	3.1	3.8
1996	358	3.1	0.16	2.8	3.5	341	3.4	0.19	3.1	3.8
1997	358	3.1	0.16	2.8	3.4	341	3.4	0.18	3.0	3.8
1998	368	3.1	0.16	2.8	3.5	354	3.5	0.19	3.2	3.9
1999	366	3.1	0.16	2.8	3.4	358	3.5	0.18	3.1	3.9
2000	373	3.1	0.16	2.8	3.4	356	3.4	0.18	3.1	3.8
2001	341	2.8	0.15	2.5	3.1	329	3.2	0.17	2.8	3.5
2002	408	3.3	0.17	3.0	3.7	389	3.7	0.19	3.4	4.1
2003	387	3.1	0.16	2.8	3.4	374	3.5	0.18	3.2	3.9
2004	408	3.3	0.16	3.0	3.6	396	3.8	0.19	3.4	4.1
2005	433	3.4	0.16	3.1	3.7	412	3.8	0.19	3.5	4.2
2006	412	3.2	0.16	2.9	3.5	390	3.6	0.18	3.2	4.0
2007	419	3.2	0.16	2.9	3.5	402	3.6	0.18	3.3	4.0
2008	390	3.0	0.15	2.7	3.3	377	3.4	0.18	3.1	3.8
2009	446	3.4	0.16	3.1	3.7	431	3.9	0.19	3.5	4.3
2010	402	3.0	0.15	2.7	3.3	394	3.4	0.18	3.1	3.8
2011	413	3.0	0.15	2.7	3.3	397	3.5	0.18	3.1	3.8
2012	451	3.3	0.16	3.0	3.6	434	3.8	0.18	3.4	4.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Skin excluding Basal and Squamous

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	385	1.0	0.05	0.9	1.1	45	0.5	0.09	0.4	0.7
1986	14	1.3	0.34	0.7	2.1	*	*	*	*	*
1987	17	1.5	0.38	0.9	2.5	*	*	*	*	*
1988	19	1.7	0.39	1.0	2.6	*	*	*	*	*
1989	20	1.7	0.39	1.0	2.6	*	*	*	*	*
1990	23	1.9	0.40	1.2	2.8	*	*	*	*	*
1991	18	1.5	0.36	0.9	2.3	*	*	*	*	*
1992	15	1.2	0.31	0.6	1.9	*	*	*	*	*
1993	12	0.9	0.27	0.5	1.6	*	*	*	*	*
1994	15	1.3	0.34	0.7	2.1	*	*	*	*	*
1995	11	0.8	0.26	0.4	1.5	*	*	*	*	*
1996	15	1.0	0.26	0.5	1.6	*	*	*	*	*
1997	15	1.1	0.29	0.6	1.8	*	*	*	*	*
1998	12	0.8	0.25	0.4	1.5	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	14	1.0	0.26	0.5	1.6	*	*	*	*	*
2001	10	0.8	0.25	0.4	1.4	*	*	*	*	*
2002	19	1.3	0.31	0.8	2.1	*	*	*	*	*
2003	12	0.9	0.26	0.4	1.5	*	*	*	*	*
2004	11	0.8	0.23	0.4	1.4	*	*	*	*	*
2005	16	1.0	0.26	0.6	1.7	*	*	*	*	*
2006	18	1.2	0.29	0.7	1.9	*	*	*	*	*
2007	14	0.9	0.24	0.5	1.5	*	*	*	*	*
2008	11	0.7	0.21	0.3	1.2	*	*	*	*	*
2009	13	0.8	0.22	0.4	1.3	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	15	0.9	0.23	0.5	1.5	*	*	*	*	*
2012	13	0.8	0.23	0.4	1.4	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Skin excluding Basal and Squamous

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	6,309	4.7	0.06	4.6	4.8	6,045	5.2	0.07	5.0	5.3
1986	175	3.9	0.31	3.3	4.5	166	4.2	0.34	3.5	4.9
1987	191	4.4	0.33	3.8	5.1	178	4.7	0.36	4.0	5.4
1988	205	4.6	0.34	4.0	5.3	190	4.9	0.37	4.2	5.6
1989	184	4.2	0.33	3.6	4.9	171	4.5	0.36	3.8	5.2
1990	202	4.5	0.33	3.9	5.2	181	4.6	0.35	3.9	5.3
1991	205	4.4	0.31	3.8	5.0	190	4.6	0.35	4.0	5.4
1992	215	4.7	0.33	4.1	5.4	204	5.1	0.37	4.4	5.9
1993	229	4.8	0.33	4.2	5.5	221	5.3	0.37	4.6	6.1
1994	215	4.6	0.32	4.0	5.3	205	5.0	0.36	4.3	5.8
1995	215	4.6	0.32	4.0	5.2	207	5.0	0.36	4.4	5.8
1996	219	4.4	0.31	3.9	5.1	208	4.9	0.34	4.2	5.6
1997	227	4.6	0.31	4.0	5.2	216	5.0	0.35	4.4	5.8
1998	237	4.8	0.32	4.2	5.5	229	5.4	0.36	4.7	6.1
1999	233	4.8	0.32	4.2	5.4	228	5.4	0.36	4.7	6.1
2000	224	4.5	0.30	3.9	5.1	215	5.0	0.34	4.3	5.7
2001	202	4.1	0.29	3.5	4.7	196	4.5	0.33	3.9	5.2
2002	264	5.1	0.32	4.5	5.8	251	5.7	0.36	5.0	6.4
2003	247	4.8	0.31	4.2	5.4	243	5.4	0.35	4.8	6.2
2004	253	4.7	0.30	4.1	5.3	246	5.3	0.34	4.7	6.1
2005	268	5.0	0.31	4.5	5.7	256	5.6	0.35	4.9	6.3
2006	263	4.8	0.30	4.2	5.4	251	5.3	0.34	4.7	6.1
2007	265	4.9	0.31	4.3	5.5	255	5.5	0.35	4.8	6.2
2008	267	4.9	0.30	4.3	5.5	264	5.6	0.35	4.9	6.3
2009	283	5.0	0.30	4.5	5.7	274	5.7	0.35	5.0	6.4
2010	262	4.6	0.29	4.1	5.2	260	5.3	0.34	4.7	6.0
2011	278	4.8	0.30	4.3	5.5	269	5.5	0.34	4.8	6.2
2012	281	4.8	0.29	4.2	5.4	271	5.4	0.33	4.8	6.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Skin excluding Basal and Squamous

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	244	1.5	0.10	1.3	1.7	20	0.4	0.11	0.3	0.7
1986	*	*	*	*	*	*	*	*	*	*
1987	13	2.7	0.76	1.4	4.6	*	*	*	*	*
1988	15	2.9	0.78	1.6	4.9	*	*	*	*	*
1989	13	2.5	0.73	1.2	4.3	*	*	*	*	*
1990	21	3.9	0.89	2.4	6.1	*	*	*	*	*
1991	15	2.9	0.82	1.5	4.9	*	*	*	*	*
1992	11	1.9	0.59	0.9	3.4	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	10	2.2	0.72	1.0	4.0	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	13	2.2	0.66	1.1	3.9	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	11	1.7	0.57	0.8	3.2	*	*	*	*	*
2006	12	1.9	0.60	0.9	3.4	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Skin excluding Basal and Squamous

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,734	2.0	0.03	2.0	2.1	3,568	2.3	0.04	2.2	2.3
1986	123	2.2	0.20	1.8	2.6	117	2.3	0.22	1.9	2.8
1987	113	1.9	0.18	1.6	2.3	108	2.1	0.20	1.7	2.5
1988	125	2.1	0.19	1.7	2.5	121	2.3	0.22	1.9	2.8
1989	122	2.0	0.18	1.6	2.4	115	2.2	0.21	1.8	2.6
1990	139	2.3	0.20	1.9	2.7	136	2.6	0.22	2.1	3.0
1991	124	2.0	0.18	1.6	2.3	121	2.2	0.20	1.8	2.6
1992	121	1.9	0.18	1.6	2.3	116	2.1	0.20	1.8	2.6
1993	141	2.2	0.19	1.9	2.6	135	2.5	0.22	2.1	2.9
1994	131	2.0	0.17	1.7	2.4	126	2.2	0.20	1.8	2.6
1995	135	2.0	0.18	1.7	2.4	130	2.3	0.20	1.9	2.7
1996	139	2.1	0.18	1.8	2.5	133	2.3	0.21	2.0	2.8
1997	131	1.9	0.17	1.6	2.3	125	2.1	0.19	1.8	2.6
1998	131	2.0	0.17	1.7	2.3	125	2.2	0.20	1.8	2.7
1999	133	1.9	0.17	1.6	2.3	130	2.2	0.20	1.8	2.6
2000	149	2.1	0.18	1.8	2.5	141	2.4	0.20	2.0	2.8
2001	139	2.0	0.17	1.7	2.3	133	2.2	0.20	1.9	2.7
2002	144	2.0	0.17	1.7	2.4	138	2.3	0.20	1.9	2.7
2003	140	1.9	0.16	1.6	2.3	131	2.1	0.19	1.8	2.6
2004	155	2.2	0.18	1.9	2.6	150	2.5	0.21	2.1	3.0
2005	165	2.2	0.18	1.9	2.6	156	2.5	0.20	2.1	2.9
2006	149	2.0	0.17	1.7	2.3	139	2.2	0.19	1.9	2.6
2007	154	2.1	0.17	1.8	2.5	147	2.4	0.20	2.0	2.9
2008	123	1.6	0.15	1.3	1.9	113	1.8	0.17	1.5	2.2
2009	163	2.2	0.18	1.9	2.6	157	2.6	0.21	2.2	3.0
2010	140	1.8	0.16	1.5	2.1	134	2.1	0.19	1.7	2.5
2011	135	1.8	0.16	1.5	2.1	128	2.0	0.19	1.7	2.4
2012	170	2.2	0.17	1.9	2.5	163	2.5	0.20	2.1	3.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Skin excluding Basal and Squamous

Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	141	0.7	0.06	0.6	0.8	25	0.6	0.12	0.3	0.8
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Melanoma of the Skin										
Both Sexes										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	7,716	2.4	0.03	2.3	2.4	7,540	2.7	0.03	2.7	2.8
1986	216	2.1	0.15	1.8	2.4	212	2.4	0.16	2.0	2.7
1987	212	2.1	0.14	1.8	2.4	207	2.3	0.16	2.0	2.6
1988	261	2.5	0.16	2.2	2.8	256	2.8	0.18	2.5	3.2
1989	220	2.1	0.14	1.8	2.4	214	2.3	0.16	2.0	2.7
1990	253	2.4	0.15	2.1	2.7	248	2.7	0.17	2.3	3.0
1991	241	2.2	0.14	1.9	2.5	236	2.5	0.16	2.2	2.8
1992	266	2.4	0.15	2.1	2.7	261	2.7	0.17	2.4	3.1
1993	282	2.5	0.15	2.2	2.8	276	2.9	0.17	2.6	3.2
1994	267	2.4	0.14	2.1	2.6	263	2.7	0.16	2.4	3.0
1995	263	2.3	0.14	2.0	2.6	257	2.6	0.16	2.3	3.0
1996	282	2.4	0.15	2.2	2.8	276	2.8	0.17	2.5	3.1
1997	270	2.3	0.14	2.0	2.6	264	2.6	0.16	2.3	3.0
1998	291	2.5	0.15	2.2	2.8	287	2.9	0.17	2.5	3.2
1999	281	2.4	0.14	2.1	2.7	276	2.7	0.16	2.4	3.0
2000	280	2.3	0.14	2.1	2.6	273	2.7	0.16	2.4	3.0
2001	259	2.1	0.13	1.9	2.4	254	2.5	0.15	2.2	2.8
2002	304	2.5	0.14	2.2	2.8	296	2.9	0.17	2.5	3.2
2003	285	2.3	0.14	2.0	2.6	277	2.6	0.16	2.3	3.0
2004	334	2.7	0.15	2.4	3.0	327	3.1	0.17	2.8	3.5
2005	336	2.7	0.14	2.4	3.0	326	3.0	0.17	2.7	3.4
2006	333	2.6	0.14	2.3	2.9	319	3.0	0.17	2.6	3.3
2007	338	2.6	0.14	2.3	2.9	332	3.0	0.17	2.7	3.4
2008	297	2.3	0.13	2.0	2.5	290	2.6	0.16	2.3	3.0
2009	359	2.7	0.15	2.4	3.0	350	3.2	0.17	2.8	3.5
2010	308	2.3	0.13	2.0	2.5	303	2.7	0.16	2.4	3.0
2011	322	2.4	0.13	2.1	2.7	314	2.8	0.16	2.5	3.1
2012	356	2.6	0.14	2.3	2.9	346	3.0	0.16	2.7	3.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Melanoma of the Skin										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	142	0.4	0.03	0.3	0.5	34	0.4	0.07	0.3	0.5
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	10	0.7	0.22	0.3	1.2	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.
*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*
** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Melanoma of the Skin										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	4,699	3.4	0.05	3.3	3.5	4,627	3.9	0.06	3.8	4.0
1986	118	2.6	0.25	2.1	3.1	116	2.9	0.28	2.4	3.5
1987	123	2.8	0.26	2.3	3.3	122	3.1	0.29	2.6	3.8
1988	156	3.5	0.29	3.0	4.1	153	3.9	0.33	3.3	4.6
1989	129	2.9	0.27	2.4	3.5	124	3.2	0.30	2.6	3.8
1990	139	3.1	0.27	2.5	3.6	136	3.4	0.30	2.8	4.0
1991	144	3.0	0.26	2.5	3.5	142	3.4	0.29	2.8	4.0
1992	165	3.6	0.29	3.1	4.2	163	4.1	0.33	3.4	4.8
1993	166	3.5	0.28	2.9	4.0	165	4.0	0.31	3.4	4.6
1994	161	3.4	0.28	2.9	4.0	159	3.9	0.31	3.3	4.5
1995	155	3.1	0.26	2.7	3.7	153	3.6	0.29	3.0	4.2
1996	173	3.5	0.27	3.0	4.0	171	3.9	0.31	3.4	4.6
1997	159	3.1	0.25	2.6	3.6	158	3.6	0.29	3.0	4.2
1998	189	3.8	0.28	3.2	4.4	187	4.3	0.32	3.7	5.0
1999	172	3.4	0.26	2.9	4.0	169	3.9	0.30	3.3	4.5
2000	166	3.3	0.26	2.8	3.8	165	3.8	0.29	3.2	4.4
2001	155	3.0	0.25	2.6	3.6	152	3.5	0.28	2.9	4.1
2002	186	3.6	0.26	3.1	4.1	182	4.0	0.30	3.5	4.7
2003	176	3.3	0.25	2.8	3.9	173	3.8	0.29	3.2	4.4
2004	205	3.8	0.27	3.3	4.3	202	4.3	0.31	3.8	5.0
2005	203	3.8	0.27	3.3	4.4	199	4.3	0.31	3.7	5.0
2006	204	3.7	0.26	3.2	4.2	198	4.2	0.30	3.6	4.8
2007	207	3.8	0.27	3.3	4.3	204	4.3	0.31	3.8	5.0
2008	205	3.7	0.26	3.2	4.2	205	4.3	0.31	3.7	5.0
2009	227	4.0	0.27	3.5	4.5	223	4.6	0.31	4.0	5.2
2010	190	3.3	0.24	2.8	3.8	189	3.9	0.28	3.3	4.5
2011	210	3.6	0.25	3.1	4.1	206	4.1	0.29	3.6	4.7
2012	216	3.6	0.25	3.1	4.1	211	4.1	0.29	3.6	4.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Melanoma of the Skin

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	59	0.4	0.05	0.3	0.5	13	0.3	0.08	0.1	0.5
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Melanoma of the Skin										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,017	1.7	0.03	1.6	1.7	2,913	1.9	0.04	1.8	2.0
1986	98	1.7	0.18	1.4	2.1	96	1.9	0.20	1.6	2.4
1987	89	1.5	0.16	1.2	1.9	85	1.7	0.19	1.3	2.1
1988	105	1.8	0.18	1.4	2.2	103	2.0	0.20	1.6	2.5
1989	91	1.5	0.16	1.2	1.9	90	1.8	0.19	1.4	2.2
1990	114	1.9	0.18	1.6	2.3	112	2.2	0.21	1.8	2.6
1991	97	1.5	0.16	1.3	1.9	94	1.7	0.18	1.4	2.1
1992	101	1.6	0.16	1.3	2.0	98	1.8	0.19	1.5	2.2
1993	116	1.9	0.17	1.5	2.2	111	2.1	0.20	1.7	2.5
1994	106	1.6	0.16	1.3	2.0	104	1.8	0.18	1.5	2.2
1995	108	1.7	0.16	1.4	2.0	104	1.9	0.19	1.5	2.3
1996	109	1.7	0.16	1.4	2.0	105	1.9	0.18	1.5	2.3
1997	111	1.6	0.16	1.3	2.0	106	1.8	0.18	1.5	2.2
1998	102	1.6	0.16	1.3	1.9	100	1.8	0.18	1.5	2.2
1999	109	1.6	0.16	1.3	2.0	107	1.9	0.18	1.5	2.3
2000	114	1.7	0.16	1.4	2.0	108	1.9	0.18	1.5	2.3
2001	104	1.5	0.15	1.2	1.8	102	1.8	0.18	1.5	2.2
2002	118	1.7	0.16	1.4	2.0	114	2.0	0.19	1.6	2.4
2003	109	1.5	0.15	1.3	1.9	104	1.8	0.17	1.4	2.1
2004	129	1.8	0.16	1.5	2.2	125	2.1	0.19	1.8	2.5
2005	133	1.8	0.16	1.5	2.2	127	2.1	0.19	1.8	2.5
2006	129	1.7	0.16	1.5	2.1	121	2.0	0.18	1.6	2.4
2007	131	1.8	0.16	1.5	2.2	128	2.2	0.19	1.8	2.6
2008	92	1.3	0.13	1.0	1.5	85	1.4	0.15	1.1	1.7
2009	132	1.8	0.16	1.5	2.2	127	2.2	0.20	1.8	2.6
2010	118	1.5	0.14	1.3	1.8	114	1.8	0.17	1.5	2.2
2011	112	1.5	0.14	1.2	1.8	108	1.8	0.17	1.4	2.1
2012	140	1.8	0.16	1.5	2.2	135	2.1	0.19	1.8	2.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Melanoma of the Skin

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	83	0.4	0.04	0.3	0.5	21	0.4	0.10	0.3	0.7
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Breast										
Both Sexes										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	53,843	16.7	0.07	16.5	16.8	44,763	16.1	0.08	16.0	16.3
1986	2,114	20.5	0.45	19.7	21.4	1,831	20.1	0.48	19.2	21.1
1987	2,101	20.3	0.45	19.4	21.2	1,824	19.9	0.47	19.0	20.8
1988	2,108	20.1	0.44	19.3	21.0	1,840	20.0	0.47	19.1	20.9
1989	2,172	20.6	0.45	19.8	21.5	1,901	20.5	0.48	19.6	21.5
1990	2,194	20.7	0.45	19.9	21.6	1,903	20.4	0.47	19.5	21.4
1991	2,284	21.1	0.45	20.3	22.0	1,978	20.9	0.47	19.9	21.8
1992	2,164	19.8	0.43	18.9	20.6	1,890	19.7	0.46	18.8	20.6
1993	2,181	19.7	0.42	18.9	20.5	1,850	19.1	0.45	18.3	20.0
1994	2,156	19.2	0.42	18.4	20.0	1,814	18.6	0.44	17.7	19.4
1995	2,149	18.9	0.41	18.1	19.8	1,818	18.4	0.43	17.6	19.3
1996	2,121	18.5	0.40	17.7	19.3	1,802	18.1	0.43	17.3	19.0
1997	2,053	17.7	0.39	17.0	18.5	1,685	16.8	0.41	16.0	17.6
1998	1,997	17.0	0.38	16.2	17.7	1,681	16.5	0.40	15.8	17.4
1999	2,030	17.1	0.38	16.3	17.8	1,648	16.0	0.40	15.3	16.8
2000	2,067	17.2	0.38	16.4	17.9	1,701	16.4	0.40	15.6	17.2
2001	1,869	15.4	0.36	14.7	16.1	1,527	14.6	0.37	13.9	15.4
2002	1,916	15.6	0.36	14.9	16.3	1,545	14.7	0.37	14.0	15.4
2003	1,887	15.1	0.35	14.5	15.8	1,509	14.2	0.37	13.5	14.9
2004	1,851	14.6	0.34	14.0	15.3	1,457	13.5	0.36	12.8	14.2
2005	1,883	14.8	0.34	14.1	15.5	1,509	13.9	0.36	13.2	14.6
2006	1,785	13.8	0.33	13.1	14.4	1,442	13.1	0.35	12.4	13.8
2007	1,825	13.9	0.33	13.3	14.6	1,454	13.1	0.35	12.4	13.8
2008	1,794	13.4	0.32	12.8	14.1	1,436	12.7	0.34	12.0	13.4
2009	1,731	12.8	0.31	12.2	13.4	1,402	12.3	0.33	11.7	13.0
2010	1,755	12.8	0.31	12.2	13.4	1,394	12.0	0.33	11.4	12.7
2011	1,829	13.1	0.31	12.5	13.7	1,474	12.5	0.33	11.9	13.2
2012	1,827	12.8	0.31	12.3	13.5	1,448	12.2	0.33	11.5	12.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Breast										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	8,522	22.5	0.25	22.1	23.0	558	5.7	0.26	5.2	6.3
1986	277	25.0	1.55	22.1	28.2	*	*	*	*	*
1987	271	24.2	1.52	21.4	27.4	*	*	*	*	*
1988	259	22.2	1.42	19.5	25.1	*	*	*	*	*
1989	264	23.0	1.46	20.2	26.0	*	*	*	*	*
1990	288	24.7	1.50	21.9	27.9	*	*	*	*	*
1991	292	24.4	1.47	21.6	27.4	14	5.5	1.51	2.9	9.5
1992	263	21.0	1.33	18.4	23.7	11	4.6	1.56	2.0	8.7
1993	321	25.6	1.47	22.8	28.6	10	5.0	1.92	2.0	10.0
1994	320	25.7	1.47	22.9	28.7	22	8.6	2.26	4.8	14.1
1995	316	24.4	1.41	21.7	27.3	15	5.2	1.41	2.8	8.8
1996	304	22.3	1.31	19.8	25.0	15	4.9	1.34	2.7	8.3
1997	352	25.7	1.40	23.1	28.6	16	4.9	1.32	2.7	8.2
1998	295	21.5	1.28	19.0	24.1	21	6.2	1.45	3.7	9.8
1999	354	26.0	1.41	23.3	28.9	28	8.5	1.81	5.3	12.8
2000	346	24.5	1.35	21.9	27.3	20	5.2	1.28	3.0	8.3
2001	320	22.7	1.30	20.2	25.4	22	5.6	1.35	3.3	8.9
2002	346	24.0	1.32	21.4	26.7	25	6.1	1.36	3.7	9.3
2003	346	23.4	1.29	21.0	26.1	32	7.4	1.40	4.9	10.7
2004	358	23.2	1.25	20.8	25.8	36	8.5	1.56	5.7	12.1
2005	343	22.4	1.24	20.1	25.0	31	6.4	1.26	4.2	9.4
2006	320	20.1	1.15	17.9	22.5	23	4.5	1.00	2.7	6.9
2007	342	21.2	1.17	19.0	23.6	29	6.2	1.26	4.0	9.2
2008	325	20.0	1.13	17.8	22.3	33	6.0	1.12	4.1	8.6
2009	306	18.6	1.09	16.6	20.9	23	4.2	0.91	2.6	6.3
2010	327	19.4	1.10	17.3	21.7	34	5.7	1.05	3.9	8.2
2011	319	18.8	1.08	16.8	21.1	36	5.9	1.04	4.1	8.3
2012	348	20.1	1.10	18.0	22.3	31	4.7	0.89	3.1	6.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Breast										
Males										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	443	0.3	0.02	0.3	0.4	348	0.3	0.02	0.3	0.3
1986	12	0.4	0.10	0.2	0.6	*	*	*	*	*
1987	11	0.3	0.09	0.1	0.5	11	0.3	0.10	0.2	0.6
1988	12	0.3	0.08	0.1	0.5	*	*	*	*	*
1989	23	0.5	0.11	0.3	0.8	19	0.5	0.11	0.3	0.7
1990	12	0.3	0.08	0.1	0.5	10	0.3	0.08	0.1	0.5
1991	14	0.4	0.10	0.2	0.6	11	0.4	0.11	0.2	0.6
1992	*	*	*	*	*	*	*	*	*	*
1993	19	0.4	0.10	0.3	0.6	16	0.4	0.10	0.2	0.7
1994	17	0.3	0.08	0.2	0.5	13	0.3	0.08	0.2	0.5
1995	23	0.5	0.11	0.3	0.8	20	0.5	0.12	0.3	0.8
1996	13	0.3	0.07	0.1	0.4	11	0.3	0.08	0.1	0.5
1997	15	0.3	0.09	0.2	0.6	10	0.3	0.08	0.1	0.5
1998	18	0.4	0.10	0.2	0.7	14	0.4	0.10	0.2	0.6
1999	23	0.5	0.10	0.3	0.7	14	0.3	0.09	0.2	0.6
2000	22	0.5	0.10	0.3	0.7	14	0.3	0.09	0.2	0.5
2001	23	0.5	0.10	0.3	0.7	19	0.4	0.10	0.3	0.7
2002	19	0.4	0.08	0.2	0.6	14	0.3	0.08	0.2	0.5
2003	15	0.3	0.08	0.2	0.5	12	0.3	0.08	0.1	0.5
2004	21	0.4	0.09	0.3	0.7	19	0.4	0.10	0.3	0.7
2005	18	0.4	0.09	0.2	0.6	15	0.4	0.09	0.2	0.6
2006	14	0.3	0.07	0.1	0.4	11	0.3	0.08	0.1	0.4
2007	19	0.4	0.08	0.2	0.6	14	0.3	0.08	0.2	0.5
2008	14	0.2	0.07	0.1	0.4	11	0.2	0.07	0.1	0.4
2009	12	0.2	0.06	0.1	0.4	*	*	*	*	*
2010	15	0.3	0.07	0.2	0.4	12	0.3	0.08	0.1	0.5
2011	14	0.3	0.07	0.1	0.4	10	0.2	0.07	0.1	0.4
2012	17	0.3	0.07	0.2	0.4	16	0.3	0.07	0.2	0.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Breast										
Males										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	90	0.7	0.08	0.5	0.8	*	*	*	*	*
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Breast										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	53,400	29.1	0.13	28.9	29.4	44,415	28.3	0.14	28.0	28.5
1986	2,102	35.6	0.79	34.1	37.2	1,822	35.0	0.84	33.4	36.7
1987	2,090	35.2	0.79	33.7	36.8	1,813	34.5	0.83	32.9	36.2
1988	2,096	35.0	0.78	33.5	36.5	1,831	34.9	0.84	33.2	36.5
1989	2,149	35.7	0.78	34.2	37.3	1,882	35.7	0.84	34.1	37.4
1990	2,182	35.9	0.78	34.4	37.5	1,893	35.6	0.84	33.9	37.2
1991	2,270	36.7	0.78	35.2	38.3	1,967	36.4	0.84	34.8	38.1
1992	2,156	34.4	0.75	32.9	35.9	1,884	34.4	0.81	32.9	36.1
1993	2,162	34.0	0.74	32.5	35.5	1,834	33.1	0.79	31.5	34.7
1994	2,139	33.3	0.73	31.9	34.7	1,801	32.3	0.77	30.8	33.9
1995	2,126	32.7	0.72	31.3	34.1	1,798	31.9	0.77	30.4	33.5
1996	2,108	32.1	0.71	30.8	33.5	1,791	31.5	0.76	30.0	33.0
1997	2,038	31.0	0.69	29.6	32.4	1,675	29.5	0.73	28.1	31.0
1998	1,979	29.5	0.67	28.2	30.8	1,667	28.9	0.72	27.5	30.3
1999	2,007	29.5	0.66	28.2	30.8	1,634	27.9	0.70	26.5	29.3
2000	2,045	29.6	0.66	28.4	31.0	1,687	28.5	0.70	27.1	29.9
2001	1,846	26.6	0.62	25.4	27.9	1,508	25.5	0.67	24.2	26.8
2002	1,897	27.1	0.63	25.9	28.4	1,531	25.7	0.67	24.4	27.1
2003	1,872	26.5	0.62	25.3	27.7	1,497	24.9	0.65	23.6	26.2
2004	1,830	25.6	0.60	24.4	26.8	1,438	23.6	0.63	22.4	24.9
2005	1,865	25.8	0.60	24.6	27.0	1,494	24.4	0.64	23.1	25.7
2006	1,771	24.3	0.58	23.2	25.5	1,431	23.3	0.63	22.1	24.5
2007	1,806	24.6	0.59	23.5	25.8	1,440	23.3	0.62	22.1	24.5
2008	1,780	23.8	0.57	22.7	24.9	1,425	22.6	0.61	21.4	23.9
2009	1,719	22.8	0.56	21.7	23.9	1,394	22.1	0.61	20.9	23.3
2010	1,740	22.6	0.55	21.5	23.7	1,382	21.4	0.59	20.2	22.5
2011	1,815	23.2	0.56	22.2	24.4	1,464	22.4	0.60	21.2	23.6
2012	1,810	22.9	0.55	21.8	24.0	1,432	21.8	0.59	20.7	23.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Breast										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	8,432	37.7	0.41	36.9	38.6	553	10.3	0.46	9.4	11.3
1986	274	42.3	2.60	37.3	47.7	*	*	*	*	*
1987	271	41.6	2.57	36.7	47.0	*	*	*	*	*
1988	256	37.5	2.39	33.0	42.5	*	*	*	*	*
1989	260	38.3	2.42	33.7	43.3	*	*	*	*	*
1990	286	41.7	2.50	36.9	46.8	*	*	*	*	*
1991	289	40.9	2.44	36.3	46.0	14	10.5	2.87	5.6	18.0
1992	262	35.6	2.23	31.4	40.2	10	7.8	2.71	3.4	15.0
1993	318	42.7	2.43	38.1	47.8	10	8.9	3.27	3.7	17.3
1994	316	42.5	2.42	37.9	47.5	22	15.7	3.88	9.0	25.0
1995	313	40.8	2.33	36.3	45.6	15	9.6	2.57	5.3	16.2
1996	302	37.8	2.20	33.6	42.3	15	9.1	2.42	4.9	15.2
1997	347	43.2	2.34	38.7	48.0	16	9.1	2.39	5.1	15.1
1998	291	35.6	2.11	31.6	40.0	21	11.5	2.62	6.9	17.8
1999	345	42.3	2.30	38.0	47.1	28	15.4	3.14	9.8	22.7
2000	339	40.2	2.20	36.0	44.7	19	8.7	2.10	5.1	13.9
2001	316	37.4	2.12	33.4	41.8	22	10.1	2.31	6.1	15.7
2002	341	39.5	2.15	35.4	43.9	25	11.0	2.35	6.9	16.6
2003	343	39.0	2.12	34.9	43.3	32	13.6	2.51	9.1	19.4
2004	356	39.3	2.10	35.3	43.6	36	15.2	2.70	10.4	21.4
2005	340	37.3	2.04	33.4	41.5	31	11.7	2.21	7.8	16.9
2006	317	33.8	1.92	30.2	37.8	23	8.2	1.79	5.1	12.5
2007	337	35.6	1.96	31.8	39.6	29	11.1	2.19	7.3	16.2
2008	322	33.6	1.89	30.0	37.5	33	11.1	2.02	7.5	15.8
2009	303	31.1	1.81	27.7	34.8	22	7.1	1.57	4.4	10.9
2010	325	32.7	1.84	29.2	36.5	33	10.2	1.86	6.9	14.5
2011	316	31.6	1.80	28.1	35.3	35	10.6	1.87	7.3	14.9
2012	347	33.8	1.84	30.3	37.6	31	8.6	1.60	5.8	12.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Female Genital System										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	32,648	17.7	0.10	17.5	17.9	27,203	17.2	0.11	17.0	17.4
1986	1,225	20.3	0.59	19.2	21.5	1,056	19.8	0.63	18.6	21.1
1987	1,217	19.8	0.58	18.7	21.0	1,057	19.5	0.62	18.4	20.8
1988	1,132	18.4	0.56	17.3	19.5	961	17.7	0.58	16.6	18.9
1989	1,176	19.2	0.57	18.1	20.3	984	18.2	0.59	17.1	19.4
1990	1,224	19.6	0.57	18.5	20.8	1,060	19.3	0.61	18.2	20.6
1991	1,206	19.1	0.56	18.0	20.2	1,005	18.2	0.59	17.1	19.4
1992	1,191	18.6	0.55	17.6	19.7	978	17.5	0.57	16.4	18.6
1993	1,209	18.6	0.54	17.5	19.7	1,011	17.8	0.57	16.7	19.0
1994	1,192	18.3	0.53	17.2	19.3	1,007	17.8	0.57	16.7	18.9
1995	1,168	17.7	0.52	16.7	18.8	997	17.4	0.56	16.4	18.6
1996	1,218	18.3	0.53	17.2	19.3	1,035	18.0	0.57	16.9	19.1
1997	1,204	18.0	0.52	17.0	19.0	1,001	17.4	0.56	16.3	18.5
1998	1,213	17.9	0.52	16.9	18.9	1,020	17.5	0.55	16.4	18.6
1999	1,202	17.7	0.51	16.7	18.7	1,017	17.4	0.55	16.3	18.5
2000	1,264	18.5	0.52	17.5	19.5	1,062	18.2	0.57	17.1	19.4
2001	1,170	17.1	0.50	16.1	18.1	948	16.3	0.53	15.3	17.4
2002	1,218	17.6	0.51	16.6	18.6	1,011	17.2	0.55	16.1	18.3
2003	1,177	16.7	0.49	15.8	17.7	972	16.2	0.53	15.2	17.3
2004	1,186	16.8	0.49	15.8	17.8	963	16.1	0.53	15.1	17.2
2005	1,227	17.1	0.49	16.1	18.1	1,006	16.5	0.53	15.5	17.6
2006	1,226	16.9	0.49	15.9	17.8	1,014	16.5	0.53	15.5	17.5
2007	1,205	16.4	0.48	15.5	17.4	1,008	16.4	0.53	15.4	17.5
2008	1,215	16.4	0.48	15.5	17.4	1,008	16.2	0.52	15.2	17.3
2009	1,178	15.8	0.47	14.9	16.7	945	15.2	0.50	14.2	16.2
2010	1,266	16.7	0.48	15.7	17.6	1,022	16.1	0.52	15.1	17.2
2011	1,264	16.5	0.47	15.5	17.4	1,020	16.0	0.51	15.0	17.1
2012	1,275	16.2	0.46	15.3	17.1	1,035	15.8	0.50	14.9	16.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Female Genital System										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	5,050	22.9	0.33	22.3	23.6	392	8.1	0.43	7.2	9.0
1986	165	25.3	2.01	21.5	29.5	*	*	*	*	*
1987	153	23.6	1.95	19.9	27.7	*	*	*	*	*
1988	166	25.1	1.99	21.3	29.3	*	*	*	*	*
1989	183	27.1	2.04	23.3	31.4	*	*	*	*	*
1990	156	22.4	1.82	19.0	26.3	*	*	*	*	*
1991	191	27.5	2.03	23.7	31.8	*	*	*	*	*
1992	201	28.2	2.02	24.4	32.4	12	9.3	2.81	4.6	16.7
1993	190	25.5	1.88	21.9	29.4	*	*	*	*	*
1994	180	24.5	1.85	21.0	28.4	*	*	*	*	*
1995	164	21.8	1.72	18.5	25.4	*	*	*	*	*
1996	168	22.1	1.72	18.8	25.7	15	12.4	3.52	6.5	21.0
1997	192	24.5	1.79	21.2	28.3	11	8.2	2.58	3.9	14.7
1998	176	22.2	1.69	19.0	25.8	17	11.8	3.23	6.3	19.6
1999	175	21.3	1.62	18.3	24.7	10	4.8	1.59	2.2	9.1
2000	188	22.9	1.68	19.7	26.4	14	7.3	2.08	3.8	12.5
2001	204	24.5	1.73	21.2	28.1	18	9.4	2.33	5.4	15.0
2002	186	21.5	1.59	18.5	24.9	21	10.1	2.33	6.1	15.7
2003	188	21.5	1.58	18.5	24.8	17	7.8	2.03	4.4	12.8
2004	207	23.7	1.66	20.5	27.1	16	6.7	1.74	3.7	11.0
2005	198	22.2	1.59	19.2	25.5	23	9.2	2.05	5.7	14.1
2006	192	21.7	1.58	18.7	25.0	20	9.1	2.14	5.4	14.2
2007	181	19.2	1.44	16.5	22.3	16	5.7	1.50	3.2	9.5
2008	183	19.4	1.45	16.7	22.4	24	9.3	2.03	5.8	14.1
2009	209	21.0	1.47	18.2	24.1	24	9.3	2.01	5.8	14.0
2010	218	22.2	1.53	19.4	25.4	26	8.5	1.74	5.5	12.6
2011	220	22.1	1.51	19.2	25.2	24	7.4	1.56	4.6	11.1
2012	216	21.1	1.46	18.3	24.1	24	7.6	1.63	4.8	11.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Cervix Uteri										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	5,538	3.2	0.04	3.1	3.3	3,848	2.6	0.04	2.6	2.7
1986	236	4.1	0.27	3.6	4.7	170	3.3	0.26	2.8	3.9
1987	217	3.9	0.27	3.3	4.4	162	3.3	0.27	2.8	3.9
1988	225	4.0	0.27	3.4	4.5	158	3.2	0.26	2.7	3.7
1989	223	3.9	0.27	3.4	4.5	150	3.1	0.26	2.6	3.6
1990	213	3.7	0.26	3.2	4.3	159	3.2	0.26	2.7	3.8
1991	243	4.1	0.27	3.6	4.7	153	3.0	0.25	2.6	3.6
1992	218	3.6	0.25	3.1	4.1	137	2.6	0.23	2.2	3.1
1993	234	3.8	0.25	3.3	4.3	159	3.0	0.24	2.6	3.6
1994	205	3.3	0.23	2.9	3.8	142	2.8	0.23	2.3	3.3
1995	201	3.2	0.23	2.8	3.7	128	2.4	0.21	2.0	2.8
1996	200	3.2	0.23	2.8	3.7	138	2.7	0.23	2.2	3.2
1997	231	3.6	0.24	3.2	4.2	167	3.2	0.25	2.7	3.7
1998	219	3.4	0.23	3.0	3.9	156	2.9	0.24	2.5	3.4
1999	218	3.3	0.23	2.9	3.8	165	3.0	0.24	2.5	3.5
2000	214	3.3	0.23	2.9	3.8	156	2.9	0.23	2.4	3.4
2001	183	2.8	0.21	2.4	3.2	122	2.3	0.21	1.9	2.7
2002	206	3.1	0.22	2.7	3.6	144	2.6	0.22	2.2	3.1
2003	185	2.8	0.21	2.4	3.2	122	2.2	0.20	1.8	2.7
2004	196	2.9	0.21	2.5	3.3	139	2.5	0.21	2.1	2.9
2005	183	2.7	0.20	2.3	3.1	124	2.2	0.20	1.8	2.6
2006	173	2.5	0.19	2.1	2.9	120	2.1	0.19	1.7	2.5
2007	189	2.7	0.20	2.3	3.1	143	2.5	0.21	2.1	2.9
2008	200	2.8	0.20	2.4	3.2	146	2.5	0.21	2.1	2.9
2009	178	2.5	0.19	2.1	2.9	120	2.1	0.19	1.7	2.5
2010	186	2.6	0.19	2.2	3.0	127	2.2	0.20	1.8	2.6
2011	195	2.7	0.19	2.3	3.1	133	2.2	0.20	1.9	2.7
2012	167	2.2	0.17	1.9	2.6	108	1.8	0.17	1.4	2.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Cervix Uteri										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,611	7.0	0.18	6.7	7.4	78	1.5	0.17	1.1	1.8
1986	65	9.8	1.25	7.5	12.6	*	*	*	*	*
1987	53	8.0	1.12	6.0	10.5	*	*	*	*	*
1988	65	9.3	1.19	7.1	11.9	*	*	*	*	*
1989	72	10.2	1.23	7.9	12.9	*	*	*	*	*
1990	51	7.2	1.02	5.3	9.4	*	*	*	*	*
1991	85	11.4	1.27	9.1	14.2	*	*	*	*	*
1992	78	10.8	1.25	8.5	13.5	*	*	*	*	*
1993	72	9.6	1.15	7.5	12.1	*	*	*	*	*
1994	63	8.1	1.04	6.2	10.4	*	*	*	*	*
1995	71	9.2	1.11	7.2	11.6	*	*	*	*	*
1996	57	7.1	0.95	5.3	9.2	*	*	*	*	*
1997	60	7.3	0.95	5.5	9.4	*	*	*	*	*
1998	58	6.9	0.92	5.3	9.0	*	*	*	*	*
1999	52	6.1	0.84	4.5	7.9	*	*	*	*	*
2000	52	5.8	0.81	4.3	7.6	*	*	*	*	*
2001	57	6.6	0.88	5.0	8.5	*	*	*	*	*
2002	58	6.4	0.84	4.9	8.3	*	*	*	*	*
2003	60	6.4	0.83	4.9	8.3	*	*	*	*	*
2004	56	6.1	0.82	4.6	7.9	*	*	*	*	*
2005	54	5.7	0.79	4.3	7.5	*	*	*	*	*
2006	50	5.4	0.77	4.0	7.2	*	*	*	*	*
2007	45	4.6	0.70	3.4	6.2	*	*	*	*	*
2008	52	5.2	0.73	3.9	6.9	*	*	*	*	*
2009	54	5.3	0.73	4.0	6.9	*	*	*	*	*
2010	58	5.7	0.76	4.3	7.4	*	*	*	*	*
2011	54	5.3	0.73	4.0	6.9	*	*	*	*	*
2012	59	5.6	0.74	4.2	7.3	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Corpus and Uterus, NOS

Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	8,793	4.7	0.05	4.6	4.8	7,128	4.4	0.05	4.3	4.5
1986	307	4.9	0.28	4.3	5.4	265	4.7	0.29	4.1	5.3
1987	324	5.1	0.28	4.5	5.7	270	4.8	0.29	4.2	5.4
1988	309	4.8	0.28	4.3	5.4	270	4.8	0.29	4.2	5.4
1989	292	4.5	0.27	4.0	5.1	246	4.3	0.27	3.7	4.8
1990	310	4.8	0.27	4.2	5.3	263	4.5	0.28	4.0	5.1
1991	314	4.9	0.28	4.3	5.5	265	4.7	0.29	4.1	5.3
1992	268	4.0	0.25	3.5	4.5	216	3.7	0.25	3.2	4.2
1993	302	4.5	0.26	4.0	5.0	248	4.2	0.27	3.7	4.7
1994	308	4.6	0.26	4.1	5.1	250	4.2	0.27	3.7	4.8
1995	304	4.5	0.26	4.0	5.0	253	4.3	0.27	3.7	4.8
1996	295	4.3	0.25	3.8	4.8	245	4.1	0.26	3.6	4.6
1997	297	4.3	0.25	3.8	4.8	246	4.1	0.27	3.6	4.7
1998	332	4.7	0.26	4.2	5.3	268	4.4	0.27	3.9	5.0
1999	307	4.4	0.25	3.9	5.0	255	4.2	0.27	3.7	4.8
2000	335	4.8	0.26	4.3	5.4	270	4.5	0.28	4.0	5.1
2001	321	4.6	0.26	4.1	5.1	263	4.4	0.27	3.8	4.9
2002	322	4.6	0.26	4.1	5.1	252	4.2	0.27	3.7	4.8
2003	301	4.2	0.24	3.8	4.7	247	4.0	0.26	3.6	4.6
2004	317	4.4	0.25	3.9	4.9	237	3.9	0.25	3.4	4.4
2005	330	4.5	0.25	4.0	5.1	260	4.2	0.26	3.7	4.7
2006	351	4.8	0.26	4.3	5.3	273	4.4	0.27	3.9	4.9
2007	331	4.5	0.25	4.0	5.0	269	4.4	0.27	3.9	5.0
2008	334	4.5	0.25	4.0	5.0	265	4.2	0.26	3.7	4.7
2009	365	4.9	0.26	4.4	5.4	272	4.4	0.27	3.9	4.9
2010	405	5.3	0.27	4.8	5.9	312	4.9	0.28	4.4	5.5
2011	389	5.0	0.26	4.5	5.5	310	4.8	0.28	4.2	5.3
2012	423	5.3	0.26	4.8	5.8	338	5.0	0.28	4.5	5.6

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Corpus and Uterus, NOS

Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,571	7.3	0.19	6.9	7.7	92	2.1	0.23	1.7	2.6
1986	41	6.3	0.99	4.5	8.5	*	*	*	*	*
1987	53	8.1	1.13	6.0	10.6	*	*	*	*	*
1988	38	6.1	1.00	4.3	8.3	*	*	*	*	*
1989	44	6.7	1.03	4.9	9.0	*	*	*	*	*
1990	47	6.9	1.02	5.1	9.2	*	*	*	*	*
1991	47	7.1	1.04	5.2	9.4	*	*	*	*	*
1992	50	7.0	1.01	5.2	9.3	*	*	*	*	*
1993	53	7.2	1.00	5.4	9.4	*	*	*	*	*
1994	58	8.1	1.07	6.1	10.4	*	*	*	*	*
1995	50	7.0	1.00	5.2	9.2	*	*	*	*	*
1996	45	6.0	0.91	4.4	8.1	*	*	*	*	*
1997	48	6.6	0.96	4.9	8.7	*	*	*	*	*
1998	58	7.7	1.02	5.8	9.9	*	*	*	*	*
1999	50	6.2	0.89	4.6	8.2	*	*	*	*	*
2000	62	7.8	1.00	6.0	10.0	*	*	*	*	*
2001	55	6.6	0.89	4.9	8.5	*	*	*	*	*
2002	61	7.2	0.93	5.5	9.3	*	*	*	*	*
2003	51	6.0	0.85	4.5	7.9	*	*	*	*	*
2004	75	8.8	1.02	6.9	11.0	*	*	*	*	*
2005	67	7.5	0.92	5.8	9.5	*	*	*	*	*
2006	73	8.5	1.00	6.7	10.7	*	*	*	*	*
2007	58	6.3	0.83	4.7	8.1	*	*	*	*	*
2008	63	6.7	0.85	5.1	8.5	*	*	*	*	*
2009	86	8.8	0.96	7.0	10.9	*	*	*	*	*
2010	82	8.4	0.94	6.7	10.5	11	3.5	1.07	1.7	6.2
2011	76	7.7	0.90	6.1	9.7	*	*	*	*	*
2012	80	8.1	0.91	6.4	10.0	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Ovary										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	16,535	8.9	0.07	8.8	9.1	14,659	9.3	0.08	9.1	9.4
1986	619	10.3	0.42	9.5	11.2	567	10.8	0.47	9.9	11.8
1987	610	9.9	0.41	9.1	10.7	569	10.5	0.45	9.6	11.4
1988	551	8.9	0.39	8.2	9.7	493	9.0	0.42	8.2	9.9
1989	617	10.1	0.41	9.3	10.9	549	10.2	0.45	9.4	11.1
1990	635	10.2	0.41	9.4	11.0	577	10.6	0.45	9.7	11.5
1991	589	9.2	0.39	8.5	10.0	532	9.5	0.42	8.7	10.4
1992	640	10.0	0.40	9.3	10.9	571	10.3	0.44	9.5	11.2
1993	595	9.2	0.38	8.4	10.0	531	9.4	0.42	8.6	10.3
1994	625	9.6	0.39	8.8	10.3	568	10.0	0.43	9.2	10.9
1995	599	9.2	0.38	8.4	9.9	558	9.9	0.42	9.0	10.7
1996	659	9.9	0.39	9.2	10.7	595	10.4	0.43	9.5	11.2
1997	619	9.2	0.37	8.5	10.0	537	9.3	0.41	8.5	10.1
1998	605	8.9	0.37	8.2	9.7	547	9.4	0.41	8.6	10.2
1999	604	8.9	0.36	8.2	9.6	530	9.1	0.40	8.3	9.9
2000	634	9.2	0.37	8.5	10.0	566	9.7	0.41	8.9	10.5
2001	594	8.7	0.36	8.0	9.4	509	8.8	0.39	8.0	9.6
2002	619	8.9	0.36	8.2	9.7	553	9.4	0.40	8.6	10.2
2003	627	8.8	0.36	8.2	9.6	547	9.1	0.39	8.3	9.9
2004	624	8.8	0.36	8.1	9.5	545	9.1	0.40	8.4	9.9
2005	657	9.1	0.36	8.4	9.8	570	9.4	0.40	8.6	10.2
2006	630	8.6	0.35	8.0	9.3	563	9.2	0.39	8.4	10.0
2007	620	8.4	0.34	7.7	9.1	536	8.6	0.38	7.9	9.4
2008	595	8.1	0.33	7.4	8.7	520	8.4	0.37	7.7	9.2
2009	578	7.7	0.32	7.0	8.3	500	7.9	0.36	7.2	8.7
2010	594	7.8	0.32	7.1	8.4	512	8.0	0.36	7.3	8.7
2011	606	7.9	0.33	7.3	8.5	512	8.1	0.36	7.4	8.8
2012	590	7.6	0.32	6.9	8.2	502	7.8	0.36	7.1	8.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Ovary										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,666	7.7	0.19	7.3	8.1	210	4.3	0.31	3.7	4.9
1986	50	7.6	1.09	5.6	10.0	*	*	*	*	*
1987	37	5.9	0.99	4.1	8.2	*	*	*	*	*
1988	56	8.6	1.17	6.4	11.2	*	*	*	*	*
1989	62	9.5	1.22	7.2	12.1	*	*	*	*	*
1990	54	7.8	1.07	5.8	10.1	*	*	*	*	*
1991	54	8.3	1.15	6.2	10.9	*	*	*	*	*
1992	62	8.7	1.12	6.7	11.2	*	*	*	*	*
1993	60	8.1	1.06	6.1	10.4	*	*	*	*	*
1994	53	7.5	1.04	5.6	9.8	*	*	*	*	*
1995	37	4.8	0.80	3.4	6.6	*	*	*	*	*
1996	59	8.1	1.06	6.1	10.4	*	*	*	*	*
1997	78	9.9	1.13	7.8	12.3	*	*	*	*	*
1998	53	6.7	0.92	5.0	8.7	*	*	*	*	*
1999	67	8.3	1.03	6.5	10.6	*	*	*	*	*
2000	64	8.0	1.01	6.1	10.2	*	*	*	*	*
2001	74	9.1	1.07	7.1	11.4	11	5.3	1.66	2.5	9.6
2002	58	6.8	0.90	5.2	8.8	*	*	*	*	*
2003	70	8.2	0.98	6.3	10.3	10	4.6	1.59	2.1	8.8
2004	70	8.1	0.97	6.3	10.2	*	*	*	*	*
2005	72	8.4	0.99	6.5	10.5	15	5.8	1.58	3.2	9.8
2006	55	6.2	0.84	4.7	8.1	12	5.5	1.68	2.7	9.7
2007	74	7.9	0.93	6.2	9.9	10	3.5	1.15	1.6	6.5
2008	60	6.6	0.86	5.0	8.5	15	5.8	1.60	3.1	9.8
2009	66	6.7	0.83	5.1	8.5	12	4.2	1.28	2.1	7.4
2010	70	7.2	0.87	5.6	9.1	12	3.8	1.14	1.9	6.7
2011	81	8.1	0.91	6.4	10.1	13	4.1	1.19	2.1	7.1
2012	70	6.7	0.82	5.2	8.5	18	5.3	1.30	3.0	8.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Male Genital System

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	37,093	31.5	0.17	31.2	31.8	29,322	28.3	0.17	28.0	28.7
1986	1,226	33.5	1.01	31.6	35.6	992	30.5	1.02	28.6	32.6
1987	1,310	35.4	1.02	33.4	37.4	1,057	31.8	1.02	29.8	33.9
1988	1,379	36.5	1.03	34.6	38.6	1,115	33.2	1.04	31.2	35.3
1989	1,423	37.7	1.04	35.6	39.7	1,131	33.6	1.04	31.6	35.7
1990	1,562	41.4	1.09	39.3	43.6	1,267	37.6	1.10	35.5	39.9
1991	1,555	40.4	1.07	38.4	42.6	1,250	36.5	1.07	34.4	38.6
1992	1,523	38.9	1.03	36.9	41.0	1,208	34.4	1.02	32.4	36.5
1993	1,637	41.6	1.07	39.5	43.7	1,308	37.4	1.07	35.3	39.5
1994	1,533	38.5	1.02	36.5	40.5	1,221	34.4	1.01	32.4	36.4
1995	1,513	37.5	0.99	35.6	39.5	1,239	34.6	1.01	32.7	36.7
1996	1,518	36.9	0.97	35.0	38.9	1,212	33.2	0.98	31.3	35.2
1997	1,491	35.8	0.95	34.0	37.7	1,190	32.3	0.96	30.5	34.3
1998	1,515	35.8	0.94	34.0	37.7	1,211	32.4	0.95	30.5	34.3
1999	1,447	33.7	0.90	31.9	35.5	1,160	30.6	0.91	28.8	32.4
2000	1,436	33.3	0.89	31.6	35.1	1,128	29.6	0.89	27.9	31.4
2001	1,427	32.5	0.87	30.8	34.2	1,139	29.4	0.88	27.7	31.1
2002	1,353	30.1	0.83	28.5	31.8	1,045	26.5	0.83	24.9	28.2
2003	1,249	27.7	0.79	26.1	29.3	977	24.7	0.80	23.1	26.3
2004	1,256	27.3	0.78	25.8	28.9	966	23.9	0.78	22.4	25.5
2005	1,290	27.4	0.77	25.9	29.0	993	24.2	0.77	22.7	25.8
2006	1,244	26.2	0.75	24.7	27.7	952	22.8	0.74	21.4	24.3
2007	1,285	26.3	0.74	24.9	27.8	993	23.3	0.74	21.8	24.8
2008	1,205	24.2	0.71	22.9	25.6	948	21.9	0.72	20.5	23.3
2009	1,211	23.9	0.69	22.5	25.3	913	20.8	0.69	19.4	22.2
2010	1,200	23.1	0.67	21.8	24.4	905	20.0	0.67	18.7	21.4
2011	1,134	21.4	0.64	20.1	22.6	892	19.4	0.66	18.2	20.8
2012	1,171	21.6	0.64	20.3	22.8	910	19.3	0.65	18.1	20.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Male Genital System

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	7,556	66.8	0.81	65.2	68.4	215	9.3	0.69	8.0	10.7
1986	232	66.3	4.66	57.5	75.9	*	*	*	*	*
1987	251	74.0	5.02	64.5	84.4	*	*	*	*	*
1988	262	73.2	4.82	64.1	83.2	*	*	*	*	*
1989	290	81.9	5.10	72.2	92.3	*	*	*	*	*
1990	294	83.7	5.24	73.8	94.5	*	*	*	*	*
1991	300	84.4	5.18	74.6	95.1	*	*	*	*	*
1992	307	88.3	5.37	78.1	99.3	*	*	*	*	*
1993	324	88.0	5.22	78.1	98.7	*	*	*	*	*
1994	309	84.6	5.11	74.9	95.1	*	*	*	*	*
1995	270	70.3	4.54	61.7	79.6	*	*	*	*	*
1996	298	78.5	4.81	69.4	88.4	*	*	*	*	*
1997	292	73.1	4.54	64.5	82.5	*	*	*	*	*
1998	295	71.9	4.42	63.5	81.0	*	*	*	*	*
1999	281	67.5	4.24	59.5	76.2	*	*	*	*	*
2000	297	72.8	4.45	64.3	81.9	11	11.7	3.90	5.4	21.3
2001	283	67.2	4.22	59.2	75.9	*	*	*	*	*
2002	298	68.2	4.15	60.3	76.7	10	7.3	2.51	3.2	13.7
2003	261	58.8	3.84	51.5	66.8	11	11.3	3.47	5.5	19.7
2004	280	64.3	4.05	56.6	72.6	10	10.2	3.43	4.6	18.5
2005	286	62.3	3.92	54.8	70.3	11	9.9	3.31	4.6	17.9
2006	286	63.6	3.96	56.1	71.8	*	*	*	*	*
2007	281	59.8	3.75	52.6	67.5	11	8.2	2.70	3.8	14.7
2008	246	50.3	3.36	44.0	57.3	11	7.1	2.44	3.2	13.1
2009	276	54.1	3.41	47.6	61.1	22	14.9	3.53	8.8	23.0
2010	281	54.7	3.40	48.2	61.7	14	10.8	3.00	5.8	17.8
2011	229	41.9	2.88	36.4	47.8	13	7.4	2.18	3.7	12.6
2012	247	45.2	2.98	39.5	51.3	14	7.5	2.18	3.9	12.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Prostate										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	36,479	31.1	0.17	30.8	31.4	28,787	27.9	0.17	27.6	28.2
1986	1,203	33.1	1.00	31.1	35.1	973	30.1	1.01	28.1	32.1
1987	1,274	34.6	1.02	32.7	36.7	1,023	31.0	1.01	29.0	33.0
1988	1,362	36.2	1.02	34.2	38.3	1,099	32.8	1.03	30.8	34.9
1989	1,395	37.0	1.04	35.0	39.1	1,107	33.1	1.04	31.1	35.1
1990	1,545	41.0	1.09	38.9	43.2	1,252	37.3	1.10	35.2	39.5
1991	1,533	40.0	1.06	37.9	42.1	1,230	36.0	1.07	33.9	38.2
1992	1,502	38.5	1.03	36.5	40.5	1,192	34.1	1.02	32.1	36.1
1993	1,609	41.1	1.06	39.0	43.2	1,284	36.9	1.06	34.8	39.0
1994	1,517	38.2	1.01	36.3	40.3	1,206	34.1	1.01	32.1	36.1
1995	1,499	37.2	0.99	35.3	39.2	1,226	34.4	1.01	32.4	36.4
1996	1,502	36.6	0.97	34.7	38.6	1,196	32.9	0.97	31.0	34.8
1997	1,465	35.3	0.95	33.5	37.2	1,167	31.8	0.95	30.0	33.7
1998	1,492	35.4	0.93	33.6	37.2	1,191	31.9	0.94	30.1	33.8
1999	1,423	33.2	0.90	31.5	35.0	1,140	30.2	0.91	28.4	32.0
2000	1,409	32.8	0.89	31.1	34.6	1,105	29.1	0.89	27.4	30.9
2001	1,402	32.0	0.87	30.3	33.7	1,116	28.8	0.87	27.1	30.6
2002	1,331	29.7	0.83	28.1	31.4	1,026	26.1	0.82	24.5	27.8
2003	1,234	27.4	0.79	25.9	29.0	966	24.4	0.79	22.9	26.1
2004	1,228	26.8	0.77	25.3	28.4	940	23.4	0.77	21.9	24.9
2005	1,270	27.1	0.77	25.6	28.6	975	23.8	0.77	22.3	25.4
2006	1,220	25.7	0.74	24.3	27.2	933	22.4	0.74	21.0	23.9
2007	1,261	25.9	0.74	24.5	27.4	974	22.9	0.74	21.4	24.4
2008	1,174	23.7	0.70	22.3	25.1	923	21.4	0.71	20.0	22.8
2009	1,189	23.5	0.69	22.2	24.9	897	20.4	0.69	19.1	21.8
2010	1,171	22.6	0.67	21.3	23.9	878	19.5	0.66	18.2	20.9
2011	1,117	21.1	0.64	19.9	22.4	876	19.1	0.65	17.9	20.4
2012	1,152	21.3	0.63	20.0	22.5	892	19.0	0.64	17.8	20.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Prostate										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	7,482	66.3	0.81	64.8	67.9	210	9.2	0.69	7.9	10.6
1986	228	65.7	4.65	56.9	75.3	*	*	*	*	*
1987	249	73.8	5.02	64.3	84.1	*	*	*	*	*
1988	261	73.2	4.82	64.0	83.1	*	*	*	*	*
1989	286	80.7	5.06	71.1	91.1	*	*	*	*	*
1990	292	83.1	5.21	73.2	93.8	*	*	*	*	*
1991	298	84.1	5.17	74.2	94.7	*	*	*	*	*
1992	302	87.1	5.34	77.0	98.0	*	*	*	*	*
1993	320	87.2	5.21	77.3	97.9	*	*	*	*	*
1994	308	84.5	5.11	74.8	95.0	*	*	*	*	*
1995	269	70.2	4.54	61.6	79.5	*	*	*	*	*
1996	298	78.5	4.81	69.4	88.4	*	*	*	*	*
1997	289	72.7	4.53	64.1	82.0	*	*	*	*	*
1998	292	71.6	4.42	63.2	80.6	*	*	*	*	*
1999	277	67.0	4.23	59.0	75.7	*	*	*	*	*
2000	293	72.3	4.44	63.8	81.4	11	11.7	3.90	5.4	21.3
2001	281	67.0	4.22	58.9	75.6	*	*	*	*	*
2002	295	67.6	4.14	59.8	76.1	10	7.3	2.51	3.2	13.7
2003	257	58.2	3.83	51.0	66.1	11	11.3	3.47	5.5	19.7
2004	278	64.1	4.05	56.4	72.4	10	10.2	3.43	4.6	18.5
2005	285	62.1	3.92	54.7	70.2	10	9.5	3.28	4.2	17.5
2006	281	63.0	3.95	55.5	71.2	*	*	*	*	*
2007	276	59.0	3.74	51.9	66.7	11	8.2	2.70	3.8	14.7
2008	242	49.8	3.35	43.4	56.7	*	*	*	*	*
2009	271	53.5	3.40	47.0	60.5	21	14.6	3.52	8.5	22.7
2010	279	54.5	3.39	48.0	61.4	14	10.8	3.00	5.8	17.8
2011	228	41.8	2.88	36.3	47.7	13	7.4	2.18	3.7	12.6
2012	247	45.2	2.98	39.5	51.3	13	7.3	2.17	3.7	12.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Testis										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	361	0.2	0.01	0.2	0.3	323	0.3	0.01	0.2	0.3
1986	10	0.2	0.06	0.1	0.3	*	*	*	*	*
1987	22	0.4	0.08	0.2	0.6	21	0.4	0.10	0.3	0.7
1988	11	0.2	0.05	0.1	0.3	10	0.2	0.06	0.1	0.4
1989	15	0.3	0.09	0.2	0.5	14	0.4	0.10	0.2	0.6
1990	11	0.2	0.06	0.1	0.4	10	0.2	0.07	0.1	0.4
1991	16	0.3	0.08	0.2	0.5	16	0.4	0.09	0.2	0.6
1992	11	0.2	0.06	0.1	0.4	10	0.2	0.07	0.1	0.4
1993	20	0.3	0.08	0.2	0.5	18	0.4	0.09	0.2	0.6
1994	11	0.2	0.05	0.1	0.3	10	0.2	0.06	0.1	0.4
1995	*	*	*	*	*	*	*	*	*	*
1996	10	0.2	0.06	0.1	0.3	10	0.2	0.07	0.1	0.4
1997	12	0.2	0.06	0.1	0.4	10	0.2	0.07	0.1	0.4
1998	15	0.3	0.07	0.1	0.4	13	0.3	0.08	0.1	0.5
1999	12	0.2	0.06	0.1	0.3	10	0.2	0.06	0.1	0.4
2000	16	0.3	0.07	0.2	0.4	13	0.3	0.07	0.1	0.4
2001	12	0.2	0.06	0.1	0.3	11	0.2	0.07	0.1	0.4
2002	14	0.2	0.07	0.1	0.4	14	0.3	0.08	0.2	0.5
2003	*	*	*	*	*	*	*	*	*	*
2004	17	0.3	0.07	0.2	0.5	16	0.3	0.08	0.2	0.5
2005	15	0.3	0.07	0.1	0.4	14	0.3	0.08	0.2	0.5
2006	14	0.2	0.06	0.1	0.4	11	0.2	0.07	0.1	0.4
2007	13	0.2	0.06	0.1	0.4	11	0.2	0.07	0.1	0.4
2008	22	0.4	0.08	0.2	0.6	19	0.4	0.09	0.2	0.6
2009	13	0.2	0.06	0.1	0.4	11	0.2	0.07	0.1	0.4
2010	17	0.3	0.06	0.1	0.4	16	0.3	0.07	0.2	0.5
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Testis										
Males										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	35	0.2	0.03	0.1	0.2	*	*	*	*	*
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary System										
Both Sexes										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	29,328	9.1	0.05	9.0	9.2	26,002	9.2	0.06	9.1	9.4
1986	975	9.5	0.31	8.9	10.1	870	9.5	0.33	8.9	10.1
1987	967	9.3	0.30	8.7	9.9	873	9.4	0.32	8.8	10.0
1988	943	8.9	0.29	8.4	9.5	827	8.8	0.31	8.2	9.5
1989	964	9.0	0.29	8.5	9.6	859	9.1	0.31	8.5	9.7
1990	1,010	9.4	0.30	8.8	10.0	895	9.4	0.32	8.7	10.0
1991	985	9.1	0.29	8.5	9.7	877	9.1	0.31	8.5	9.7
1992	1,028	9.3	0.29	8.8	9.9	924	9.4	0.31	8.8	10.1
1993	1,066	9.5	0.29	8.9	10.1	961	9.7	0.31	9.1	10.4
1994	1,124	9.9	0.30	9.4	10.5	993	9.9	0.32	9.3	10.6
1995	1,047	9.1	0.28	8.6	9.7	942	9.4	0.31	8.8	10.0
1996	1,018	8.8	0.28	8.3	9.4	891	8.8	0.29	8.2	9.4
1997	1,035	8.9	0.28	8.4	9.5	898	8.8	0.29	8.2	9.4
1998	1,094	9.3	0.28	8.7	9.9	973	9.4	0.30	8.9	10.0
1999	1,068	9.0	0.27	8.4	9.5	943	9.1	0.30	8.5	9.7
2000	1,140	9.5	0.28	8.9	10.0	1,006	9.6	0.30	9.0	10.2
2001	1,110	9.2	0.27	8.6	9.7	980	9.3	0.30	8.7	9.9
2002	1,071	8.8	0.27	8.3	9.3	952	9.0	0.29	8.5	9.6
2003	1,153	9.3	0.28	8.8	9.9	1,014	9.5	0.30	8.9	10.1
2004	1,135	9.1	0.27	8.6	9.6	1,008	9.4	0.30	8.8	10.0
2005	1,143	9.0	0.27	8.5	9.5	1,023	9.4	0.29	8.8	9.9
2006	1,134	8.9	0.26	8.4	9.4	998	9.1	0.29	8.5	9.7
2007	1,150	8.9	0.26	8.4	9.4	997	9.0	0.29	8.4	9.6
2008	1,226	9.3	0.27	8.8	9.9	1,101	9.8	0.30	9.2	10.4
2009	1,179	8.8	0.26	8.3	9.4	1,034	9.1	0.29	8.5	9.6
2010	1,169	8.6	0.26	8.1	9.1	1,011	8.8	0.28	8.2	9.3
2011	1,176	8.6	0.25	8.1	9.1	1,060	9.1	0.28	8.5	9.6
2012	1,218	8.7	0.25	8.2	9.2	1,092	9.2	0.28	8.7	9.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary System

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,112	8.9	0.16	8.6	9.2	214	3.1	0.23	2.6	3.5
1986	102	9.7	1.01	7.8	11.9	*	*	*	*	*
1987	91	8.5	0.93	6.8	10.5	*	*	*	*	*
1988	114	10.6	1.02	8.7	12.8	*	*	*	*	*
1989	102	9.3	0.95	7.5	11.4	*	*	*	*	*
1990	112	10.6	1.04	8.7	12.8	*	*	*	*	*
1991	105	9.7	0.97	7.9	11.8	*	*	*	*	*
1992	103	9.1	0.92	7.3	11.0	*	*	*	*	*
1993	101	8.5	0.88	6.9	10.4	*	*	*	*	*
1994	122	10.1	0.94	8.4	12.1	*	*	*	*	*
1995	103	8.7	0.88	7.1	10.6	*	*	*	*	*
1996	120	9.8	0.92	8.1	11.8	*	*	*	*	*
1997	130	10.2	0.92	8.5	12.2	*	*	*	*	*
1998	111	8.4	0.82	6.9	10.2	10	3.4	1.15	1.5	6.4
1999	119	9.1	0.85	7.5	10.9	*	*	*	*	*
2000	127	9.8	0.89	8.2	11.7	*	*	*	*	*
2001	123	9.2	0.85	7.6	11.0	*	*	*	*	*
2002	110	8.5	0.83	6.9	10.2	*	*	*	*	*
2003	131	9.5	0.85	7.9	11.2	*	*	*	*	*
2004	118	8.1	0.76	6.7	9.8	*	*	*	*	*
2005	109	7.6	0.75	6.2	9.2	11	4.2	1.30	2.1	7.3
2006	124	8.4	0.77	6.9	10.0	12	3.1	0.96	1.5	5.5
2007	138	9.4	0.82	7.9	11.1	15	3.5	0.97	1.9	5.9
2008	114	7.6	0.73	6.2	9.1	11	3.0	0.97	1.4	5.4
2009	131	8.7	0.78	7.3	10.4	14	2.9	0.83	1.5	4.9
2010	143	9.1	0.78	7.6	10.7	15	3.3	0.92	1.8	5.6
2011	103	6.6	0.67	5.4	8.1	13	2.9	0.84	1.5	5.0
2012	106	6.4	0.64	5.2	7.8	20	4.0	0.96	2.4	6.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary System

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	18,692	14.6	0.11	14.4	14.8	16,831	15.1	0.12	14.8	15.3
1986	625	15.8	0.66	14.5	17.1	569	16.1	0.71	14.8	17.6
1987	603	15.0	0.64	13.7	16.3	548	15.3	0.68	14.0	16.7
1988	605	14.8	0.63	13.6	16.1	524	14.6	0.66	13.3	15.9
1989	594	14.4	0.61	13.2	15.6	541	14.7	0.66	13.5	16.1
1990	623	15.1	0.63	13.9	16.4	566	15.4	0.67	14.1	16.8
1991	627	15.0	0.62	13.8	16.3	562	15.2	0.67	14.0	16.6
1992	638	15.0	0.62	13.8	16.3	586	15.6	0.67	14.4	17.0
1993	682	15.7	0.62	14.5	17.0	622	16.3	0.67	15.0	17.7
1994	715	16.2	0.63	15.0	17.5	646	16.7	0.67	15.4	18.1
1995	663	14.7	0.59	13.6	15.9	594	15.0	0.63	13.8	16.3
1996	639	14.3	0.58	13.2	15.5	574	14.6	0.62	13.4	15.9
1997	646	14.1	0.57	13.0	15.3	572	14.3	0.61	13.2	15.6
1998	686	14.9	0.58	13.8	16.1	619	15.4	0.63	14.2	16.7
1999	661	14.2	0.56	13.1	15.3	585	14.4	0.60	13.2	15.6
2000	713	15.1	0.58	14.0	16.3	642	15.6	0.62	14.4	16.9
2001	691	14.3	0.55	13.2	15.4	627	14.9	0.60	13.8	16.1
2002	681	13.9	0.54	12.9	15.0	620	14.6	0.59	13.4	15.8
2003	749	15.1	0.56	14.0	16.3	672	15.6	0.61	14.5	16.9
2004	721	14.2	0.54	13.2	15.3	655	15.0	0.59	13.8	16.2
2005	754	14.8	0.55	13.8	15.9	680	15.4	0.60	14.2	16.6
2006	735	14.3	0.54	13.2	15.4	658	14.8	0.58	13.7	16.0
2007	736	13.8	0.52	12.8	14.9	649	14.1	0.56	13.0	15.3
2008	784	14.9	0.54	13.8	16.0	715	15.7	0.59	14.6	16.9
2009	779	14.3	0.52	13.3	15.3	694	14.7	0.57	13.6	15.9
2010	745	13.3	0.50	12.4	14.4	656	13.7	0.54	12.6	14.8
2011	791	14.0	0.51	13.0	15.0	723	14.9	0.56	13.8	16.0
2012	806	13.9	0.50	13.0	15.0	732	14.8	0.56	13.7	15.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary System

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,733	12.5	0.32	11.9	13.2	128	4.3	0.43	3.5	5.2
1986	54	12.8	1.90	9.4	17.0	*	*	*	*	*
1987	53	12.5	1.86	9.2	16.6	*	*	*	*	*
1988	79	18.1	2.18	14.1	22.8	*	*	*	*	*
1989	50	12.0	1.83	8.7	16.1	*	*	*	*	*
1990	55	13.8	2.01	10.1	18.2	*	*	*	*	*
1991	65	14.7	1.95	11.1	18.9	*	*	*	*	*
1992	51	10.5	1.56	7.6	13.9	*	*	*	*	*
1993	57	11.8	1.67	8.8	15.5	*	*	*	*	*
1994	64	13.1	1.74	9.9	16.9	*	*	*	*	*
1995	67	14.2	1.85	10.8	18.2	*	*	*	*	*
1996	61	12.3	1.68	9.3	16.0	*	*	*	*	*
1997	70	13.1	1.67	10.0	16.7	*	*	*	*	*
1998	63	11.5	1.55	8.7	14.9	*	*	*	*	*
1999	73	14.0	1.75	10.8	17.8	*	*	*	*	*
2000	67	13.6	1.77	10.4	17.5	*	*	*	*	*
2001	61	10.8	1.49	8.1	14.1	*	*	*	*	*
2002	53	10.6	1.57	7.7	14.0	*	*	*	*	*
2003	72	12.9	1.65	9.9	16.5	*	*	*	*	*
2004	61	10.5	1.44	7.9	13.7	*	*	*	*	*
2005	70	13.6	1.76	10.3	17.4	*	*	*	*	*
2006	74	13.0	1.66	9.9	16.6	*	*	*	*	*
2007	75	13.8	1.72	10.7	17.5	12	6.5	2.03	3.1	11.5
2008	63	10.5	1.40	7.9	13.5	*	*	*	*	*
2009	73	12.6	1.58	9.7	16.0	12	6.0	1.93	2.9	10.9
2010	79	13.3	1.58	10.4	16.7	10	5.1	1.80	2.2	9.6
2011	61	10.2	1.36	7.7	13.2	*	*	*	*	*
2012	62	9.8	1.32	7.4	12.7	12	6.7	2.09	3.3	11.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary System

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	10,636	5.5	0.05	5.4	5.6	9,171	5.4	0.06	5.3	5.6
1986	350	5.6	0.30	5.0	6.2	301	5.4	0.31	4.8	6.0
1987	364	5.8	0.31	5.2	6.4	325	5.8	0.32	5.1	6.4
1988	338	5.3	0.29	4.7	5.9	303	5.3	0.31	4.7	5.9
1989	370	5.8	0.31	5.3	6.5	318	5.6	0.32	5.0	6.3
1990	387	6.0	0.31	5.4	6.6	329	5.7	0.32	5.1	6.4
1991	358	5.4	0.29	4.9	6.0	315	5.4	0.31	4.8	6.0
1992	390	5.8	0.30	5.3	6.4	338	5.7	0.31	5.1	6.3
1993	384	5.7	0.29	5.1	6.3	339	5.6	0.31	5.1	6.3
1994	409	6.0	0.30	5.4	6.6	347	5.7	0.31	5.1	6.4
1995	384	5.6	0.29	5.1	6.2	348	5.8	0.32	5.2	6.5
1996	379	5.4	0.28	4.8	6.0	317	5.1	0.29	4.6	5.7
1997	389	5.5	0.28	5.0	6.1	326	5.3	0.30	4.7	5.9
1998	408	5.7	0.29	5.2	6.3	354	5.7	0.31	5.1	6.3
1999	407	5.6	0.28	5.1	6.2	358	5.7	0.31	5.1	6.3
2000	427	5.9	0.29	5.3	6.5	364	5.8	0.31	5.2	6.4
2001	419	5.7	0.28	5.2	6.3	353	5.6	0.30	5.0	6.2
2002	390	5.3	0.27	4.8	5.9	332	5.3	0.29	4.7	5.9
2003	404	5.5	0.27	4.9	6.0	342	5.3	0.29	4.8	6.0
2004	414	5.6	0.28	5.1	6.2	353	5.5	0.30	4.9	6.1
2005	389	5.2	0.27	4.7	5.7	343	5.3	0.29	4.8	5.9
2006	399	5.3	0.27	4.8	5.9	340	5.2	0.29	4.7	5.8
2007	414	5.4	0.27	4.9	6.0	348	5.3	0.29	4.8	5.9
2008	442	5.7	0.27	5.1	6.2	386	5.8	0.30	5.2	6.5
2009	400	5.1	0.26	4.6	5.6	340	5.1	0.28	4.5	5.6
2010	424	5.3	0.26	4.8	5.9	355	5.2	0.29	4.7	5.8
2011	385	4.8	0.25	4.3	5.3	337	5.0	0.28	4.4	5.5
2012	412	5.1	0.25	4.6	5.6	360	5.2	0.28	4.7	5.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary System

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,379	6.6	0.18	6.2	6.9	86	2.2	0.25	1.8	2.8
1986	48	7.7	1.14	5.6	10.2	*	*	*	*	*
1987	38	6.0	1.00	4.2	8.2	*	*	*	*	*
1988	35	5.6	0.97	3.9	7.8	*	*	*	*	*
1989	52	7.8	1.10	5.8	10.3	*	*	*	*	*
1990	57	8.9	1.19	6.7	11.5	*	*	*	*	*
1991	40	6.3	1.01	4.5	8.6	*	*	*	*	*
1992	52	7.8	1.10	5.8	10.3	*	*	*	*	*
1993	44	6.3	0.96	4.5	8.4	*	*	*	*	*
1994	58	8.1	1.08	6.1	10.5	*	*	*	*	*
1995	36	5.2	0.87	3.6	7.2	*	*	*	*	*
1996	59	7.9	1.04	6.0	10.2	*	*	*	*	*
1997	60	8.1	1.05	6.1	10.4	*	*	*	*	*
1998	48	6.3	0.91	4.6	8.3	*	*	*	*	*
1999	46	6.0	0.88	4.4	7.9	*	*	*	*	*
2000	60	7.6	0.99	5.8	9.8	*	*	*	*	*
2001	62	7.8	1.00	6.0	10.0	*	*	*	*	*
2002	57	7.2	0.96	5.4	9.3	*	*	*	*	*
2003	59	7.1	0.94	5.4	9.2	*	*	*	*	*
2004	57	6.7	0.89	5.0	8.6	*	*	*	*	*
2005	39	4.4	0.71	3.1	6.1	*	*	*	*	*
2006	50	5.7	0.82	4.3	7.6	*	*	*	*	*
2007	63	7.0	0.89	5.4	9.0	*	*	*	*	*
2008	51	5.6	0.79	4.2	7.4	*	*	*	*	*
2009	58	6.4	0.84	4.8	8.2	*	*	*	*	*
2010	64	6.7	0.84	5.1	8.5	*	*	*	*	*
2011	42	4.4	0.68	3.1	5.9	*	*	*	*	*
2012	44	4.5	0.69	3.3	6.1	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary Bladder										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	14,356	4.5	0.04	4.4	4.5	12,894	4.6	0.04	4.5	4.6
1986	485	4.8	0.22	4.4	5.3	439	4.8	0.23	4.4	5.3
1987	460	4.5	0.21	4.1	4.9	412	4.5	0.22	4.1	4.9
1988	469	4.5	0.21	4.1	4.9	416	4.4	0.22	4.0	4.9
1989	447	4.3	0.20	3.9	4.7	396	4.2	0.21	3.8	4.7
1990	498	4.6	0.21	4.2	5.1	440	4.6	0.22	4.2	5.0
1991	464	4.3	0.20	3.9	4.7	413	4.3	0.21	3.9	4.7
1992	516	4.7	0.21	4.3	5.1	462	4.7	0.22	4.3	5.2
1993	503	4.5	0.20	4.1	4.9	462	4.6	0.22	4.2	5.1
1994	525	4.6	0.20	4.3	5.1	471	4.7	0.22	4.3	5.1
1995	501	4.4	0.20	4.0	4.8	462	4.5	0.21	4.1	5.0
1996	512	4.4	0.20	4.1	4.8	452	4.4	0.21	4.0	4.8
1997	505	4.3	0.19	4.0	4.7	453	4.4	0.21	4.0	4.8
1998	519	4.4	0.19	4.0	4.8	468	4.5	0.21	4.1	4.9
1999	560	4.7	0.20	4.3	5.1	501	4.8	0.21	4.4	5.2
2000	541	4.5	0.19	4.1	4.9	486	4.6	0.21	4.2	5.0
2001	520	4.3	0.19	3.9	4.7	467	4.4	0.20	4.0	4.8
2002	511	4.2	0.19	3.8	4.6	463	4.4	0.20	4.0	4.8
2003	557	4.5	0.19	4.2	4.9	510	4.8	0.21	4.4	5.2
2004	574	4.6	0.19	4.2	5.0	521	4.8	0.21	4.4	5.2
2005	560	4.4	0.19	4.1	4.8	508	4.6	0.21	4.2	5.0
2006	579	4.5	0.19	4.2	4.9	522	4.7	0.21	4.3	5.1
2007	574	4.5	0.19	4.1	4.8	504	4.5	0.20	4.1	5.0
2008	615	4.7	0.19	4.3	5.1	562	5.0	0.21	4.5	5.4
2009	597	4.5	0.19	4.1	4.9	531	4.6	0.20	4.3	5.1
2010	589	4.4	0.18	4.0	4.7	518	4.5	0.20	4.1	4.9
2011	579	4.2	0.18	3.9	4.6	515	4.4	0.20	4.0	4.8
2012	596	4.3	0.18	3.9	4.6	540	4.5	0.20	4.1	4.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary Bladder										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,375	4.2	0.12	4.0	4.4	87	1.4	0.16	1.1	1.8
1986	45	4.6	0.71	3.3	6.2	*	*	*	*	*
1987	47	4.8	0.72	3.5	6.4	*	*	*	*	*
1988	52	5.0	0.72	3.7	6.6	*	*	*	*	*
1989	49	5.0	0.73	3.6	6.6	*	*	*	*	*
1990	56	5.7	0.78	4.2	7.4	*	*	*	*	*
1991	50	5.0	0.73	3.7	6.6	*	*	*	*	*
1992	53	5.0	0.70	3.7	6.5	*	*	*	*	*
1993	39	3.6	0.59	2.5	4.9	*	*	*	*	*
1994	51	4.6	0.65	3.4	6.0	*	*	*	*	*
1995	38	3.4	0.57	2.4	4.7	*	*	*	*	*
1996	57	5.0	0.68	3.8	6.5	*	*	*	*	*
1997	49	4.1	0.59	3.0	5.4	*	*	*	*	*
1998	51	4.1	0.58	3.0	5.4	*	*	*	*	*
1999	56	4.6	0.62	3.4	5.9	*	*	*	*	*
2000	53	4.5	0.63	3.3	5.9	*	*	*	*	*
2001	51	4.1	0.59	3.0	5.4	*	*	*	*	*
2002	47	3.8	0.57	2.8	5.1	*	*	*	*	*
2003	44	3.5	0.55	2.6	4.8	*	*	*	*	*
2004	48	3.4	0.50	2.5	4.5	*	*	*	*	*
2005	46	3.5	0.53	2.5	4.7	*	*	*	*	*
2006	55	4.1	0.56	3.1	5.3	*	*	*	*	*
2007	64	4.6	0.58	3.5	5.8	*	*	*	*	*
2008	49	3.5	0.51	2.6	4.6	*	*	*	*	*
2009	64	4.5	0.57	3.5	5.8	*	*	*	*	*
2010	63	4.3	0.55	3.3	5.5	*	*	*	*	*
2011	56	3.7	0.51	2.8	4.9	*	*	*	*	*
2012	42	2.7	0.42	1.9	3.7	14	3.0	0.84	1.6	5.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary Bladder

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	9,625	7.9	0.08	7.8	8.1	8,884	8.3	0.09	8.1	8.5
1986	322	8.8	0.51	7.8	9.9	300	9.2	0.55	8.1	10.3
1987	307	8.1	0.48	7.2	9.1	281	8.3	0.52	7.3	9.4
1988	310	8.0	0.48	7.1	9.0	277	8.1	0.50	7.1	9.1
1989	288	7.4	0.45	6.5	8.3	262	7.5	0.48	6.6	8.5
1990	322	8.3	0.49	7.4	9.3	294	8.5	0.52	7.5	9.6
1991	318	8.2	0.48	7.3	9.2	286	8.2	0.51	7.3	9.3
1992	339	8.5	0.48	7.6	9.5	319	9.0	0.52	8.0	10.0
1993	354	8.7	0.48	7.8	9.7	330	9.1	0.52	8.1	10.2
1994	355	8.5	0.47	7.6	9.5	331	9.0	0.51	8.0	10.0
1995	350	8.1	0.45	7.3	9.1	324	8.5	0.48	7.6	9.5
1996	318	7.6	0.44	6.7	8.5	295	7.9	0.47	7.0	8.9
1997	338	7.7	0.43	6.9	8.6	312	8.1	0.47	7.2	9.0
1998	338	7.7	0.43	6.9	8.5	312	8.0	0.46	7.1	9.0
1999	360	8.1	0.43	7.3	9.0	330	8.4	0.47	7.5	9.4
2000	355	7.9	0.43	7.1	8.8	331	8.3	0.46	7.4	9.3
2001	339	7.4	0.41	6.6	8.2	317	7.8	0.44	7.0	8.7
2002	342	7.4	0.40	6.6	8.2	322	7.9	0.44	7.0	8.8
2003	378	8.0	0.42	7.2	8.9	358	8.7	0.46	7.8	9.6
2004	382	7.8	0.41	7.0	8.6	359	8.4	0.45	7.6	9.3
2005	370	7.6	0.40	6.8	8.4	345	8.0	0.44	7.2	8.9
2006	383	7.8	0.41	7.1	8.7	353	8.3	0.44	7.4	9.2
2007	391	7.7	0.39	6.9	8.5	350	7.9	0.43	7.1	8.8
2008	428	8.4	0.41	7.6	9.3	402	9.1	0.46	8.2	10.0
2009	413	7.9	0.39	7.1	8.7	375	8.2	0.43	7.4	9.1
2010	407	7.6	0.38	6.8	8.4	367	7.9	0.42	7.1	8.7
2011	398	7.3	0.37	6.6	8.1	364	7.7	0.41	6.9	8.5
2012	420	7.6	0.37	6.9	8.3	388	8.1	0.42	7.3	9.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary Bladder

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	695	5.7	0.23	5.3	6.2	46	1.8	0.30	1.3	2.5
1986	21	5.9	1.38	3.5	9.1	*	*	*	*	*
1987	26	7.1	1.50	4.5	10.6	*	*	*	*	*
1988	32	8.1	1.55	5.4	11.6	*	*	*	*	*
1989	24	6.6	1.42	4.1	9.8	*	*	*	*	*
1990	27	7.6	1.58	4.9	11.2	*	*	*	*	*
1991	32	8.5	1.60	5.7	12.1	*	*	*	*	*
1992	19	4.7	1.17	2.7	7.5	*	*	*	*	*
1993	23	5.1	1.15	3.1	7.8	*	*	*	*	*
1994	23	5.1	1.12	3.1	7.7	*	*	*	*	*
1995	25	5.8	1.23	3.6	8.6	*	*	*	*	*
1996	23	5.2	1.16	3.2	7.9	*	*	*	*	*
1997	25	5.1	1.08	3.2	7.6	*	*	*	*	*
1998	26	5.3	1.11	3.3	7.8	*	*	*	*	*
1999	28	6.1	1.21	3.9	8.9	*	*	*	*	*
2000	23	5.7	1.25	3.5	8.6	*	*	*	*	*
2001	21	4.4	1.04	2.6	6.8	*	*	*	*	*
2002	19	4.1	0.99	2.4	6.4	*	*	*	*	*
2003	18	4.1	1.05	2.3	6.6	*	*	*	*	*
2004	21	3.3	0.76	2.0	5.1	*	*	*	*	*
2005	24	5.7	1.22	3.5	8.5	*	*	*	*	*
2006	30	6.5	1.28	4.2	9.3	*	*	*	*	*
2007	36	7.2	1.29	4.9	10.1	*	*	*	*	*
2008	24	4.6	0.99	2.9	6.9	*	*	*	*	*
2009	36	7.0	1.23	4.8	9.7	*	*	*	*	*
2010	35	6.7	1.18	4.6	9.3	*	*	*	*	*
2011	31	5.5	1.03	3.7	7.9	*	*	*	*	*
2012	23	4.0	0.88	2.5	6.1	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary Bladder										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	4,731	2.4	0.03	2.3	2.5	4,010	2.3	0.04	2.2	2.4
1986	163	2.6	0.20	2.2	3.0	139	2.4	0.20	2.0	2.8
1987	153	2.4	0.20	2.0	2.8	131	2.3	0.20	1.9	2.7
1988	159	2.4	0.19	2.1	2.8	139	2.3	0.20	2.0	2.8
1989	159	2.4	0.20	2.1	2.9	134	2.3	0.20	1.9	2.7
1990	176	2.6	0.20	2.3	3.1	146	2.4	0.20	2.1	2.9
1991	146	2.1	0.18	1.8	2.5	127	2.1	0.18	1.7	2.5
1992	177	2.6	0.20	2.2	3.0	143	2.3	0.19	1.9	2.7
1993	149	2.1	0.17	1.8	2.5	132	2.1	0.18	1.7	2.5
1994	170	2.4	0.19	2.1	2.8	140	2.2	0.19	1.9	2.6
1995	151	2.1	0.17	1.8	2.5	138	2.2	0.19	1.8	2.6
1996	194	2.7	0.19	2.3	3.1	157	2.4	0.20	2.1	2.9
1997	167	2.3	0.18	2.0	2.7	141	2.2	0.18	1.8	2.6
1998	181	2.5	0.18	2.1	2.8	156	2.4	0.19	2.0	2.8
1999	200	2.7	0.19	2.3	3.1	171	2.6	0.20	2.2	3.0
2000	186	2.5	0.18	2.1	2.8	155	2.3	0.19	2.0	2.7
2001	181	2.4	0.18	2.0	2.8	150	2.3	0.19	1.9	2.7
2002	169	2.2	0.17	1.9	2.6	141	2.2	0.18	1.8	2.5
2003	179	2.4	0.18	2.0	2.8	152	2.3	0.19	2.0	2.7
2004	192	2.5	0.18	2.2	2.9	162	2.4	0.19	2.0	2.8
2005	190	2.5	0.18	2.1	2.9	163	2.5	0.20	2.1	2.9
2006	196	2.5	0.18	2.2	2.9	169	2.5	0.20	2.1	2.9
2007	183	2.3	0.18	2.0	2.7	154	2.3	0.19	1.9	2.7
2008	187	2.3	0.17	2.0	2.7	160	2.3	0.19	2.0	2.7
2009	184	2.3	0.17	2.0	2.7	156	2.3	0.19	1.9	2.7
2010	182	2.2	0.17	1.9	2.6	151	2.1	0.18	1.8	2.5
2011	181	2.2	0.17	1.9	2.6	151	2.2	0.19	1.9	2.6
2012	176	2.1	0.16	1.8	2.4	152	2.1	0.17	1.8	2.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Urinary Bladder

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	680	3.4	0.13	3.1	3.6	41	1.2	0.19	0.8	1.6
1986	24	3.9	0.82	2.5	5.9	*	*	*	*	*
1987	21	3.5	0.77	2.1	5.3	*	*	*	*	*
1988	20	3.2	0.73	1.9	4.9	*	*	*	*	*
1989	25	4.0	0.82	2.6	5.9	*	*	*	*	*
1990	29	4.6	0.87	3.1	6.6	*	*	*	*	*
1991	18	3.0	0.71	1.8	4.7	*	*	*	*	*
1992	34	5.1	0.89	3.5	7.2	*	*	*	*	*
1993	16	2.5	0.63	1.4	4.0	*	*	*	*	*
1994	28	4.1	0.78	2.7	5.9	*	*	*	*	*
1995	13	2.0	0.55	1.0	3.3	*	*	*	*	*
1996	34	4.7	0.81	3.2	6.5	*	*	*	*	*
1997	24	3.3	0.68	2.1	4.9	*	*	*	*	*
1998	25	3.3	0.67	2.2	4.9	*	*	*	*	*
1999	28	3.7	0.70	2.4	5.3	*	*	*	*	*
2000	30	3.9	0.72	2.6	5.6	*	*	*	*	*
2001	30	3.9	0.72	2.6	5.6	*	*	*	*	*
2002	28	3.6	0.69	2.4	5.2	*	*	*	*	*
2003	26	3.3	0.65	2.2	4.8	*	*	*	*	*
2004	27	3.3	0.63	2.2	4.8	*	*	*	*	*
2005	22	2.6	0.55	1.6	3.9	*	*	*	*	*
2006	25	3.0	0.60	1.9	4.4	*	*	*	*	*
2007	28	3.2	0.61	2.1	4.6	*	*	*	*	*
2008	25	2.9	0.57	1.8	4.2	*	*	*	*	*
2009	28	3.2	0.60	2.1	4.6	*	*	*	*	*
2010	28	3.1	0.59	2.0	4.4	*	*	*	*	*
2011	25	2.6	0.53	1.7	3.9	*	*	*	*	*
2012	19	2.0	0.47	1.2	3.2	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Kidney and Renal Pelvis

Both Sexes

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	14,276	4.4	0.04	4.3	4.5	12,487	4.5	0.04	4.4	4.5
1986	463	4.4	0.21	4.0	4.8	408	4.4	0.22	4.0	4.8
1987	485	4.6	0.21	4.2	5.0	442	4.7	0.23	4.3	5.2
1988	457	4.3	0.20	3.9	4.7	396	4.2	0.22	3.8	4.7
1989	501	4.7	0.21	4.2	5.1	448	4.7	0.23	4.3	5.2
1990	495	4.6	0.21	4.2	5.0	441	4.6	0.22	4.2	5.1
1991	495	4.5	0.21	4.1	5.0	444	4.6	0.22	4.2	5.1
1992	489	4.4	0.20	4.0	4.8	441	4.5	0.22	4.1	5.0
1993	530	4.7	0.20	4.3	5.1	470	4.8	0.22	4.4	5.2
1994	571	5.0	0.21	4.6	5.5	495	5.0	0.23	4.6	5.5
1995	526	4.6	0.20	4.2	5.0	461	4.6	0.22	4.2	5.1
1996	494	4.3	0.19	3.9	4.7	428	4.3	0.21	3.9	4.7
1997	509	4.4	0.19	4.0	4.8	428	4.3	0.21	3.9	4.7
1998	550	4.7	0.20	4.3	5.1	483	4.7	0.22	4.3	5.2
1999	490	4.1	0.19	3.8	4.5	425	4.1	0.20	3.8	4.6
2000	575	4.8	0.20	4.4	5.2	498	4.8	0.22	4.4	5.3
2001	565	4.7	0.20	4.3	5.1	489	4.7	0.21	4.3	5.1
2002	540	4.4	0.19	4.1	4.8	472	4.5	0.21	4.1	5.0
2003	566	4.6	0.19	4.2	5.0	474	4.5	0.21	4.1	4.9
2004	537	4.3	0.19	3.9	4.7	466	4.4	0.20	4.0	4.8
2005	554	4.4	0.19	4.0	4.7	488	4.5	0.20	4.1	4.9
2006	530	4.2	0.18	3.8	4.5	454	4.2	0.20	3.8	4.6
2007	543	4.2	0.18	3.8	4.5	466	4.2	0.20	3.8	4.6
2008	581	4.4	0.19	4.1	4.8	510	4.6	0.21	4.2	5.0
2009	545	4.1	0.18	3.7	4.4	473	4.2	0.19	3.8	4.6
2010	547	4.0	0.17	3.7	4.4	464	4.0	0.19	3.7	4.4
2011	556	4.0	0.17	3.7	4.4	509	4.4	0.20	4.0	4.8
2012	582	4.1	0.17	3.8	4.5	514	4.4	0.20	4.0	4.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Kidney and Renal Pelvis

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,666	4.5	0.11	4.3	4.7	123	1.6	0.16	1.3	1.9
1986	53	4.7	0.68	3.5	6.2	*	*	*	*	*
1987	41	3.5	0.57	2.5	4.8	*	*	*	*	*
1988	60	5.3	0.71	4.0	6.9	*	*	*	*	*
1989	52	4.3	0.61	3.2	5.6	*	*	*	*	*
1990	53	4.7	0.66	3.5	6.2	*	*	*	*	*
1991	49	4.1	0.60	3.0	5.5	*	*	*	*	*
1992	48	3.9	0.58	2.9	5.2	*	*	*	*	*
1993	58	4.6	0.61	3.4	5.9	*	*	*	*	*
1994	70	5.5	0.67	4.2	6.9	*	*	*	*	*
1995	64	5.2	0.67	4.0	6.7	*	*	*	*	*
1996	62	4.7	0.61	3.6	6.0	*	*	*	*	*
1997	77	5.8	0.69	4.6	7.3	*	*	*	*	*
1998	57	4.1	0.56	3.1	5.4	10	3.4	1.15	1.5	6.4
1999	62	4.5	0.58	3.4	5.8	*	*	*	*	*
2000	73	5.3	0.63	4.1	6.7	*	*	*	*	*
2001	71	5.0	0.61	3.9	6.3	*	*	*	*	*
2002	60	4.4	0.59	3.4	5.7	*	*	*	*	*
2003	87	5.9	0.65	4.7	7.3	*	*	*	*	*
2004	67	4.5	0.56	3.5	5.8	*	*	*	*	*
2005	61	4.0	0.52	3.0	5.1	*	*	*	*	*
2006	66	4.1	0.52	3.2	5.2	10	2.4	0.81	1.1	4.5
2007	69	4.5	0.56	3.5	5.8	*	*	*	*	*
2008	64	4.0	0.51	3.0	5.1	*	*	*	*	*
2009	62	3.9	0.50	3.0	5.0	10	1.9	0.62	0.8	3.5
2010	76	4.6	0.54	3.6	5.8	*	*	*	*	*
2011	42	2.6	0.41	1.9	3.5	*	*	*	*	*
2012	62	3.6	0.47	2.8	4.7	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Kidney and Renal Pelvis

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	8,700	6.4	0.07	6.3	6.5	7,607	6.5	0.08	6.3	6.6
1986	291	6.7	0.41	5.9	7.5	257	6.7	0.43	5.8	7.6
1987	286	6.6	0.41	5.9	7.5	259	6.8	0.44	6.0	7.7
1988	288	6.6	0.41	5.9	7.5	241	6.3	0.42	5.5	7.2
1989	296	6.8	0.41	6.0	7.6	269	7.0	0.44	6.1	7.9
1990	291	6.6	0.40	5.8	7.4	262	6.7	0.42	5.9	7.6
1991	297	6.6	0.40	5.9	7.4	266	6.8	0.43	5.9	7.6
1992	288	6.3	0.38	5.5	7.1	257	6.4	0.41	5.6	7.3
1993	314	6.7	0.39	6.0	7.5	279	6.8	0.42	6.0	7.7
1994	346	7.4	0.41	6.6	8.2	302	7.4	0.43	6.6	8.3
1995	302	6.4	0.37	5.6	7.1	260	6.3	0.39	5.5	7.1
1996	314	6.6	0.38	5.9	7.4	273	6.6	0.40	5.8	7.4
1997	297	6.1	0.36	5.5	6.9	250	6.0	0.38	5.3	6.8
1998	334	6.9	0.38	6.2	7.7	294	7.0	0.41	6.2	7.9
1999	294	6.0	0.35	5.3	6.7	249	5.8	0.37	5.1	6.6
2000	343	6.9	0.38	6.2	7.7	297	6.9	0.41	6.2	7.8
2001	336	6.6	0.37	5.9	7.4	295	6.8	0.40	6.0	7.6
2002	330	6.3	0.35	5.7	7.1	290	6.5	0.38	5.7	7.3
2003	360	6.8	0.37	6.1	7.6	303	6.7	0.39	6.0	7.5
2004	326	6.2	0.35	5.5	6.9	284	6.2	0.37	5.5	7.0
2005	365	6.9	0.36	6.2	7.6	317	6.9	0.39	6.2	7.7
2006	340	6.2	0.34	5.5	6.9	293	6.3	0.37	5.6	7.1
2007	324	5.7	0.32	5.1	6.4	281	5.8	0.35	5.1	6.5
2008	340	6.1	0.34	5.5	6.8	297	6.3	0.37	5.6	7.0
2009	350	6.1	0.33	5.4	6.7	304	6.2	0.36	5.5	6.9
2010	319	5.4	0.31	4.8	6.1	270	5.4	0.33	4.8	6.1
2011	371	6.3	0.34	5.7	7.0	340	6.8	0.37	6.1	7.6
2012	358	5.9	0.32	5.3	6.5	318	6.1	0.35	5.5	6.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Kidney and Renal Pelvis

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,013	6.6	0.22	6.2	7.1	80	2.4	0.31	1.8	3.1
1986	33	7.0	1.31	4.6	10.0	*	*	*	*	*
1987	25	5.1	1.07	3.2	7.6	*	*	*	*	*
1988	46	9.6	1.49	6.9	12.9	*	*	*	*	*
1989	26	5.5	1.15	3.4	8.1	*	*	*	*	*
1990	28	6.1	1.24	4.0	9.0	*	*	*	*	*
1991	31	5.8	1.07	3.9	8.2	*	*	*	*	*
1992	31	5.6	1.03	3.8	8.0	*	*	*	*	*
1993	33	6.4	1.17	4.3	9.1	*	*	*	*	*
1994	40	7.6	1.27	5.3	10.5	*	*	*	*	*
1995	41	8.2	1.37	5.7	11.3	*	*	*	*	*
1996	37	6.8	1.17	4.7	9.4	*	*	*	*	*
1997	44	7.8	1.27	5.5	10.6	*	*	*	*	*
1998	36	6.0	1.06	4.1	8.5	*	*	*	*	*
1999	44	7.8	1.25	5.5	10.6	*	*	*	*	*
2000	44	7.9	1.26	5.7	10.7	*	*	*	*	*
2001	39	6.3	1.06	4.4	8.7	*	*	*	*	*
2002	33	6.2	1.18	4.1	8.9	*	*	*	*	*
2003	54	8.9	1.28	6.5	11.7	*	*	*	*	*
2004	39	7.0	1.21	4.8	9.7	*	*	*	*	*
2005	45	7.5	1.21	5.4	10.2	*	*	*	*	*
2006	44	6.5	1.06	4.6	8.9	*	*	*	*	*
2007	36	6.2	1.11	4.2	8.7	*	*	*	*	*
2008	39	5.8	0.99	4.1	8.1	*	*	*	*	*
2009	37	5.7	0.99	3.9	7.9	*	*	*	*	*
2010	44	6.6	1.05	4.7	9.0	*	*	*	*	*
2011	27	4.2	0.85	2.7	6.2	*	*	*	*	*
2012	37	5.6	0.98	3.8	7.8	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Kidney and Renal Pelvis

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	5,576	3.0	0.04	2.9	3.0	4,880	3.0	0.04	2.9	3.1
1986	172	2.8	0.22	2.4	3.2	151	2.8	0.23	2.3	3.3
1987	199	3.2	0.23	2.8	3.7	183	3.3	0.25	2.8	3.8
1988	169	2.7	0.21	2.3	3.1	155	2.8	0.23	2.4	3.3
1989	205	3.3	0.23	2.9	3.8	179	3.2	0.25	2.8	3.8
1990	204	3.3	0.23	2.8	3.7	179	3.2	0.24	2.7	3.7
1991	198	3.1	0.22	2.7	3.5	178	3.1	0.24	2.7	3.6
1992	201	3.1	0.22	2.6	3.5	184	3.2	0.24	2.7	3.7
1993	216	3.3	0.22	2.8	3.7	191	3.3	0.24	2.9	3.8
1994	225	3.4	0.23	2.9	3.8	193	3.3	0.24	2.8	3.8
1995	224	3.4	0.23	3.0	3.9	201	3.5	0.25	3.0	4.1
1996	180	2.6	0.20	2.3	3.1	155	2.6	0.21	2.2	3.0
1997	212	3.1	0.21	2.7	3.6	178	3.0	0.23	2.6	3.5
1998	216	3.1	0.21	2.7	3.6	189	3.1	0.23	2.7	3.6
1999	196	2.8	0.20	2.4	3.3	176	2.9	0.22	2.5	3.4
2000	232	3.3	0.22	2.9	3.8	201	3.3	0.24	2.9	3.8
2001	229	3.2	0.22	2.8	3.7	194	3.2	0.23	2.8	3.7
2002	210	2.9	0.20	2.6	3.4	182	3.0	0.22	2.6	3.4
2003	206	2.8	0.20	2.5	3.3	171	2.7	0.21	2.3	3.2
2004	211	2.9	0.20	2.6	3.4	182	3.0	0.22	2.5	3.4
2005	189	2.6	0.19	2.2	3.0	171	2.7	0.21	2.3	3.2
2006	190	2.6	0.19	2.2	3.0	161	2.6	0.21	2.2	3.0
2007	219	2.9	0.20	2.5	3.4	185	2.9	0.22	2.5	3.4
2008	241	3.2	0.21	2.8	3.6	213	3.3	0.23	2.9	3.8
2009	195	2.5	0.18	2.2	2.9	169	2.6	0.20	2.2	3.0
2010	228	2.9	0.20	2.6	3.4	194	3.0	0.22	2.6	3.4
2011	185	2.3	0.17	2.0	2.7	169	2.5	0.20	2.1	2.9
2012	224	2.8	0.19	2.5	3.2	196	3.0	0.22	2.6	3.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Kidney and Renal Pelvis

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	653	3.0	0.12	2.8	3.2	43	1.0	0.16	0.7	1.4
1986	20	3.1	0.71	1.8	4.8	*	*	*	*	*
1987	16	2.4	0.61	1.3	3.8	*	*	*	*	*
1988	14	2.2	0.61	1.2	3.7	*	*	*	*	*
1989	26	3.7	0.73	2.4	5.4	*	*	*	*	*
1990	25	3.8	0.77	2.5	5.6	*	*	*	*	*
1991	18	2.8	0.65	1.6	4.3	*	*	*	*	*
1992	17	2.6	0.63	1.5	4.1	*	*	*	*	*
1993	25	3.4	0.68	2.2	4.9	*	*	*	*	*
1994	30	4.0	0.74	2.7	5.7	*	*	*	*	*
1995	23	3.2	0.68	2.0	4.8	*	*	*	*	*
1996	25	3.2	0.65	2.1	4.7	*	*	*	*	*
1997	33	4.3	0.76	3.0	6.1	*	*	*	*	*
1998	21	2.7	0.60	1.7	4.1	*	*	*	*	*
1999	18	2.3	0.54	1.3	3.6	*	*	*	*	*
2000	29	3.6	0.67	2.4	5.1	*	*	*	*	*
2001	32	3.9	0.70	2.7	5.5	*	*	*	*	*
2002	27	3.3	0.65	2.2	4.8	*	*	*	*	*
2003	33	3.8	0.68	2.6	5.4	*	*	*	*	*
2004	28	3.1	0.60	2.1	4.5	*	*	*	*	*
2005	16	1.8	0.44	1.0	2.9	*	*	*	*	*
2006	22	2.4	0.52	1.5	3.7	*	*	*	*	*
2007	33	3.6	0.63	2.5	5.1	*	*	*	*	*
2008	25	2.6	0.53	1.7	3.9	*	*	*	*	*
2009	25	2.7	0.54	1.7	4.0	*	*	*	*	*
2010	32	3.2	0.58	2.2	4.6	*	*	*	*	*
2011	15	1.6	0.41	0.9	2.6	*	*	*	*	*
2012	25	2.5	0.51	1.6	3.7	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Brain and Other Nervous System

Both Sexes

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	13,717	4.2	0.04	4.2	4.3	12,532	4.6	0.04	4.5	4.6
1986	491	4.6	0.21	4.2	5.0	451	4.8	0.23	4.4	5.3
1987	442	4.1	0.20	3.7	4.5	414	4.4	0.22	4.0	4.9
1988	473	4.4	0.20	4.0	4.8	440	4.7	0.23	4.3	5.2
1989	527	4.9	0.21	4.5	5.3	471	5.0	0.23	4.6	5.5
1990	505	4.6	0.21	4.2	5.0	467	4.9	0.23	4.5	5.4
1991	489	4.4	0.20	4.0	4.8	446	4.7	0.22	4.2	5.1
1992	537	4.7	0.21	4.4	5.2	489	5.0	0.23	4.6	5.5
1993	531	4.7	0.20	4.3	5.1	490	5.0	0.23	4.6	5.5
1994	519	4.5	0.20	4.1	4.9	478	4.9	0.22	4.4	5.3
1995	479	4.2	0.19	3.8	4.6	432	4.4	0.21	4.0	4.8
1996	471	4.1	0.19	3.7	4.4	438	4.4	0.21	4.0	4.9
1997	505	4.3	0.19	3.9	4.7	476	4.8	0.22	4.4	5.2
1998	475	4.0	0.18	3.7	4.4	438	4.4	0.21	4.0	4.8
1999	527	4.4	0.19	4.0	4.8	493	4.9	0.22	4.4	5.3
2000	502	4.2	0.19	3.8	4.6	458	4.5	0.21	4.1	4.9
2001	553	4.6	0.19	4.2	5.0	498	4.9	0.22	4.5	5.3
2002	467	3.8	0.18	3.5	4.2	421	4.1	0.20	3.7	4.5
2003	501	4.1	0.18	3.7	4.4	464	4.5	0.21	4.1	4.9
2004	511	4.1	0.18	3.7	4.5	467	4.5	0.21	4.1	4.9
2005	445	3.5	0.17	3.2	3.9	401	3.8	0.19	3.5	4.2
2006	510	4.0	0.18	3.7	4.4	469	4.5	0.21	4.1	4.9
2007	497	3.8	0.17	3.5	4.2	449	4.2	0.20	3.8	4.6
2008	501	3.8	0.17	3.5	4.1	458	4.2	0.20	3.8	4.6
2009	570	4.3	0.18	3.9	4.7	511	4.6	0.21	4.2	5.1
2010	545	4.0	0.18	3.7	4.4	488	4.4	0.20	4.0	4.8
2011	550	4.0	0.17	3.7	4.3	492	4.3	0.20	4.0	4.7
2012	594	4.2	0.18	3.9	4.6	533	4.6	0.20	4.2	5.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Brain and Other Nervous System

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,027	2.5	0.08	2.4	2.7	158	1.6	0.14	1.3	1.9
1986	39	3.1	0.52	2.1	4.2	*	*	*	*	*
1987	28	2.3	0.44	1.5	3.3	*	*	*	*	*
1988	29	2.3	0.44	1.5	3.3	*	*	*	*	*
1989	54	4.3	0.61	3.2	5.6	*	*	*	*	*
1990	35	2.7	0.48	1.8	3.8	*	*	*	*	*
1991	37	2.7	0.47	1.9	3.8	*	*	*	*	*
1992	47	3.3	0.50	2.4	4.4	*	*	*	*	*
1993	37	2.6	0.44	1.8	3.6	*	*	*	*	*
1994	41	2.9	0.47	2.1	4.0	*	*	*	*	*
1995	43	3.2	0.51	2.3	4.3	*	*	*	*	*
1996	29	2.0	0.37	1.3	2.8	*	*	*	*	*
1997	28	2.0	0.38	1.3	2.9	*	*	*	*	*
1998	33	2.1	0.38	1.5	3.0	*	*	*	*	*
1999	32	2.1	0.39	1.4	3.0	*	*	*	*	*
2000	34	2.3	0.40	1.6	3.2	10	2.8	0.94	1.3	5.3
2001	49	3.5	0.51	2.5	4.6	*	*	*	*	*
2002	39	2.5	0.40	1.7	3.4	*	*	*	*	*
2003	27	1.7	0.34	1.1	2.5	10	2.5	0.85	1.1	4.7
2004	34	2.4	0.41	1.6	3.3	10	2.3	0.81	1.0	4.4
2005	36	2.4	0.41	1.7	3.3	*	*	*	*	*
2006	37	2.3	0.38	1.6	3.1	*	*	*	*	*
2007	39	2.3	0.38	1.6	3.2	*	*	*	*	*
2008	35	1.9	0.32	1.3	2.6	*	*	*	*	*
2009	51	3.0	0.43	2.2	4.0	*	*	*	*	*
2010	47	2.6	0.39	1.9	3.5	10	1.5	0.49	0.7	2.9
2011	45	2.4	0.36	1.7	3.2	13	1.9	0.55	1.0	3.3
2012	42	2.4	0.38	1.7	3.3	19	3.0	0.71	1.7	4.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Brain and Other Nervous System

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	7,503	5.2	0.06	5.1	5.3	6,875	5.6	0.07	5.4	5.7
1986	269	5.8	0.37	5.1	6.5	250	6.1	0.40	5.4	7.0
1987	231	4.8	0.33	4.2	5.5	214	5.1	0.36	4.5	5.9
1988	248	5.3	0.35	4.7	6.0	231	5.6	0.38	4.9	6.4
1989	291	6.2	0.38	5.5	7.0	261	6.4	0.41	5.7	7.3
1990	277	5.6	0.34	4.9	6.3	259	6.0	0.38	5.3	6.8
1991	268	5.5	0.34	4.8	6.2	247	5.9	0.38	5.1	6.6
1992	300	6.0	0.36	5.3	6.7	274	6.3	0.39	5.6	7.1
1993	285	5.8	0.35	5.1	6.5	261	6.1	0.39	5.4	6.9
1994	269	5.3	0.33	4.7	6.0	246	5.7	0.37	5.0	6.4
1995	265	5.2	0.33	4.6	5.9	237	5.4	0.36	4.8	6.2
1996	261	5.0	0.31	4.4	5.6	253	5.7	0.36	5.0	6.4
1997	298	5.7	0.33	5.1	6.4	283	6.3	0.38	5.6	7.1
1998	262	5.0	0.32	4.4	5.7	239	5.4	0.35	4.7	6.1
1999	291	5.5	0.33	4.9	6.2	272	6.1	0.37	5.4	6.8
2000	265	4.9	0.31	4.3	5.6	246	5.4	0.34	4.7	6.1
2001	309	5.8	0.33	5.1	6.5	282	6.2	0.37	5.5	6.9
2002	263	4.8	0.30	4.2	5.4	239	5.1	0.33	4.5	5.8
2003	237	4.3	0.28	3.7	4.8	217	4.6	0.31	4.0	5.3
2004	277	5.0	0.31	4.4	5.7	253	5.4	0.34	4.7	6.1
2005	240	4.3	0.28	3.8	4.9	217	4.6	0.31	4.0	5.2
2006	291	5.1	0.30	4.5	5.7	271	5.6	0.35	5.0	6.4
2007	258	4.4	0.28	3.8	5.0	233	4.7	0.31	4.1	5.3
2008	284	4.8	0.29	4.2	5.4	262	5.3	0.33	4.6	6.0
2009	314	5.1	0.30	4.6	5.8	283	5.5	0.33	4.9	6.2
2010	304	5.0	0.29	4.5	5.6	270	5.3	0.33	4.7	6.0
2011	299	4.8	0.28	4.3	5.4	267	5.2	0.32	4.6	5.8
2012	347	5.4	0.29	4.8	6.0	308	5.7	0.33	5.0	6.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Brain and Other Nervous System

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	540	3.1	0.14	2.8	3.4	88	1.9	0.23	1.5	2.4
1986	19	3.4	0.85	2.0	5.5	*	*	*	*	*
1987	17	3.1	0.76	1.8	5.0	*	*	*	*	*
1988	14	3.1	0.92	1.6	5.4	*	*	*	*	*
1989	29	5.4	1.08	3.5	7.9	*	*	*	*	*
1990	16	2.4	0.66	1.3	4.1	*	*	*	*	*
1991	20	3.5	0.83	2.1	5.5	*	*	*	*	*
1992	26	4.4	0.96	2.7	6.7	*	*	*	*	*
1993	22	3.6	0.82	2.2	5.6	*	*	*	*	*
1994	23	3.8	0.85	2.3	5.8	*	*	*	*	*
1995	24	4.2	0.94	2.5	6.4	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	14	2.4	0.67	1.3	4.1	*	*	*	*	*
1998	20	2.9	0.67	1.7	4.5	*	*	*	*	*
1999	17	2.3	0.58	1.3	3.8	*	*	*	*	*
2000	15	2.5	0.69	1.3	4.2	*	*	*	*	*
2001	25	4.2	0.89	2.6	6.3	*	*	*	*	*
2002	19	3.2	0.76	1.8	5.0	*	*	*	*	*
2003	15	2.2	0.62	1.2	3.8	*	*	*	*	*
2004	17	2.8	0.73	1.6	4.6	*	*	*	*	*
2005	18	3.1	0.76	1.8	5.0	*	*	*	*	*
2006	17	2.3	0.59	1.3	3.8	*	*	*	*	*
2007	22	3.1	0.71	1.9	4.8	*	*	*	*	*
2008	17	2.0	0.50	1.1	3.2	*	*	*	*	*
2009	27	3.4	0.69	2.2	5.1	*	*	*	*	*
2010	27	3.3	0.68	2.1	5.0	*	*	*	*	*
2011	28	3.4	0.67	2.2	5.0	*	*	*	*	*
2012	25	3.3	0.70	2.1	5.0	14	5.2	1.54	2.6	9.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Brain and Other Nervous System

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	6,214	3.5	0.04	3.4	3.5	5,657	3.7	0.05	3.6	3.8
1986	222	3.7	0.25	3.2	4.2	201	3.9	0.28	3.3	4.5
1987	211	3.5	0.25	3.1	4.0	200	3.9	0.28	3.3	4.4
1988	225	3.7	0.25	3.3	4.3	209	4.0	0.28	3.5	4.6
1989	236	4.0	0.26	3.5	4.5	210	4.1	0.29	3.6	4.7
1990	228	3.8	0.25	3.3	4.3	208	4.0	0.28	3.4	4.6
1991	221	3.5	0.24	3.1	4.0	199	3.6	0.26	3.1	4.2
1992	237	3.7	0.24	3.2	4.2	215	3.9	0.27	3.4	4.5
1993	246	3.9	0.25	3.4	4.4	229	4.2	0.28	3.7	4.8
1994	250	3.9	0.25	3.4	4.4	232	4.2	0.28	3.7	4.8
1995	214	3.3	0.23	2.9	3.8	195	3.5	0.26	3.0	4.1
1996	210	3.3	0.23	2.8	3.7	185	3.4	0.25	2.9	3.9
1997	207	3.2	0.22	2.7	3.6	193	3.5	0.25	3.0	4.0
1998	213	3.2	0.22	2.8	3.7	199	3.6	0.26	3.1	4.1
1999	236	3.5	0.23	3.1	4.0	221	3.9	0.27	3.4	4.5
2000	237	3.6	0.23	3.1	4.1	212	3.8	0.26	3.3	4.3
2001	244	3.6	0.23	3.2	4.1	216	3.9	0.27	3.4	4.4
2002	204	3.0	0.21	2.6	3.5	182	3.2	0.24	2.8	3.7
2003	264	3.9	0.24	3.4	4.4	247	4.4	0.28	3.8	4.9
2004	234	3.4	0.22	3.0	3.9	214	3.8	0.26	3.3	4.3
2005	205	2.9	0.21	2.5	3.4	184	3.2	0.24	2.7	3.7
2006	219	3.2	0.22	2.8	3.6	198	3.5	0.25	3.0	4.0
2007	239	3.4	0.22	3.0	3.9	216	3.7	0.26	3.2	4.2
2008	217	3.0	0.20	2.6	3.4	196	3.2	0.23	2.8	3.7
2009	256	3.6	0.23	3.1	4.0	228	3.9	0.26	3.4	4.4
2010	241	3.3	0.21	2.9	3.7	218	3.6	0.25	3.1	4.1
2011	251	3.3	0.21	2.9	3.8	225	3.7	0.25	3.2	4.2
2012	247	3.3	0.21	2.9	3.7	225	3.7	0.25	3.2	4.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Brain and Other Nervous System

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	487	2.1	0.10	1.9	2.3	70	1.4	0.17	1.1	1.7
1986	20	2.7	0.64	1.6	4.3	*	*	*	*	*
1987	11	1.6	0.51	0.8	2.9	*	*	*	*	*
1988	15	1.9	0.50	1.0	3.1	*	*	*	*	*
1989	25	3.5	0.73	2.3	5.2	*	*	*	*	*
1990	19	2.7	0.64	1.6	4.2	*	*	*	*	*
1991	17	2.2	0.54	1.3	3.5	*	*	*	*	*
1992	21	2.6	0.57	1.6	3.9	*	*	*	*	*
1993	15	1.8	0.49	1.0	3.0	*	*	*	*	*
1994	18	2.3	0.55	1.3	3.6	*	*	*	*	*
1995	19	2.5	0.58	1.5	3.9	*	*	*	*	*
1996	22	2.7	0.57	1.6	4.0	*	*	*	*	*
1997	14	1.7	0.46	0.9	2.9	*	*	*	*	*
1998	13	1.5	0.43	0.8	2.6	*	*	*	*	*
1999	15	1.8	0.48	1.0	3.0	*	*	*	*	*
2000	19	2.2	0.51	1.3	3.4	*	*	*	*	*
2001	24	2.9	0.59	1.8	4.3	*	*	*	*	*
2002	20	2.1	0.48	1.3	3.3	*	*	*	*	*
2003	12	1.4	0.39	0.7	2.4	*	*	*	*	*
2004	17	2.0	0.49	1.2	3.2	*	*	*	*	*
2005	18	1.9	0.45	1.1	3.0	*	*	*	*	*
2006	20	2.1	0.48	1.3	3.3	*	*	*	*	*
2007	17	1.8	0.43	1.0	2.8	*	*	*	*	*
2008	18	1.8	0.42	1.1	2.8	*	*	*	*	*
2009	24	2.5	0.52	1.6	3.8	*	*	*	*	*
2010	20	2.0	0.44	1.2	3.0	*	*	*	*	*
2011	17	1.6	0.38	0.9	2.5	*	*	*	*	*
2012	17	1.8	0.43	1.0	2.8	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Endocrine System										
Both Sexes										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,636	0.8	0.02	0.8	0.8	2,260	0.8	0.02	0.8	0.9
1986	71	0.7	0.08	0.5	0.8	61	0.6	0.08	0.5	0.8
1987	87	0.8	0.09	0.6	1.0	75	0.8	0.09	0.6	1.0
1988	84	0.8	0.08	0.6	1.0	73	0.8	0.09	0.6	1.0
1989	97	0.9	0.09	0.7	1.1	86	0.9	0.10	0.7	1.2
1990	91	0.8	0.09	0.7	1.0	80	0.8	0.09	0.7	1.0
1991	89	0.8	0.09	0.6	1.0	79	0.8	0.09	0.6	1.0
1992	98	0.9	0.09	0.7	1.0	85	0.9	0.09	0.7	1.1
1993	108	0.9	0.09	0.8	1.1	97	1.0	0.10	0.8	1.2
1994	87	0.8	0.08	0.6	0.9	76	0.8	0.09	0.6	1.0
1995	81	0.7	0.08	0.6	0.9	72	0.7	0.09	0.6	0.9
1996	88	0.8	0.08	0.6	0.9	76	0.8	0.09	0.6	1.0
1997	112	1.0	0.09	0.8	1.1	95	0.9	0.10	0.8	1.2
1998	93	0.8	0.08	0.6	1.0	79	0.8	0.09	0.6	1.0
1999	94	0.8	0.08	0.6	1.0	78	0.8	0.09	0.6	1.0
2000	90	0.8	0.08	0.6	0.9	76	0.7	0.08	0.6	0.9
2001	95	0.8	0.08	0.6	1.0	77	0.8	0.09	0.6	0.9
2002	99	0.8	0.08	0.7	1.0	86	0.8	0.09	0.7	1.0
2003	106	0.9	0.08	0.7	1.0	90	0.9	0.09	0.7	1.1
2004	107	0.9	0.08	0.7	1.0	93	0.9	0.09	0.7	1.1
2005	105	0.8	0.08	0.7	1.0	93	0.9	0.09	0.7	1.1
2006	107	0.8	0.08	0.7	1.0	93	0.9	0.09	0.7	1.1
2007	89	0.7	0.07	0.5	0.8	71	0.6	0.08	0.5	0.8
2008	96	0.8	0.08	0.6	0.9	79	0.7	0.08	0.6	0.9
2009	111	0.9	0.08	0.7	1.0	91	0.9	0.09	0.7	1.0
2010	111	0.8	0.08	0.7	1.0	91	0.8	0.08	0.6	1.0
2011	109	0.8	0.08	0.7	1.0	104	0.9	0.09	0.7	1.1
2012	131	0.9	0.08	0.8	1.1	104	0.9	0.09	0.7	1.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Endocrine System										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	313	0.8	0.05	0.7	0.9	63	0.8	0.11	0.6	1.0
1986	10	0.8	0.26	0.3	1.5	*	*	*	*	*
1987	11	0.8	0.25	0.4	1.5	*	*	*	*	*
1988	10	0.7	0.25	0.3	1.4	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	13	0.8	0.25	0.4	1.5	*	*	*	*	*
1993	10	0.9	0.28	0.4	1.6	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	10	0.7	0.22	0.3	1.2	*	*	*	*	*
1997	15	1.0	0.28	0.6	1.7	*	*	*	*	*
1998	12	0.9	0.26	0.4	1.5	*	*	*	*	*
1999	13	0.7	0.20	0.4	1.2	*	*	*	*	*
2000	11	0.8	0.24	0.4	1.4	*	*	*	*	*
2001	15	1.0	0.26	0.5	1.6	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	15	1.0	0.28	0.6	1.7	*	*	*	*	*
2004	12	0.7	0.22	0.4	1.3	*	*	*	*	*
2005	10	0.6	0.21	0.3	1.2	*	*	*	*	*
2006	12	0.8	0.23	0.4	1.4	*	*	*	*	*
2007	15	0.9	0.24	0.5	1.5	*	*	*	*	*
2008	14	0.9	0.26	0.5	1.6	*	*	*	*	*
2009	15	1.0	0.25	0.5	1.6	*	*	*	*	*
2010	16	0.9	0.24	0.5	1.5	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	20	1.1	0.25	0.7	1.7	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Endocrine System										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,111	0.8	0.02	0.7	0.8	967	0.8	0.03	0.8	0.9
1986	25	0.5	0.11	0.3	0.8	23	0.5	0.12	0.3	0.8
1987	37	0.8	0.14	0.6	1.1	33	0.8	0.15	0.6	1.2
1988	37	0.8	0.14	0.6	1.1	31	0.8	0.15	0.5	1.1
1989	42	1.0	0.15	0.7	1.3	38	1.0	0.17	0.7	1.4
1990	37	0.8	0.14	0.6	1.2	32	0.8	0.15	0.6	1.2
1991	39	0.8	0.13	0.6	1.1	32	0.7	0.13	0.5	1.1
1992	41	0.8	0.12	0.5	1.0	36	0.8	0.13	0.6	1.1
1993	43	0.9	0.14	0.6	1.2	39	0.9	0.15	0.6	1.3
1994	37	0.8	0.13	0.5	1.0	31	0.7	0.13	0.5	1.0
1995	31	0.6	0.12	0.4	0.9	28	0.7	0.13	0.5	1.0
1996	32	0.6	0.11	0.4	0.9	29	0.7	0.12	0.4	1.0
1997	60	1.2	0.15	0.9	1.5	53	1.2	0.17	0.9	1.6
1998	40	0.8	0.13	0.6	1.1	35	0.8	0.14	0.6	1.1
1999	48	0.9	0.13	0.7	1.2	45	1.0	0.15	0.7	1.4
2000	36	0.7	0.11	0.5	0.9	33	0.7	0.13	0.5	1.0
2001	34	0.6	0.11	0.4	0.9	28	0.6	0.12	0.4	0.9
2002	42	0.8	0.13	0.6	1.1	38	0.9	0.14	0.6	1.2
2003	54	1.0	0.14	0.7	1.3	47	1.0	0.15	0.8	1.4
2004	38	0.6	0.11	0.5	0.9	34	0.7	0.12	0.5	1.0
2005	48	0.9	0.13	0.7	1.2	42	0.9	0.15	0.7	1.3
2006	37	0.7	0.12	0.5	1.0	34	0.8	0.13	0.5	1.1
2007	39	0.7	0.11	0.5	0.9	32	0.7	0.12	0.4	0.9
2008	39	0.7	0.11	0.5	0.9	32	0.7	0.12	0.5	0.9
2009	40	0.7	0.10	0.5	0.9	34	0.7	0.12	0.5	1.0
2010	41	0.7	0.11	0.5	1.0	32	0.7	0.12	0.4	0.9
2011	55	0.9	0.12	0.7	1.2	52	1.0	0.14	0.7	1.3
2012	59	0.9	0.12	0.7	1.2	44	0.8	0.12	0.6	1.1

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Endocrine System

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	112	0.6	0.06	0.5	0.8	32	0.8	0.16	0.5	1.2
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	10	1.4	0.46	0.6	2.6	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Endocrine System										
Females										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,525	0.8	0.02	0.8	0.9	1,293	0.8	0.02	0.8	0.9
1986	46	0.7	0.11	0.5	1.0	38	0.7	0.11	0.5	0.9
1987	50	0.8	0.12	0.6	1.1	42	0.8	0.13	0.6	1.1
1988	47	0.8	0.11	0.6	1.0	42	0.8	0.12	0.6	1.0
1989	55	0.9	0.12	0.7	1.2	48	0.9	0.13	0.7	1.2
1990	54	0.8	0.11	0.6	1.1	48	0.8	0.12	0.6	1.1
1991	50	0.8	0.11	0.6	1.0	47	0.8	0.13	0.6	1.1
1992	57	0.9	0.12	0.7	1.1	49	0.9	0.13	0.7	1.2
1993	65	1.0	0.13	0.8	1.3	58	1.1	0.14	0.8	1.4
1994	50	0.7	0.11	0.6	1.0	45	0.8	0.12	0.6	1.0
1995	50	0.7	0.10	0.5	1.0	44	0.7	0.11	0.5	1.0
1996	56	0.8	0.11	0.6	1.1	47	0.8	0.12	0.6	1.1
1997	52	0.8	0.11	0.6	1.0	42	0.7	0.11	0.5	1.0
1998	53	0.8	0.11	0.6	1.0	44	0.8	0.12	0.6	1.1
1999	46	0.7	0.10	0.5	0.9	33	0.6	0.10	0.4	0.8
2000	54	0.8	0.11	0.6	1.0	43	0.7	0.11	0.5	1.0
2001	61	0.9	0.12	0.7	1.2	49	0.9	0.12	0.6	1.1
2002	57	0.8	0.11	0.6	1.1	48	0.8	0.12	0.6	1.1
2003	52	0.7	0.10	0.6	1.0	43	0.7	0.11	0.5	1.0
2004	69	1.0	0.12	0.7	1.2	59	1.0	0.13	0.7	1.3
2005	57	0.8	0.11	0.6	1.0	51	0.8	0.12	0.6	1.1
2006	70	1.0	0.12	0.8	1.2	59	1.0	0.13	0.7	1.3
2007	50	0.7	0.10	0.5	0.9	39	0.6	0.10	0.5	0.9
2008	57	0.8	0.10	0.6	1.0	47	0.8	0.11	0.6	1.0
2009	71	1.0	0.12	0.8	1.2	57	0.9	0.13	0.7	1.2
2010	70	0.9	0.11	0.7	1.2	59	0.9	0.12	0.7	1.2
2011	54	0.7	0.10	0.5	0.9	52	0.8	0.11	0.6	1.1
2012	72	1.0	0.11	0.7	1.2	60	1.0	0.13	0.7	1.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Endocrine System										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	201	0.9	0.06	0.8	1.0	31	0.8	0.14	0.5	1.1
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	11	1.1	0.34	0.6	2.0	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	10	1.2	0.37	0.6	2.1	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	11	1.2	0.37	0.6	2.2	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	10	1.1	0.36	0.5	2.0	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	10	1.0	0.31	0.5	1.8	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Thyroid										
Both Sexes										
Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,524	0.5	0.01	0.5	0.5	1,335	0.5	0.01	0.5	0.5
1986	39	0.4	0.06	0.3	0.5	36	0.4	0.06	0.3	0.5
1987	48	0.5	0.07	0.3	0.6	42	0.5	0.07	0.3	0.6
1988	42	0.4	0.06	0.3	0.6	38	0.4	0.07	0.3	0.6
1989	51	0.5	0.07	0.4	0.7	45	0.5	0.07	0.4	0.7
1990	54	0.5	0.07	0.4	0.7	48	0.5	0.07	0.4	0.7
1991	53	0.5	0.07	0.4	0.6	50	0.5	0.07	0.4	0.7
1992	52	0.5	0.06	0.4	0.6	47	0.5	0.07	0.4	0.6
1993	54	0.5	0.07	0.4	0.6	51	0.5	0.07	0.4	0.7
1994	55	0.5	0.07	0.4	0.6	50	0.5	0.07	0.4	0.7
1995	49	0.4	0.06	0.3	0.6	45	0.5	0.07	0.3	0.6
1996	52	0.5	0.06	0.3	0.6	43	0.4	0.07	0.3	0.6
1997	58	0.5	0.07	0.4	0.6	50	0.5	0.07	0.4	0.7
1998	55	0.5	0.06	0.4	0.6	47	0.5	0.07	0.3	0.6
1999	41	0.3	0.05	0.3	0.5	36	0.4	0.06	0.2	0.5
2000	54	0.5	0.06	0.3	0.6	48	0.5	0.07	0.3	0.6
2001	53	0.4	0.06	0.3	0.6	46	0.4	0.07	0.3	0.6
2002	62	0.5	0.06	0.4	0.7	57	0.5	0.07	0.4	0.7
2003	57	0.5	0.06	0.4	0.6	48	0.5	0.07	0.3	0.6
2004	65	0.5	0.06	0.4	0.7	55	0.5	0.07	0.4	0.7
2005	64	0.5	0.06	0.4	0.7	58	0.5	0.07	0.4	0.7
2006	75	0.6	0.07	0.5	0.7	63	0.6	0.07	0.4	0.7
2007	56	0.4	0.06	0.3	0.6	43	0.4	0.06	0.3	0.5
2008	56	0.4	0.06	0.3	0.6	46	0.4	0.06	0.3	0.6
2009	69	0.5	0.07	0.4	0.7	56	0.5	0.07	0.4	0.7
2010	64	0.5	0.06	0.4	0.6	53	0.5	0.06	0.3	0.6
2011	72	0.5	0.06	0.4	0.7	70	0.6	0.07	0.5	0.8
2012	74	0.5	0.06	0.4	0.7	64	0.5	0.07	0.4	0.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Thyroid										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	152	0.4	0.04	0.4	0.5	37	0.5	0.09	0.4	0.7
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	10	0.7	0.21	0.3	1.2	*	*	*	*	*
2007	10	0.7	0.21	0.3	1.2	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Thyroid										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	559	0.4	0.02	0.4	0.5	503	0.4	0.02	0.4	0.5
1986	14	0.3	0.09	0.2	0.5	14	0.4	0.10	0.2	0.6
1987	18	0.4	0.10	0.2	0.6	16	0.4	0.10	0.2	0.6
1988	16	0.4	0.11	0.2	0.7	13	0.4	0.11	0.2	0.7
1989	19	0.5	0.12	0.3	0.8	18	0.5	0.13	0.3	0.8
1990	17	0.4	0.11	0.3	0.7	15	0.4	0.12	0.2	0.7
1991	20	0.5	0.10	0.3	0.7	17	0.4	0.10	0.2	0.7
1992	15	0.3	0.08	0.2	0.5	14	0.3	0.09	0.2	0.6
1993	20	0.4	0.10	0.3	0.7	20	0.5	0.11	0.3	0.8
1994	20	0.4	0.10	0.3	0.7	18	0.4	0.10	0.3	0.7
1995	15	0.3	0.09	0.2	0.5	14	0.4	0.10	0.2	0.6
1996	17	0.3	0.08	0.2	0.6	15	0.4	0.09	0.2	0.6
1997	27	0.6	0.11	0.4	0.8	23	0.5	0.11	0.3	0.8
1998	23	0.5	0.10	0.3	0.7	22	0.5	0.11	0.3	0.8
1999	23	0.5	0.10	0.3	0.7	22	0.5	0.11	0.3	0.8
2000	20	0.4	0.09	0.2	0.6	20	0.5	0.10	0.3	0.7
2001	14	0.3	0.08	0.2	0.5	13	0.3	0.08	0.2	0.5
2002	27	0.6	0.11	0.4	0.8	26	0.6	0.12	0.4	0.9
2003	20	0.4	0.09	0.2	0.6	17	0.4	0.09	0.2	0.6
2004	15	0.3	0.07	0.2	0.4	14	0.3	0.08	0.2	0.5
2005	26	0.5	0.10	0.3	0.8	24	0.6	0.11	0.4	0.8
2006	23	0.4	0.09	0.3	0.7	20	0.4	0.10	0.3	0.7
2007	23	0.4	0.09	0.3	0.6	19	0.4	0.09	0.2	0.6
2008	20	0.4	0.08	0.2	0.6	17	0.4	0.09	0.2	0.6
2009	19	0.3	0.07	0.2	0.5	17	0.3	0.08	0.2	0.6
2010	21	0.4	0.08	0.2	0.6	18	0.4	0.09	0.2	0.6
2011	32	0.5	0.10	0.4	0.8	30	0.6	0.11	0.4	0.8
2012	35	0.6	0.10	0.4	0.8	27	0.5	0.10	0.3	0.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Thyroid										
Males										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	37	0.3	0.04	0.2	0.4	19	0.6	0.15	0.3	0.9
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Thyroid										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	965	0.5	0.02	0.5	0.5	832	0.5	0.02	0.5	0.5
1986	25	0.4	0.08	0.3	0.6	22	0.4	0.08	0.2	0.6
1987	30	0.5	0.09	0.3	0.7	26	0.5	0.10	0.3	0.7
1988	26	0.4	0.08	0.3	0.6	25	0.4	0.09	0.3	0.6
1989	32	0.5	0.09	0.4	0.7	27	0.5	0.10	0.3	0.7
1990	37	0.6	0.09	0.4	0.8	33	0.6	0.10	0.4	0.8
1991	33	0.5	0.09	0.4	0.7	33	0.6	0.11	0.4	0.8
1992	37	0.6	0.09	0.4	0.8	33	0.6	0.10	0.4	0.8
1993	34	0.5	0.09	0.4	0.7	31	0.5	0.10	0.4	0.8
1994	35	0.5	0.09	0.4	0.7	32	0.5	0.09	0.4	0.7
1995	34	0.5	0.08	0.3	0.7	31	0.5	0.09	0.3	0.7
1996	35	0.5	0.09	0.4	0.7	28	0.5	0.09	0.3	0.7
1997	31	0.4	0.08	0.3	0.6	27	0.4	0.09	0.3	0.7
1998	32	0.5	0.08	0.3	0.7	25	0.4	0.08	0.3	0.6
1999	18	0.3	0.06	0.2	0.4	14	0.2	0.06	0.1	0.4
2000	34	0.5	0.08	0.3	0.7	28	0.5	0.09	0.3	0.7
2001	39	0.6	0.09	0.4	0.8	33	0.5	0.10	0.4	0.8
2002	35	0.5	0.08	0.3	0.7	31	0.5	0.09	0.3	0.7
2003	37	0.5	0.09	0.4	0.7	31	0.5	0.09	0.3	0.7
2004	50	0.7	0.09	0.5	0.9	41	0.6	0.10	0.4	0.9
2005	38	0.5	0.09	0.4	0.7	34	0.5	0.09	0.4	0.8
2006	52	0.7	0.10	0.5	1.0	43	0.7	0.11	0.5	1.0
2007	33	0.5	0.08	0.3	0.6	24	0.4	0.08	0.2	0.6
2008	36	0.5	0.08	0.3	0.7	29	0.5	0.09	0.3	0.7
2009	50	0.7	0.10	0.5	0.9	39	0.6	0.10	0.4	0.9
2010	43	0.5	0.08	0.4	0.7	35	0.5	0.09	0.4	0.7
2011	40	0.5	0.08	0.4	0.7	40	0.6	0.09	0.4	0.8
2012	39	0.5	0.08	0.4	0.7	37	0.6	0.10	0.4	0.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Thyroid										
Females										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	115	0.6	0.05	0.5	0.7	18	0.5	0.12	0.3	0.8
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lymphoma										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	26,528	8.2	0.05	8.1	8.3	24,079	8.6	0.06	8.5	8.7
1986	838	7.9	0.28	7.3	8.4	774	8.2	0.30	7.6	8.8
1987	836	7.8	0.28	7.3	8.4	759	8.1	0.30	7.5	8.7
1988	893	8.4	0.28	7.8	8.9	817	8.7	0.31	8.1	9.3
1989	937	8.6	0.29	8.1	9.2	860	9.0	0.31	8.4	9.6
1990	935	8.6	0.28	8.1	9.2	851	8.9	0.31	8.3	9.5
1991	1,002	9.1	0.29	8.5	9.6	918	9.5	0.31	8.9	10.1
1992	1,032	9.2	0.29	8.7	9.8	945	9.6	0.32	9.0	10.3
1993	1,041	9.2	0.29	8.7	9.8	945	9.6	0.31	9.0	10.2
1994	1,092	9.6	0.29	9.0	10.2	981	9.9	0.32	9.3	10.5
1995	1,138	9.9	0.29	9.3	10.5	1,030	10.2	0.32	9.6	10.9
1996	1,118	9.6	0.29	9.1	10.2	1,009	10.0	0.31	9.4	10.6
1997	1,099	9.4	0.28	8.8	10.0	1,001	9.8	0.31	9.2	10.5
1998	1,080	9.1	0.28	8.6	9.7	970	9.4	0.30	8.8	10.0
1999	1,118	9.4	0.28	8.8	10.0	1,026	9.9	0.31	9.3	10.5
2000	1,065	8.9	0.27	8.3	9.4	977	9.4	0.30	8.8	10.0
2001	1,041	8.6	0.27	8.1	9.1	950	9.1	0.29	8.5	9.7
2002	1,008	8.2	0.26	7.7	8.8	911	8.6	0.29	8.1	9.2
2003	979	7.9	0.25	7.4	8.4	895	8.4	0.28	7.8	8.9
2004	991	8.0	0.25	7.5	8.5	908	8.5	0.28	8.0	9.1
2005	1,000	8.0	0.25	7.5	8.5	905	8.3	0.28	7.8	8.9
2006	943	7.4	0.24	7.0	7.9	855	7.9	0.27	7.3	8.4
2007	868	6.7	0.23	6.3	7.2	783	7.1	0.26	6.6	7.6
2008	904	6.9	0.23	6.5	7.4	817	7.3	0.26	6.8	7.9
2009	952	7.2	0.24	6.7	7.7	838	7.4	0.26	6.9	8.0
2010	882	6.6	0.22	6.2	7.0	793	7.0	0.25	6.5	7.5
2011	883	6.5	0.22	6.1	6.9	786	6.8	0.25	6.3	7.3
2012	853	6.2	0.22	5.8	6.7	775	6.7	0.24	6.2	7.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lymphoma										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,156	5.7	0.12	5.4	5.9	293	3.8	0.24	3.4	4.3
1986	60	5.3	0.71	4.0	6.9	*	*	*	*	*
1987	72	6.0	0.74	4.6	7.6	*	*	*	*	*
1988	65	5.1	0.65	3.9	6.6	11	8.7	3.06	3.8	16.4
1989	68	5.3	0.67	4.1	6.8	*	*	*	*	*
1990	81	6.8	0.79	5.4	8.5	*	*	*	*	*
1991	79	6.0	0.71	4.7	7.6	*	*	*	*	*
1992	85	6.6	0.75	5.2	8.2	*	*	*	*	*
1993	91	7.0	0.76	5.6	8.6	*	*	*	*	*
1994	101	7.7	0.79	6.2	9.4	10	5.5	1.90	2.4	10.3
1995	98	7.4	0.78	6.0	9.1	10	5.5	1.94	2.4	10.3
1996	102	7.6	0.78	6.2	9.3	*	*	*	*	*
1997	83	6.0	0.68	4.8	7.5	15	5.7	1.61	3.0	9.7
1998	101	7.4	0.76	6.0	9.1	*	*	*	*	*
1999	86	6.2	0.68	4.9	7.7	*	*	*	*	*
2000	75	5.3	0.63	4.1	6.7	13	3.6	1.16	1.7	6.6
2001	80	5.5	0.62	4.3	6.8	11	2.6	0.84	1.2	4.9
2002	82	5.6	0.63	4.4	6.9	15	4.3	1.19	2.3	7.2
2003	65	4.4	0.56	3.4	5.7	19	6.9	1.72	4.0	10.9
2004	67	4.4	0.55	3.4	5.7	16	4.2	1.12	2.3	6.9
2005	77	5.1	0.59	4.0	6.4	18	5.5	1.38	3.2	8.8
2006	76	5.0	0.58	3.9	6.2	12	3.1	0.93	1.5	5.4
2007	69	4.5	0.56	3.5	5.7	16	3.9	1.04	2.1	6.4
2008	74	4.9	0.58	3.8	6.1	13	3.0	0.87	1.5	5.1
2009	94	5.8	0.61	4.6	7.1	20	4.6	1.10	2.7	7.2
2010	76	4.5	0.53	3.6	5.7	13	2.4	0.71	1.2	4.2
2011	83	4.9	0.55	3.9	6.1	14	3.0	0.84	1.6	5.0
2012	66	3.9	0.49	3.0	5.0	12	2.5	0.77	1.3	4.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.
*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*
** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lymphoma										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	13,905	10.5	0.09	10.3	10.7	12,542	10.9	0.10	10.7	11.1
1986	422	9.4	0.48	8.5	10.4	382	9.7	0.51	8.7	10.7
1987	431	9.6	0.48	8.7	10.6	380	9.7	0.51	8.7	10.7
1988	457	10.4	0.51	9.4	11.5	417	10.9	0.55	9.8	12.0
1989	488	10.8	0.51	9.8	11.8	442	11.2	0.55	10.1	12.3
1990	527	11.8	0.53	10.7	12.9	481	12.2	0.58	11.1	13.4
1991	520	11.3	0.51	10.3	12.3	472	11.7	0.56	10.7	12.9
1992	550	12.0	0.53	10.9	13.0	495	12.3	0.57	11.2	13.5
1993	513	11.0	0.50	10.1	12.1	461	11.4	0.54	10.3	12.5
1994	571	12.2	0.53	11.2	13.3	508	12.5	0.57	11.4	13.7
1995	585	12.4	0.52	11.4	13.4	529	12.8	0.57	11.7	14.0
1996	581	12.1	0.52	11.2	13.2	515	12.3	0.55	11.3	13.5
1997	560	11.5	0.50	10.6	12.5	496	11.8	0.54	10.7	12.9
1998	584	12.2	0.51	11.2	13.2	519	12.5	0.56	11.4	13.6
1999	603	12.4	0.51	11.4	13.5	551	13.1	0.56	12.0	14.2
2000	570	11.5	0.49	10.5	12.4	520	12.1	0.54	11.1	13.2
2001	547	10.9	0.47	10.0	11.9	502	11.6	0.52	10.6	12.7
2002	513	10.4	0.47	9.5	11.3	468	10.9	0.51	9.9	12.0
2003	494	9.8	0.45	8.9	10.7	446	10.2	0.49	9.3	11.2
2004	515	10.1	0.45	9.3	11.1	474	10.8	0.50	9.9	11.9
2005	517	10.2	0.46	9.3	11.1	467	10.6	0.50	9.7	11.7
2006	515	9.8	0.44	8.9	10.7	465	10.3	0.48	9.4	11.3
2007	456	8.8	0.42	8.0	9.6	416	9.3	0.46	8.4	10.2
2008	467	8.8	0.41	8.0	9.6	422	9.3	0.46	8.4	10.2
2009	480	8.8	0.41	8.0	9.6	424	9.1	0.45	8.2	10.0
2010	482	8.7	0.40	7.9	9.5	430	9.1	0.45	8.3	10.0
2011	475	8.5	0.40	7.8	9.3	422	8.9	0.44	8.1	9.8
2012	482	8.4	0.39	7.6	9.2	438	9.0	0.44	8.1	9.9

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lymphoma										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,197	7.4	0.23	7.0	7.9	166	5.0	0.44	4.2	5.9
1986	37	7.7	1.30	5.3	10.6	*	*	*	*	*
1987	48	9.2	1.43	6.6	12.4	*	*	*	*	*
1988	34	6.0	1.07	4.1	8.5	*	*	*	*	*
1989	40	7.0	1.19	4.9	9.8	*	*	*	*	*
1990	44	8.7	1.44	6.1	11.9	*	*	*	*	*
1991	44	7.8	1.28	5.5	10.7	*	*	*	*	*
1992	55	10.3	1.52	7.5	13.7	*	*	*	*	*
1993	49	8.6	1.32	6.2	11.5	*	*	*	*	*
1994	56	9.7	1.38	7.2	12.8	*	*	*	*	*
1995	52	9.0	1.35	6.5	12.0	*	*	*	*	*
1996	62	11.8	1.65	8.8	15.4	*	*	*	*	*
1997	51	8.2	1.21	6.0	11.0	13	11.4	3.56	5.6	20.4
1998	58	10.0	1.38	7.4	13.0	*	*	*	*	*
1999	48	8.2	1.26	5.9	11.0	*	*	*	*	*
2000	41	6.8	1.15	4.7	9.4	*	*	*	*	*
2001	39	6.2	1.04	4.3	8.5	*	*	*	*	*
2002	35	5.7	1.02	3.8	8.0	10	5.4	1.95	2.3	10.6
2003	38	6.2	1.10	4.3	8.7	10	8.3	2.91	3.6	15.5
2004	34	5.5	1.01	3.7	7.8	*	*	*	*	*
2005	42	6.7	1.10	4.8	9.2	*	*	*	*	*
2006	41	6.3	1.04	4.4	8.6	*	*	*	*	*
2007	35	6.0	1.10	4.0	8.5	*	*	*	*	*
2008	38	5.9	1.03	4.1	8.2	*	*	*	*	*
2009	44	6.8	1.10	4.8	9.2	12	7.3	2.30	3.5	13.0
2010	46	6.3	0.98	4.6	8.6	*	*	*	*	*
2011	45	6.4	1.00	4.6	8.6	*	*	*	*	*
2012	41	5.8	0.95	4.1	8.0	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lymphoma										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	12,623	6.6	0.06	6.5	6.7	11,537	7.0	0.07	6.8	7.1
1986	416	6.6	0.33	6.0	7.3	392	7.1	0.36	6.4	7.9
1987	405	6.5	0.33	5.8	7.1	379	6.8	0.36	6.1	7.6
1988	436	6.9	0.33	6.3	7.6	400	7.1	0.36	6.5	7.9
1989	449	7.1	0.34	6.4	7.8	418	7.4	0.37	6.7	8.2
1990	408	6.2	0.31	5.6	6.9	370	6.4	0.33	5.7	7.0
1991	482	7.3	0.34	6.7	8.0	446	7.6	0.37	6.9	8.4
1992	482	7.3	0.33	6.6	7.9	450	7.7	0.37	7.0	8.5
1993	528	7.8	0.34	7.2	8.5	484	8.2	0.38	7.4	8.9
1994	521	7.8	0.34	7.1	8.5	473	8.1	0.38	7.4	8.9
1995	553	8.1	0.34	7.4	8.8	501	8.3	0.38	7.6	9.1
1996	537	7.8	0.34	7.2	8.5	494	8.2	0.37	7.5	9.0
1997	539	7.7	0.34	7.1	8.4	505	8.3	0.37	7.6	9.1
1998	496	7.0	0.32	6.4	7.7	451	7.4	0.35	6.7	8.1
1999	515	7.3	0.32	6.6	7.9	475	7.7	0.36	7.0	8.4
2000	495	7.0	0.32	6.4	7.6	457	7.4	0.35	6.8	8.2
2001	494	6.9	0.31	6.3	7.5	448	7.2	0.34	6.5	7.9
2002	495	6.8	0.31	6.2	7.5	443	7.1	0.34	6.4	7.8
2003	485	6.5	0.30	5.9	7.1	449	7.0	0.33	6.3	7.7
2004	476	6.5	0.30	6.0	7.2	434	6.9	0.34	6.3	7.6
2005	483	6.5	0.30	5.9	7.1	438	6.8	0.33	6.1	7.4
2006	428	5.7	0.28	5.2	6.3	390	6.1	0.31	5.5	6.7
2007	412	5.4	0.27	4.8	5.9	367	5.6	0.30	5.0	6.2
2008	437	5.6	0.27	5.1	6.2	395	6.0	0.31	5.4	6.6
2009	472	6.1	0.28	5.5	6.7	414	6.2	0.31	5.6	6.9
2010	400	5.1	0.26	4.6	5.6	363	5.4	0.29	4.8	6.0
2011	408	5.1	0.26	4.6	5.6	364	5.3	0.29	4.8	5.9
2012	371	4.6	0.25	4.2	5.1	337	5.0	0.28	4.4	5.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Lymphoma										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	959	4.4	0.14	4.1	4.7	127	2.9	0.28	2.4	3.5
1986	23	3.5	0.75	2.2	5.3	*	*	*	*	*
1987	24	3.6	0.76	2.3	5.4	*	*	*	*	*
1988	31	4.4	0.80	2.9	6.2	*	*	*	*	*
1989	28	4.0	0.77	2.6	5.8	*	*	*	*	*
1990	37	5.4	0.91	3.8	7.5	*	*	*	*	*
1991	35	4.7	0.82	3.3	6.6	*	*	*	*	*
1992	30	4.1	0.77	2.7	5.9	*	*	*	*	*
1993	42	5.7	0.89	4.1	7.7	*	*	*	*	*
1994	45	6.0	0.92	4.4	8.1	*	*	*	*	*
1995	46	6.2	0.92	4.5	8.2	*	*	*	*	*
1996	40	5.0	0.81	3.6	6.9	*	*	*	*	*
1997	32	4.2	0.75	2.8	5.9	*	*	*	*	*
1998	43	5.5	0.85	4.0	7.4	*	*	*	*	*
1999	38	4.7	0.78	3.3	6.5	*	*	*	*	*
2000	34	4.2	0.72	2.9	5.8	*	*	*	*	*
2001	41	4.9	0.77	3.5	6.6	*	*	*	*	*
2002	47	5.5	0.81	4.0	7.3	*	*	*	*	*
2003	27	3.2	0.63	2.1	4.7	*	*	*	*	*
2004	33	3.8	0.66	2.6	5.3	*	*	*	*	*
2005	35	3.9	0.67	2.7	5.5	10	4.8	1.60	2.2	8.9
2006	35	4.0	0.67	2.7	5.5	*	*	*	*	*
2007	34	3.7	0.63	2.5	5.1	11	4.4	1.39	2.1	8.0
2008	36	4.1	0.68	2.8	5.6	*	*	*	*	*
2009	50	5.2	0.75	3.9	6.9	*	*	*	*	*
2010	30	3.1	0.58	2.1	4.5	*	*	*	*	*
2011	38	3.9	0.63	2.7	5.3	*	*	*	*	*
2012	25	2.6	0.52	1.7	3.8	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Hodgkin Lymphoma										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,634	0.5	0.01	0.5	0.5	1,398	0.5	0.01	0.5	0.5
1986	103	0.9	0.09	0.7	1.1	94	1.0	0.10	0.8	1.2
1987	90	0.8	0.09	0.6	1.0	77	0.8	0.09	0.6	1.0
1988	73	0.7	0.08	0.5	0.8	62	0.7	0.09	0.5	0.9
1989	70	0.6	0.07	0.5	0.8	60	0.6	0.08	0.5	0.8
1990	71	0.6	0.07	0.5	0.8	59	0.6	0.08	0.4	0.8
1991	72	0.6	0.07	0.5	0.8	59	0.6	0.08	0.5	0.8
1992	67	0.6	0.07	0.4	0.7	56	0.6	0.08	0.4	0.7
1993	65	0.6	0.07	0.4	0.7	56	0.6	0.08	0.4	0.7
1994	68	0.6	0.07	0.5	0.7	56	0.6	0.07	0.4	0.7
1995	58	0.5	0.07	0.4	0.6	48	0.5	0.07	0.4	0.6
1996	60	0.5	0.07	0.4	0.7	51	0.5	0.07	0.4	0.7
1997	60	0.5	0.07	0.4	0.7	53	0.5	0.07	0.4	0.7
1998	61	0.5	0.07	0.4	0.7	49	0.5	0.07	0.4	0.6
1999	57	0.5	0.06	0.4	0.6	51	0.5	0.07	0.4	0.7
2000	61	0.5	0.06	0.4	0.6	53	0.5	0.07	0.4	0.7
2001	54	0.4	0.06	0.3	0.6	48	0.5	0.07	0.3	0.6
2002	65	0.5	0.07	0.4	0.7	57	0.6	0.07	0.4	0.7
2003	52	0.4	0.06	0.3	0.6	46	0.4	0.07	0.3	0.6
2004	56	0.4	0.06	0.3	0.6	49	0.5	0.07	0.3	0.6
2005	50	0.4	0.06	0.3	0.5	41	0.4	0.06	0.3	0.5
2006	58	0.5	0.06	0.4	0.6	47	0.5	0.07	0.3	0.6
2007	48	0.4	0.05	0.3	0.5	44	0.4	0.06	0.3	0.6
2008	50	0.4	0.06	0.3	0.5	39	0.4	0.06	0.3	0.5
2009	49	0.4	0.05	0.3	0.5	43	0.4	0.06	0.3	0.5
2010	35	0.3	0.05	0.2	0.4	31	0.3	0.05	0.2	0.4
2011	38	0.3	0.05	0.2	0.4	32	0.3	0.05	0.2	0.4
2012	43	0.3	0.05	0.2	0.4	37	0.3	0.06	0.2	0.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Hodgkin Lymphoma

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	216	0.5	0.03	0.4	0.6	20	0.2	0.04	0.1	0.3
1986	*	*	*	*	*	*	*	*	*	*
1987	12	0.7	0.22	0.4	1.3	*	*	*	*	*
1988	10	0.6	0.21	0.3	1.2	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	12	0.8	0.24	0.4	1.4	*	*	*	*	*
1991	12	0.7	0.21	0.4	1.3	*	*	*	*	*
1992	11	0.6	0.20	0.3	1.2	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	10	0.6	0.20	0.3	1.2	*	*	*	*	*
1995	10	0.6	0.20	0.3	1.1	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	11	0.8	0.23	0.4	1.4	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	10	0.6	0.20	0.3	1.1	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Hodgkin Lymphoma

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	915	0.6	0.02	0.6	0.7	779	0.6	0.02	0.6	0.7
1986	47	0.9	0.14	0.7	1.2	43	1.0	0.15	0.7	1.3
1987	56	1.0	0.14	0.8	1.3	45	1.0	0.15	0.7	1.3
1988	40	0.8	0.13	0.6	1.1	35	0.8	0.14	0.6	1.1
1989	40	0.8	0.13	0.6	1.1	35	0.8	0.14	0.6	1.2
1990	46	0.9	0.13	0.6	1.2	38	0.8	0.14	0.6	1.2
1991	40	0.8	0.13	0.5	1.1	33	0.7	0.13	0.5	1.1
1992	36	0.7	0.12	0.5	1.0	29	0.7	0.12	0.4	0.9
1993	25	0.5	0.10	0.3	0.7	19	0.4	0.10	0.3	0.7
1994	37	0.7	0.11	0.5	0.9	31	0.7	0.12	0.4	0.9
1995	35	0.6	0.11	0.4	0.9	30	0.7	0.12	0.4	0.9
1996	35	0.7	0.12	0.5	1.0	31	0.7	0.13	0.5	1.0
1997	30	0.6	0.11	0.4	0.9	24	0.6	0.12	0.4	0.9
1998	43	0.8	0.12	0.6	1.1	33	0.7	0.13	0.5	1.0
1999	36	0.7	0.12	0.5	1.0	33	0.8	0.13	0.5	1.1
2000	34	0.6	0.11	0.4	0.9	29	0.6	0.12	0.4	0.9
2001	31	0.6	0.10	0.4	0.8	30	0.6	0.12	0.4	0.9
2002	38	0.7	0.12	0.5	1.0	32	0.7	0.13	0.5	1.0
2003	28	0.5	0.10	0.3	0.7	24	0.5	0.11	0.3	0.8
2004	37	0.7	0.11	0.5	0.9	35	0.7	0.13	0.5	1.0
2005	24	0.4	0.09	0.3	0.6	18	0.4	0.09	0.2	0.6
2006	35	0.6	0.11	0.4	0.9	28	0.6	0.11	0.4	0.9
2007	21	0.4	0.08	0.2	0.6	20	0.4	0.10	0.3	0.7
2008	29	0.5	0.09	0.3	0.7	22	0.5	0.10	0.3	0.7
2009	25	0.4	0.09	0.3	0.6	23	0.5	0.10	0.3	0.7
2010	18	0.3	0.08	0.2	0.5	16	0.4	0.09	0.2	0.6
2011	21	0.4	0.08	0.2	0.6	20	0.4	0.09	0.3	0.6
2012	28	0.5	0.09	0.3	0.7	23	0.5	0.10	0.3	0.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Hodgkin Lymphoma

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	122	0.6	0.06	0.5	0.8	14	0.3	0.08	0.2	0.5
1986	*	*	*	*	*	*	*	*	*	*
1987	10	1.3	0.44	0.6	2.6	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Hodgkin Lymphoma										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	719	0.4	0.01	0.4	0.4	619	0.4	0.02	0.4	0.4
1986	56	0.9	0.12	0.7	1.2	51	1.0	0.14	0.7	1.3
1987	34	0.6	0.10	0.4	0.8	32	0.6	0.12	0.4	0.9
1988	33	0.5	0.09	0.4	0.8	27	0.5	0.10	0.3	0.8
1989	30	0.5	0.09	0.3	0.7	25	0.5	0.09	0.3	0.7
1990	25	0.4	0.08	0.2	0.6	21	0.4	0.08	0.2	0.5
1991	32	0.5	0.09	0.3	0.7	26	0.5	0.09	0.3	0.7
1992	31	0.5	0.09	0.3	0.7	27	0.5	0.10	0.3	0.7
1993	40	0.6	0.10	0.4	0.8	37	0.7	0.11	0.5	0.9
1994	31	0.5	0.09	0.3	0.7	25	0.4	0.09	0.3	0.7
1995	23	0.4	0.07	0.2	0.5	18	0.3	0.07	0.2	0.5
1996	25	0.4	0.08	0.3	0.6	20	0.4	0.08	0.2	0.6
1997	30	0.4	0.08	0.3	0.6	29	0.5	0.09	0.3	0.7
1998	18	0.3	0.06	0.2	0.4	16	0.3	0.07	0.2	0.4
1999	21	0.3	0.07	0.2	0.5	18	0.3	0.08	0.2	0.5
2000	27	0.4	0.08	0.3	0.6	24	0.4	0.09	0.3	0.7
2001	23	0.3	0.07	0.2	0.5	18	0.3	0.07	0.2	0.5
2002	27	0.4	0.08	0.3	0.6	25	0.5	0.09	0.3	0.7
2003	24	0.4	0.07	0.2	0.5	22	0.4	0.08	0.3	0.6
2004	19	0.3	0.07	0.2	0.4	14	0.3	0.07	0.1	0.4
2005	26	0.4	0.08	0.3	0.6	23	0.4	0.09	0.3	0.6
2006	23	0.3	0.07	0.2	0.5	19	0.3	0.08	0.2	0.5
2007	27	0.4	0.07	0.3	0.6	24	0.4	0.08	0.3	0.6
2008	21	0.3	0.07	0.2	0.5	17	0.3	0.07	0.2	0.5
2009	24	0.3	0.07	0.2	0.5	20	0.3	0.07	0.2	0.5
2010	17	0.2	0.06	0.1	0.4	15	0.3	0.07	0.1	0.4
2011	17	0.2	0.06	0.1	0.4	12	0.2	0.06	0.1	0.4
2012	15	0.2	0.05	0.1	0.3	14	0.2	0.06	0.1	0.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Hodgkin Lymphoma

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	94	0.4	0.04	0.3	0.5	*	*	*	*	*
1986	*	*	*	*	*	*	*	*	*	*
1987	*	*	*	*	*	*	*	*	*	*
1988	*	*	*	*	*	*	*	*	*	*
1989	*	*	*	*	*	*	*	*	*	*
1990	*	*	*	*	*	*	*	*	*	*
1991	*	*	*	*	*	*	*	*	*	*
1992	*	*	*	*	*	*	*	*	*	*
1993	*	*	*	*	*	*	*	*	*	*
1994	*	*	*	*	*	*	*	*	*	*
1995	*	*	*	*	*	*	*	*	*	*
1996	*	*	*	*	*	*	*	*	*	*
1997	*	*	*	*	*	*	*	*	*	*
1998	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*
2003	*	*	*	*	*	*	*	*	*	*
2004	*	*	*	*	*	*	*	*	*	*
2005	*	*	*	*	*	*	*	*	*	*
2006	*	*	*	*	*	*	*	*	*	*
2007	*	*	*	*	*	*	*	*	*	*
2008	*	*	*	*	*	*	*	*	*	*
2009	*	*	*	*	*	*	*	*	*	*
2010	*	*	*	*	*	*	*	*	*	*
2011	*	*	*	*	*	*	*	*	*	*
2012	*	*	*	*	*	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Non-Hodgkin Lymphoma

Both Sexes

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	24,894	7.7	0.05	7.6	7.8	22,681	8.1	0.05	8.0	8.2
1986	735	7.0	0.26	6.5	7.5	680	7.3	0.28	6.7	7.8
1987	746	7.1	0.26	6.6	7.6	682	7.3	0.28	6.7	7.9
1988	820	7.7	0.27	7.2	8.2	755	8.0	0.29	7.4	8.6
1989	867	8.0	0.28	7.5	8.6	800	8.4	0.30	7.8	9.0
1990	864	8.0	0.27	7.5	8.6	792	8.3	0.30	7.7	8.9
1991	930	8.4	0.28	7.9	9.0	859	8.9	0.30	8.3	9.5
1992	965	8.6	0.28	8.1	9.2	889	9.1	0.31	8.5	9.7
1993	976	8.7	0.28	8.1	9.2	889	9.0	0.30	8.4	9.6
1994	1,024	9.0	0.28	8.5	9.6	925	9.3	0.31	8.7	9.9
1995	1,080	9.4	0.29	8.8	10.0	982	9.8	0.31	9.1	10.4
1996	1,058	9.1	0.28	8.6	9.7	958	9.5	0.31	8.9	10.1
1997	1,039	8.9	0.28	8.4	9.4	948	9.3	0.30	8.7	9.9
1998	1,019	8.6	0.27	8.1	9.2	921	8.9	0.29	8.4	9.5
1999	1,061	8.9	0.27	8.4	9.5	975	9.4	0.30	8.8	10.0
2000	1,004	8.4	0.26	7.9	8.9	924	8.9	0.29	8.3	9.5
2001	987	8.1	0.26	7.6	8.7	902	8.6	0.29	8.0	9.2
2002	943	7.7	0.25	7.2	8.2	854	8.1	0.28	7.5	8.6
2003	927	7.5	0.25	7.0	8.0	849	7.9	0.27	7.4	8.5
2004	935	7.5	0.25	7.1	8.0	859	8.0	0.28	7.5	8.6
2005	950	7.6	0.25	7.1	8.1	864	7.9	0.27	7.4	8.5
2006	885	7.0	0.23	6.5	7.4	808	7.4	0.26	6.9	7.9
2007	820	6.4	0.22	5.9	6.8	739	6.7	0.25	6.2	7.2
2008	854	6.6	0.23	6.1	7.0	778	7.0	0.25	6.5	7.5
2009	903	6.8	0.23	6.4	7.3	795	7.0	0.25	6.6	7.6
2010	847	6.3	0.22	5.9	6.8	762	6.7	0.25	6.2	7.2
2011	845	6.2	0.22	5.8	6.6	754	6.5	0.24	6.0	7.0
2012	810	5.9	0.21	5.5	6.3	738	6.4	0.24	5.9	6.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Non-Hodgkin Lymphoma

Both Sexes

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,940	5.2	0.12	4.9	5.4	273	3.6	0.24	3.2	4.1
1986	51	4.7	0.67	3.5	6.2	*	*	*	*	*
1987	60	5.2	0.70	4.0	6.8	*	*	*	*	*
1988	55	4.5	0.62	3.3	5.8	10	8.3	3.03	3.5	15.9
1989	59	4.7	0.64	3.6	6.2	*	*	*	*	*
1990	69	6.0	0.75	4.6	7.6	*	*	*	*	*
1991	67	5.3	0.68	4.1	6.8	*	*	*	*	*
1992	74	6.0	0.72	4.6	7.5	*	*	*	*	*
1993	82	6.4	0.73	5.0	8.0	*	*	*	*	*
1994	91	7.1	0.76	5.6	8.7	*	*	*	*	*
1995	88	6.9	0.75	5.5	8.5	10	5.5	1.94	2.4	10.3
1996	94	7.1	0.75	5.7	8.7	*	*	*	*	*
1997	78	5.7	0.66	4.5	7.1	13	5.2	1.57	2.6	9.1
1998	90	6.7	0.72	5.3	8.2	*	*	*	*	*
1999	80	5.8	0.67	4.6	7.3	*	*	*	*	*
2000	68	4.8	0.60	3.7	6.2	12	3.5	1.15	1.6	6.4
2001	75	5.2	0.61	4.1	6.5	10	2.4	0.82	1.1	4.6
2002	76	5.2	0.61	4.1	6.6	13	3.9	1.16	2.0	6.8
2003	60	4.1	0.54	3.1	5.3	18	6.8	1.71	3.8	10.8
2004	60	4.1	0.54	3.1	5.3	16	4.2	1.12	2.3	6.9
2005	69	4.6	0.57	3.6	5.9	17	5.4	1.38	3.0	8.6
2006	67	4.3	0.54	3.3	5.5	10	2.6	0.87	1.2	4.8
2007	65	4.3	0.55	3.3	5.5	16	3.9	1.04	2.1	6.4
2008	64	4.3	0.54	3.3	5.4	12	2.8	0.85	1.4	4.9
2009	88	5.3	0.58	4.3	6.6	20	4.6	1.10	2.7	7.2
2010	72	4.3	0.52	3.4	5.5	13	2.4	0.71	1.2	4.2
2011	78	4.6	0.53	3.6	5.7	13	2.9	0.83	1.5	4.9
2012	60	3.6	0.47	2.7	4.6	12	2.5	0.77	1.3	4.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Non-Hodgkin Lymphoma

Males

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	12,990	9.9	0.09	9.7	10.0	11,763	10.3	0.10	10.1	10.5
1986	375	8.5	0.46	7.6	9.5	339	8.7	0.49	7.8	9.7
1987	375	8.5	0.46	7.7	9.5	335	8.7	0.49	7.7	9.7
1988	417	9.6	0.49	8.7	10.6	382	10.1	0.53	9.0	11.2
1989	448	10.0	0.49	9.1	11.0	407	10.3	0.53	9.3	11.4
1990	481	10.9	0.52	9.9	12.0	443	11.4	0.56	10.3	12.5
1991	480	10.5	0.50	9.6	11.5	439	11.0	0.54	10.0	12.1
1992	514	11.3	0.52	10.3	12.3	466	11.7	0.56	10.6	12.8
1993	488	10.6	0.49	9.6	11.6	442	11.0	0.54	9.9	12.1
1994	534	11.6	0.51	10.6	12.6	477	11.9	0.56	10.8	13.0
1995	550	11.7	0.51	10.7	12.8	499	12.2	0.56	11.1	13.3
1996	546	11.4	0.50	10.5	12.5	484	11.6	0.54	10.6	12.7
1997	530	10.9	0.48	10.0	11.9	472	11.2	0.52	10.2	12.3
1998	541	11.4	0.50	10.4	12.4	486	11.8	0.54	10.7	12.9
1999	567	11.7	0.50	10.8	12.7	518	12.3	0.55	11.3	13.4
2000	536	10.8	0.47	9.9	11.8	491	11.5	0.52	10.5	12.5
2001	516	10.4	0.46	9.5	11.3	472	11.0	0.51	10.0	12.0
2002	475	9.6	0.45	8.8	10.5	436	10.2	0.49	9.3	11.2
2003	466	9.3	0.43	8.4	10.1	422	9.7	0.48	8.8	10.7
2004	478	9.5	0.44	8.6	10.4	439	10.1	0.49	9.2	11.1
2005	493	9.8	0.45	8.9	10.7	449	10.3	0.49	9.3	11.3
2006	480	9.2	0.42	8.3	10.0	437	9.7	0.47	8.8	10.7
2007	435	8.4	0.41	7.6	9.2	396	8.8	0.45	8.0	9.7
2008	438	8.3	0.40	7.5	9.1	400	8.8	0.44	8.0	9.7
2009	455	8.4	0.40	7.6	9.2	401	8.6	0.43	7.8	9.5
2010	464	8.3	0.39	7.6	9.2	414	8.8	0.44	7.9	9.7
2011	454	8.2	0.39	7.4	9.0	402	8.5	0.43	7.7	9.4
2012	454	7.9	0.38	7.2	8.7	415	8.5	0.42	7.7	9.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Non-Hodgkin Lymphoma

Males

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,075	6.8	0.22	6.4	7.2	152	4.7	0.43	3.9	5.6
1986	33	7.0	1.25	4.8	9.9	*	*	*	*	*
1987	38	7.8	1.36	5.4	10.9	*	*	*	*	*
1988	30	5.5	1.04	3.7	7.9	*	*	*	*	*
1989	36	6.6	1.17	4.5	9.3	*	*	*	*	*
1990	36	7.5	1.37	5.1	10.6	*	*	*	*	*
1991	38	6.9	1.22	4.7	9.7	*	*	*	*	*
1992	48	9.3	1.47	6.6	12.6	*	*	*	*	*
1993	43	7.6	1.22	5.4	10.3	*	*	*	*	*
1994	52	9.1	1.35	6.7	12.1	*	*	*	*	*
1995	47	8.4	1.32	6.0	11.3	*	*	*	*	*
1996	59	11.4	1.62	8.4	15.0	*	*	*	*	*
1997	46	7.5	1.16	5.4	10.1	12	10.8	3.51	5.1	19.7
1998	49	8.4	1.28	6.1	11.3	*	*	*	*	*
1999	45	7.8	1.24	5.6	10.6	*	*	*	*	*
2000	36	5.9	1.07	4.0	8.4	*	*	*	*	*
2001	38	6.1	1.04	4.2	8.4	*	*	*	*	*
2002	31	5.2	0.99	3.4	7.5	*	*	*	*	*
2003	34	5.7	1.07	3.8	8.1	10	8.3	2.91	3.6	15.5
2004	32	5.3	1.00	3.5	7.5	*	*	*	*	*
2005	36	6.0	1.05	4.1	8.3	*	*	*	*	*
2006	36	5.5	0.98	3.7	7.7	*	*	*	*	*
2007	34	5.9	1.09	3.9	8.4	*	*	*	*	*
2008	32	5.0	0.96	3.3	7.2	*	*	*	*	*
2009	42	6.5	1.09	4.6	9.0	12	7.3	2.30	3.5	13.0
2010	44	6.1	0.96	4.4	8.3	*	*	*	*	*
2011	44	6.2	0.99	4.5	8.5	*	*	*	*	*
2012	36	5.3	0.92	3.6	7.4	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Non-Hodgkin Lymphoma

Females

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	11,904	6.2	0.06	6.1	6.3	10,918	6.6	0.06	6.4	6.7
1986	360	5.7	0.31	5.1	6.4	341	6.1	0.34	5.5	6.8
1987	371	5.9	0.31	5.3	6.5	347	6.2	0.34	5.5	6.9
1988	403	6.4	0.32	5.7	7.0	373	6.6	0.35	6.0	7.3
1989	419	6.6	0.33	6.0	7.3	393	7.0	0.36	6.3	7.7
1990	383	5.9	0.30	5.3	6.5	349	6.0	0.33	5.4	6.7
1991	450	6.8	0.33	6.2	7.5	420	7.2	0.35	6.5	7.9
1992	451	6.8	0.32	6.2	7.4	423	7.2	0.36	6.5	8.0
1993	488	7.2	0.33	6.6	7.9	447	7.5	0.36	6.8	8.2
1994	490	7.3	0.33	6.7	8.0	448	7.6	0.37	6.9	8.4
1995	530	7.7	0.34	7.1	8.4	483	8.0	0.37	7.3	8.8
1996	512	7.4	0.33	6.8	8.1	474	7.9	0.37	7.2	8.6
1997	509	7.3	0.33	6.7	8.0	476	7.8	0.36	7.1	8.5
1998	478	6.8	0.31	6.2	7.4	435	7.1	0.34	6.4	7.8
1999	494	6.9	0.31	6.3	7.6	457	7.3	0.35	6.7	8.1
2000	468	6.6	0.31	6.0	7.2	433	7.0	0.34	6.3	7.7
2001	471	6.5	0.30	5.9	7.2	430	6.9	0.34	6.2	7.6
2002	468	6.4	0.30	5.8	7.0	418	6.6	0.33	6.0	7.3
2003	461	6.1	0.29	5.6	6.7	427	6.6	0.32	6.0	7.2
2004	457	6.2	0.30	5.7	6.9	420	6.7	0.33	6.0	7.3
2005	457	6.1	0.29	5.5	6.7	415	6.4	0.32	5.7	7.0
2006	405	5.4	0.27	4.9	5.9	371	5.7	0.30	5.2	6.4
2007	385	5.0	0.26	4.5	5.5	343	5.2	0.29	4.6	5.8
2008	416	5.3	0.27	4.8	5.9	378	5.7	0.30	5.1	6.3
2009	448	5.8	0.28	5.2	6.3	394	5.9	0.31	5.3	6.5
2010	383	4.8	0.25	4.3	5.3	348	5.1	0.28	4.6	5.7
2011	391	4.8	0.25	4.4	5.4	352	5.1	0.28	4.6	5.7
2012	356	4.4	0.24	4.0	4.9	323	4.8	0.27	4.2	5.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Non-Hodgkin Lymphoma

Females

Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	865	4.0	0.14	3.7	4.3	121	2.9	0.28	2.4	3.5
1986	18	2.9	0.69	1.7	4.5	*	*	*	*	*
1987	22	3.4	0.74	2.1	5.1	*	*	*	*	*
1988	25	3.6	0.74	2.3	5.4	*	*	*	*	*
1989	23	3.3	0.71	2.1	5.0	*	*	*	*	*
1990	33	4.9	0.87	3.4	6.9	*	*	*	*	*
1991	29	4.1	0.78	2.7	6.0	*	*	*	*	*
1992	26	3.7	0.74	2.4	5.4	*	*	*	*	*
1993	39	5.3	0.87	3.8	7.3	*	*	*	*	*
1994	39	5.4	0.88	3.8	7.4	*	*	*	*	*
1995	41	5.6	0.88	4.0	7.6	*	*	*	*	*
1996	35	4.4	0.76	3.1	6.2	*	*	*	*	*
1997	32	4.2	0.75	2.8	5.9	*	*	*	*	*
1998	41	5.3	0.84	3.8	7.2	*	*	*	*	*
1999	35	4.4	0.75	3.1	6.1	*	*	*	*	*
2000	32	4.0	0.71	2.7	5.6	*	*	*	*	*
2001	37	4.5	0.74	3.1	6.2	*	*	*	*	*
2002	45	5.3	0.80	3.9	7.1	*	*	*	*	*
2003	26	3.1	0.61	2.0	4.5	*	*	*	*	*
2004	28	3.3	0.63	2.2	4.8	*	*	*	*	*
2005	33	3.7	0.65	2.6	5.2	*	*	*	*	*
2006	31	3.5	0.63	2.4	4.9	*	*	*	*	*
2007	31	3.4	0.61	2.3	4.8	11	4.4	1.39	2.1	8.0
2008	32	3.6	0.65	2.5	5.1	*	*	*	*	*
2009	46	4.8	0.71	3.5	6.4	*	*	*	*	*
2010	28	2.9	0.56	1.9	4.3	*	*	*	*	*
2011	34	3.4	0.60	2.4	4.8	*	*	*	*	*
2012	24	2.5	0.51	1.6	3.7	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Myeloma										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	12,127	3.8	0.03	3.7	3.8	9,487	3.4	0.03	3.3	3.4
1986	407	3.9	0.20	3.5	4.3	324	3.5	0.20	3.1	3.9
1987	385	3.6	0.19	3.3	4.0	298	3.1	0.18	2.8	3.5
1988	381	3.6	0.18	3.2	3.9	316	3.3	0.19	3.0	3.7
1989	422	3.9	0.19	3.5	4.3	329	3.4	0.19	3.1	3.8
1990	384	3.6	0.18	3.2	4.0	309	3.2	0.19	2.9	3.6
1991	415	3.8	0.19	3.5	4.2	332	3.4	0.19	3.1	3.8
1992	447	4.0	0.19	3.7	4.4	355	3.6	0.19	3.2	4.0
1993	457	4.1	0.19	3.7	4.4	347	3.5	0.19	3.1	3.9
1994	439	3.9	0.18	3.5	4.2	353	3.5	0.19	3.2	3.9
1995	492	4.3	0.19	3.9	4.7	378	3.7	0.19	3.4	4.1
1996	442	3.8	0.18	3.5	4.2	341	3.4	0.18	3.0	3.7
1997	448	3.8	0.18	3.5	4.2	352	3.5	0.18	3.1	3.8
1998	485	4.1	0.19	3.8	4.5	359	3.5	0.18	3.1	3.9
1999	483	4.1	0.18	3.7	4.4	379	3.6	0.19	3.3	4.0
2000	501	4.2	0.19	3.8	4.6	382	3.7	0.19	3.3	4.1
2001	469	3.9	0.18	3.6	4.3	367	3.5	0.18	3.2	3.9
2002	473	3.9	0.18	3.5	4.2	378	3.6	0.18	3.2	3.9
2003	482	3.9	0.18	3.6	4.3	378	3.5	0.18	3.2	3.9
2004	442	3.6	0.17	3.2	3.9	345	3.2	0.18	2.9	3.6
2005	481	3.8	0.18	3.5	4.2	375	3.5	0.18	3.1	3.8
2006	462	3.7	0.17	3.3	4.0	358	3.3	0.18	3.0	3.7
2007	468	3.7	0.17	3.4	4.0	365	3.3	0.18	3.0	3.7
2008	437	3.4	0.16	3.1	3.7	344	3.1	0.17	2.8	3.4
2009	443	3.4	0.16	3.1	3.7	348	3.1	0.17	2.8	3.5
2010	446	3.3	0.16	3.0	3.6	341	2.9	0.16	2.6	3.3
2011	456	3.3	0.16	3.0	3.7	363	3.1	0.17	2.8	3.5
2012	480	3.5	0.16	3.2	3.8	371	3.2	0.17	2.9	3.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Myeloma										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,528	7.2	0.15	6.9	7.5	112	1.4	0.14	1.1	1.7
1986	82	7.5	0.86	6.0	9.4	*	*	*	*	*
1987	87	8.4	0.94	6.7	10.4	*	*	*	*	*
1988	64	5.9	0.76	4.5	7.6	*	*	*	*	*
1989	90	8.3	0.91	6.7	10.3	*	*	*	*	*
1990	75	7.1	0.84	5.5	8.9	*	*	*	*	*
1991	82	7.7	0.88	6.1	9.6	*	*	*	*	*
1992	90	7.9	0.85	6.3	9.7	*	*	*	*	*
1993	103	8.8	0.89	7.2	10.8	*	*	*	*	*
1994	82	6.9	0.78	5.4	8.5	*	*	*	*	*
1995	113	9.5	0.91	7.8	11.4	*	*	*	*	*
1996	98	8.1	0.84	6.5	9.9	*	*	*	*	*
1997	85	6.7	0.74	5.3	8.3	11	4.5	1.45	2.1	8.1
1998	123	9.9	0.91	8.2	11.8	*	*	*	*	*
1999	99	7.8	0.79	6.3	9.5	*	*	*	*	*
2000	115	8.8	0.84	7.2	10.6	*	*	*	*	*
2001	98	7.4	0.75	6.0	9.0	*	*	*	*	*
2002	89	6.5	0.71	5.2	8.1	*	*	*	*	*
2003	101	7.7	0.77	6.2	9.3	*	*	*	*	*
2004	91	6.6	0.70	5.3	8.1	*	*	*	*	*
2005	101	6.9	0.70	5.6	8.4	*	*	*	*	*
2006	99	6.8	0.69	5.5	8.3	*	*	*	*	*
2007	97	6.7	0.69	5.4	8.1	*	*	*	*	*
2008	86	5.7	0.63	4.5	7.1	*	*	*	*	*
2009	89	5.9	0.64	4.7	7.3	*	*	*	*	*
2010	98	6.4	0.66	5.2	7.8	*	*	*	*	*
2011	88	5.5	0.60	4.4	6.8	*	*	*	*	*
2012	103	6.4	0.64	5.2	7.8	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Myeloma										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	6,035	4.6	0.06	4.5	4.7	4,780	4.2	0.06	4.1	4.3
1986	208	4.9	0.35	4.2	5.6	164	4.3	0.35	3.7	5.1
1987	198	4.7	0.35	4.0	5.4	156	4.2	0.35	3.5	4.9
1988	192	4.6	0.34	3.9	5.3	164	4.4	0.36	3.7	5.2
1989	199	4.6	0.34	4.0	5.4	164	4.4	0.35	3.7	5.1
1990	182	4.2	0.32	3.6	4.9	150	3.9	0.33	3.3	4.7
1991	205	4.9	0.35	4.2	5.6	164	4.4	0.35	3.7	5.1
1992	209	4.6	0.33	4.0	5.3	164	4.2	0.33	3.5	4.9
1993	205	4.3	0.31	3.7	4.9	149	3.6	0.30	3.0	4.2
1994	217	4.8	0.33	4.2	5.5	172	4.3	0.34	3.7	5.0
1995	250	5.5	0.35	4.8	6.2	194	4.9	0.36	4.2	5.6
1996	225	4.9	0.33	4.2	5.6	172	4.3	0.33	3.7	5.0
1997	217	4.5	0.31	3.9	5.1	171	4.1	0.32	3.5	4.8
1998	230	4.9	0.33	4.3	5.6	167	4.1	0.32	3.5	4.7
1999	222	4.6	0.32	4.0	5.3	180	4.3	0.33	3.7	5.0
2000	258	5.4	0.34	4.7	6.1	207	5.0	0.35	4.3	5.7
2001	222	4.5	0.31	4.0	5.2	178	4.2	0.32	3.6	4.9
2002	245	5.0	0.32	4.4	5.6	194	4.5	0.33	3.9	5.2
2003	216	4.3	0.30	3.8	4.9	173	4.0	0.31	3.4	4.6
2004	222	4.4	0.30	3.8	5.0	174	4.0	0.30	3.4	4.6
2005	237	4.6	0.30	4.0	5.2	189	4.3	0.31	3.7	4.9
2006	235	4.5	0.30	4.0	5.1	189	4.2	0.31	3.7	4.9
2007	232	4.5	0.30	3.9	5.1	187	4.2	0.31	3.6	4.8
2008	248	4.7	0.30	4.1	5.3	195	4.3	0.31	3.7	4.9
2009	241	4.4	0.29	3.9	5.0	196	4.2	0.30	3.6	4.8
2010	245	4.4	0.29	3.9	5.0	187	4.0	0.29	3.4	4.6
2011	227	4.0	0.27	3.5	4.5	179	3.7	0.28	3.2	4.3
2012	248	4.3	0.28	3.8	4.9	201	4.1	0.29	3.5	4.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Myeloma										
Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,189	8.6	0.26	8.1	9.1	66	1.8	0.25	1.4	2.4
1986	43	9.8	1.61	7.0	13.4	*	*	*	*	*
1987	42	10.3	1.72	7.2	14.2	*	*	*	*	*
1988	28	6.9	1.38	4.4	10.0	*	*	*	*	*
1989	34	7.2	1.30	4.9	10.2	*	*	*	*	*
1990	32	7.2	1.36	4.7	10.2	*	*	*	*	*
1991	40	9.9	1.67	6.9	13.6	*	*	*	*	*
1992	44	9.1	1.44	6.5	12.3	*	*	*	*	*
1993	51	10.2	1.50	7.5	13.6	*	*	*	*	*
1994	42	9.2	1.54	6.4	12.6	*	*	*	*	*
1995	55	11.4	1.64	8.4	15.0	*	*	*	*	*
1996	50	10.7	1.61	7.7	14.2	*	*	*	*	*
1997	40	6.9	1.13	4.9	9.5	*	*	*	*	*
1998	61	13.0	1.79	9.8	16.9	*	*	*	*	*
1999	38	7.4	1.26	5.2	10.3	*	*	*	*	*
2000	49	9.5	1.43	6.9	12.6	*	*	*	*	*
2001	42	7.9	1.27	5.6	10.7	*	*	*	*	*
2002	47	9.2	1.42	6.6	12.3	*	*	*	*	*
2003	43	8.3	1.36	5.9	11.4	*	*	*	*	*
2004	44	8.5	1.35	6.1	11.5	*	*	*	*	*
2005	45	7.7	1.23	5.5	10.5	*	*	*	*	*
2006	44	7.8	1.25	5.6	10.6	*	*	*	*	*
2007	43	7.5	1.18	5.4	10.1	*	*	*	*	*
2008	49	9.0	1.34	6.5	11.9	*	*	*	*	*
2009	41	7.1	1.15	5.0	9.7	*	*	*	*	*
2010	52	8.6	1.25	6.3	11.3	*	*	*	*	*
2011	45	7.1	1.12	5.1	9.6	*	*	*	*	*
2012	45	7.1	1.10	5.1	9.6	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Myeloma										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	6,092	3.2	0.04	3.1	3.3	4,707	2.8	0.04	2.7	2.9
1986	199	3.2	0.23	2.8	3.7	160	2.9	0.23	2.5	3.4
1987	187	2.9	0.22	2.5	3.4	142	2.5	0.21	2.1	2.9
1988	189	3.0	0.22	2.6	3.5	152	2.7	0.22	2.3	3.2
1989	223	3.4	0.23	3.0	3.9	165	2.8	0.22	2.4	3.3
1990	202	3.1	0.22	2.7	3.6	159	2.7	0.22	2.3	3.2
1991	210	3.2	0.22	2.8	3.7	168	2.9	0.23	2.5	3.4
1992	238	3.5	0.23	3.1	4.0	191	3.2	0.23	2.7	3.7
1993	252	3.7	0.24	3.3	4.2	198	3.3	0.24	2.9	3.8
1994	222	3.3	0.22	2.9	3.7	181	3.0	0.22	2.6	3.5
1995	242	3.5	0.23	3.1	4.0	184	3.1	0.23	2.6	3.6
1996	217	3.1	0.21	2.7	3.6	169	2.8	0.21	2.4	3.2
1997	231	3.3	0.22	2.9	3.8	181	2.9	0.22	2.5	3.4
1998	255	3.7	0.23	3.2	4.1	192	3.2	0.23	2.7	3.6
1999	261	3.7	0.23	3.3	4.2	199	3.2	0.23	2.8	3.7
2000	243	3.4	0.22	3.0	3.9	175	2.8	0.22	2.4	3.3
2001	247	3.5	0.22	3.1	3.9	189	3.1	0.22	2.6	3.5
2002	228	3.1	0.21	2.7	3.6	184	2.9	0.21	2.5	3.3
2003	266	3.6	0.22	3.2	4.1	205	3.2	0.23	2.8	3.7
2004	220	3.0	0.20	2.6	3.4	171	2.7	0.21	2.3	3.1
2005	244	3.3	0.21	2.9	3.8	186	2.9	0.22	2.5	3.4
2006	227	3.1	0.21	2.7	3.5	169	2.7	0.21	2.3	3.1
2007	236	3.2	0.21	2.8	3.6	178	2.8	0.21	2.4	3.2
2008	189	2.5	0.18	2.1	2.9	149	2.3	0.19	1.9	2.7
2009	202	2.6	0.19	2.3	3.0	152	2.3	0.19	2.0	2.8
2010	201	2.5	0.18	2.1	2.9	154	2.2	0.18	1.8	2.5
2011	229	2.9	0.19	2.5	3.3	184	2.7	0.20	2.3	3.1
2012	232	2.9	0.20	2.6	3.4	170	2.6	0.20	2.2	3.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Myeloma										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,339	6.4	0.18	6.0	6.7	46	1.0	0.16	0.7	1.4
1986	39	6.1	0.99	4.3	8.4	*	*	*	*	*
1987	45	7.2	1.10	5.2	9.7	*	*	*	*	*
1988	36	5.5	0.93	3.8	7.6	*	*	*	*	*
1989	56	8.8	1.20	6.6	11.5	*	*	*	*	*
1990	43	7.0	1.07	5.0	9.4	*	*	*	*	*
1991	42	6.5	1.01	4.7	8.8	*	*	*	*	*
1992	46	6.9	1.03	5.0	9.2	*	*	*	*	*
1993	52	7.6	1.07	5.7	10.0	*	*	*	*	*
1994	40	5.6	0.90	4.0	7.7	*	*	*	*	*
1995	58	8.1	1.07	6.1	10.5	*	*	*	*	*
1996	48	6.5	0.95	4.8	8.6	*	*	*	*	*
1997	45	6.2	0.92	4.5	8.2	*	*	*	*	*
1998	62	8.1	1.04	6.2	10.4	*	*	*	*	*
1999	61	7.9	1.01	6.0	10.1	*	*	*	*	*
2000	66	8.4	1.03	6.4	10.6	*	*	*	*	*
2001	56	7.0	0.93	5.2	9.0	*	*	*	*	*
2002	42	5.1	0.79	3.7	6.8	*	*	*	*	*
2003	58	7.3	0.96	5.5	9.4	*	*	*	*	*
2004	47	5.5	0.81	4.0	7.3	*	*	*	*	*
2005	56	6.4	0.86	4.8	8.3	*	*	*	*	*
2006	55	6.2	0.85	4.7	8.1	*	*	*	*	*
2007	54	6.0	0.83	4.5	7.9	*	*	*	*	*
2008	37	3.9	0.65	2.7	5.4	*	*	*	*	*
2009	48	5.2	0.75	3.8	6.9	*	*	*	*	*
2010	46	5.1	0.75	3.7	6.7	*	*	*	*	*
2011	43	4.5	0.69	3.2	6.0	*	*	*	*	*
2012	58	5.8	0.78	4.4	7.6	*	*	*	*	*

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Leukemia										
Both Sexes										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	25,665	8.0	0.05	7.9	8.1	22,789	8.2	0.05	8.1	8.3
1986	872	8.3	0.29	7.8	8.9	777	8.4	0.30	7.8	9.0
1987	810	7.7	0.27	7.2	8.2	731	7.9	0.29	7.3	8.4
1988	849	8.0	0.28	7.5	8.6	750	8.0	0.30	7.4	8.6
1989	911	8.5	0.28	8.0	9.1	825	8.8	0.31	8.2	9.4
1990	929	8.6	0.28	8.0	9.1	830	8.7	0.30	8.1	9.3
1991	948	8.6	0.28	8.1	9.2	854	8.8	0.30	8.3	9.5
1992	945	8.5	0.28	8.0	9.1	849	8.7	0.30	8.1	9.3
1993	970	8.6	0.28	8.0	9.1	872	8.8	0.30	8.2	9.4
1994	940	8.2	0.27	7.7	8.8	836	8.4	0.29	7.8	9.0
1995	967	8.4	0.27	7.9	9.0	862	8.6	0.29	8.0	9.2
1996	933	8.0	0.26	7.5	8.6	828	8.2	0.28	7.6	8.8
1997	956	8.2	0.26	7.7	8.7	846	8.3	0.29	7.7	8.9
1998	930	7.8	0.26	7.3	8.4	802	7.8	0.28	7.3	8.4
1999	979	8.2	0.26	7.7	8.7	879	8.5	0.29	7.9	9.0
2000	989	8.2	0.26	7.7	8.7	889	8.5	0.29	8.0	9.1
2001	1,013	8.3	0.26	7.8	8.9	893	8.5	0.29	8.0	9.1
2002	1,003	8.2	0.26	7.7	8.7	892	8.5	0.28	7.9	9.1
2003	993	8.0	0.26	7.5	8.6	875	8.2	0.28	7.6	8.7
2004	945	7.6	0.25	7.1	8.1	849	8.0	0.27	7.4	8.5
2005	978	7.8	0.25	7.3	8.3	863	8.0	0.27	7.5	8.6
2006	953	7.5	0.24	7.0	8.0	859	7.9	0.27	7.4	8.4
2007	908	7.1	0.24	6.6	7.6	813	7.4	0.26	6.9	8.0
2008	960	7.4	0.24	6.9	7.9	852	7.7	0.27	7.2	8.3
2009	1,049	8.0	0.25	7.5	8.5	919	8.3	0.28	7.8	8.9
2010	979	7.3	0.24	6.9	7.8	853	7.6	0.26	7.1	8.1
2011	914	6.8	0.23	6.3	7.2	790	6.9	0.25	6.5	7.4
2012	1,042	7.6	0.24	7.1	8.1	901	7.8	0.26	7.3	8.3

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Leukemia										
Both Sexes										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	2,561	7.0	0.14	6.7	7.2	315	3.7	0.24	3.3	4.2
1986	90	8.4	0.94	6.7	10.5	*	*	*	*	*
1987	78	7.2	0.85	5.6	9.0	*	*	*	*	*
1988	94	8.6	0.93	6.9	10.6	*	*	*	*	*
1989	76	6.2	0.74	4.8	7.8	10	4.0	1.35	1.8	7.9
1990	93	7.8	0.85	6.2	9.6	*	*	*	*	*
1991	85	7.2	0.82	5.7	9.0	*	*	*	*	*
1992	86	7.0	0.79	5.6	8.7	10	6.3	2.26	2.7	12.0
1993	93	7.3	0.78	5.8	9.0	*	*	*	*	*
1994	95	7.6	0.81	6.1	9.3	*	*	*	*	*
1995	95	7.4	0.79	6.0	9.1	10	3.6	1.32	1.5	7.1
1996	94	7.5	0.80	6.0	9.2	11	4.0	1.46	1.7	7.8
1997	105	8.0	0.81	6.5	9.7	*	*	*	*	*
1998	115	8.4	0.81	6.9	10.2	13	4.9	1.50	2.4	8.6
1999	93	6.7	0.71	5.3	8.2	*	*	*	*	*
2000	89	6.1	0.66	4.8	7.5	11	3.1	0.99	1.4	5.7
2001	103	7.3	0.74	5.9	8.8	17	4.6	1.26	2.5	7.7
2002	97	7.1	0.74	5.7	8.7	14	3.5	1.03	1.8	6.1
2003	108	7.6	0.75	6.2	9.2	10	2.3	0.77	1.0	4.3
2004	85	6.1	0.68	4.9	7.6	11	2.6	0.84	1.2	4.7
2005	99	6.7	0.69	5.4	8.2	16	4.8	1.32	2.6	8.0
2006	89	6.0	0.66	4.8	7.4	*	*	*	*	*
2007	77	5.1	0.59	4.0	6.3	18	3.8	1.00	2.1	6.3
2008	94	5.9	0.63	4.8	7.3	14	3.0	0.88	1.6	5.2
2009	115	7.5	0.72	6.2	9.1	15	2.5	0.70	1.4	4.3
2010	103	6.4	0.65	5.2	7.8	23	4.4	0.98	2.7	6.7
2011	97	6.0	0.62	4.8	7.3	27	5.7	1.15	3.6	8.3
2012	113	6.8	0.66	5.6	8.2	28	5.8	1.16	3.8	8.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Leukemia										
Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	13,957	10.7	0.09	10.5	10.9	12,430	11.0	0.10	10.8	11.2
1986	475	11.3	0.55	10.3	12.5	425	11.5	0.58	10.3	12.7
1987	416	9.8	0.50	8.8	10.8	377	10.1	0.54	9.1	11.2
1988	444	10.5	0.52	9.5	11.5	388	10.4	0.55	9.4	11.5
1989	504	11.4	0.53	10.4	12.5	454	11.8	0.57	10.7	13.0
1990	486	11.1	0.53	10.1	12.2	439	11.4	0.57	10.3	12.6
1991	472	10.5	0.51	9.6	11.6	430	10.9	0.54	9.8	12.0
1992	528	11.9	0.54	10.9	13.0	482	12.4	0.58	11.3	13.6
1993	543	12.1	0.54	11.1	13.2	480	12.3	0.58	11.2	13.4
1994	524	11.4	0.51	10.4	12.4	466	11.6	0.55	10.5	12.7
1995	527	11.6	0.52	10.6	12.7	468	11.8	0.56	10.7	13.0
1996	500	11.1	0.51	10.1	12.1	446	11.3	0.55	10.2	12.4
1997	524	11.2	0.50	10.2	12.2	466	11.4	0.54	10.4	12.5
1998	478	10.0	0.46	9.1	10.9	424	10.2	0.50	9.2	11.2
1999	513	10.8	0.49	9.9	11.8	461	11.2	0.53	10.1	12.2
2000	523	10.7	0.47	9.7	11.6	471	11.1	0.52	10.1	12.2
2001	549	11.2	0.49	10.3	12.2	478	11.3	0.52	10.3	12.4
2002	552	11.3	0.49	10.3	12.3	495	11.7	0.53	10.7	12.8
2003	566	11.5	0.49	10.6	12.5	499	11.7	0.53	10.7	12.8
2004	534	10.6	0.47	9.8	11.6	483	11.2	0.51	10.2	12.2
2005	537	10.6	0.46	9.7	11.5	476	10.9	0.50	9.9	11.9
2006	518	10.0	0.45	9.1	10.9	465	10.4	0.49	9.5	11.4
2007	493	9.5	0.43	8.7	10.4	447	10.0	0.48	9.1	11.0
2008	542	10.1	0.44	9.3	11.0	484	10.6	0.49	9.6	11.6
2009	588	10.8	0.45	10.0	11.8	514	11.1	0.50	10.2	12.1
2010	542	9.9	0.43	9.1	10.8	476	10.2	0.47	9.3	11.2
2011	517	9.3	0.42	8.5	10.1	442	9.3	0.45	8.5	10.3
2012	562	9.9	0.42	9.1	10.8	494	10.2	0.47	9.3	11.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Leukemia										
Males										
Year	Black					Asian or Other Races				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,348	9.1	0.27	8.6	9.7	179	4.8	0.43	4.0	5.7
1986	49	11.7	1.87	8.3	15.9	*	*	*	*	*
1987	39	8.3	1.44	5.8	11.6	*	*	*	*	*
1988	53	11.5	1.74	8.4	15.4	*	*	*	*	*
1989	42	8.1	1.34	5.7	11.1	*	*	*	*	*
1990	44	8.7	1.47	6.0	12.0	*	*	*	*	*
1991	38	8.3	1.53	5.6	11.8	*	*	*	*	*
1992	40	8.0	1.40	5.5	11.2	*	*	*	*	*
1993	60	12.1	1.68	9.0	15.8	*	*	*	*	*
1994	53	10.8	1.63	7.8	14.4	*	*	*	*	*
1995	53	10.6	1.56	7.8	14.1	*	*	*	*	*
1996	49	9.8	1.55	7.0	13.3	*	*	*	*	*
1997	56	11.0	1.58	8.1	14.4	*	*	*	*	*
1998	47	8.0	1.23	5.7	10.7	*	*	*	*	*
1999	47	8.6	1.35	6.1	11.6	*	*	*	*	*
2000	44	7.0	1.11	5.0	9.5	*	*	*	*	*
2001	57	10.2	1.48	7.5	13.4	14	7.2	2.09	3.7	12.7
2002	47	8.9	1.41	6.3	12.0	10	5.7	2.00	2.5	11.0
2003	60	10.4	1.42	7.8	13.5	*	*	*	*	*
2004	44	8.0	1.31	5.6	10.9	*	*	*	*	*
2005	53	8.6	1.25	6.3	11.3	*	*	*	*	*
2006	51	8.3	1.26	6.0	11.1	*	*	*	*	*
2007	37	6.2	1.09	4.3	8.7	*	*	*	*	*
2008	49	7.9	1.21	5.7	10.6	*	*	*	*	*
2009	65	10.7	1.42	8.1	13.8	*	*	*	*	*
2010	55	8.4	1.20	6.2	11.0	11	5.4	1.82	2.4	9.9
2011	59	9.4	1.29	7.0	12.2	16	7.2	1.97	3.9	12.0
2012	57	8.8	1.23	6.5	11.5	11	6.1	1.99	2.9	11.0

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Leukemia										
Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	11,708	6.1	0.06	6.0	6.3	10,359	6.3	0.06	6.2	6.4
1986	397	6.4	0.33	5.8	7.1	352	6.4	0.35	5.8	7.1
1987	394	6.4	0.33	5.8	7.1	354	6.5	0.35	5.8	7.2
1988	405	6.5	0.33	5.9	7.1	362	6.6	0.35	5.9	7.3
1989	407	6.5	0.32	5.8	7.1	371	6.6	0.35	6.0	7.4
1990	443	7.0	0.33	6.3	7.6	391	6.9	0.36	6.2	7.7
1991	476	7.3	0.34	6.6	8.0	424	7.4	0.37	6.7	8.1
1992	417	6.4	0.32	5.8	7.0	367	6.4	0.34	5.7	7.1
1993	427	6.4	0.31	5.8	7.0	392	6.7	0.34	6.1	7.4
1994	416	6.2	0.31	5.6	6.8	370	6.3	0.33	5.7	7.0
1995	440	6.4	0.31	5.8	7.1	394	6.6	0.34	6.0	7.3
1996	433	6.3	0.30	5.7	6.9	382	6.4	0.33	5.8	7.1
1997	432	6.2	0.30	5.7	6.9	380	6.3	0.33	5.7	7.0
1998	452	6.5	0.31	5.9	7.1	378	6.2	0.32	5.6	6.9
1999	466	6.6	0.31	6.0	7.2	418	6.8	0.34	6.2	7.5
2000	466	6.5	0.30	5.9	7.1	418	6.8	0.34	6.1	7.5
2001	464	6.5	0.30	5.9	7.1	415	6.7	0.34	6.1	7.4
2002	451	6.2	0.30	5.7	6.9	397	6.4	0.33	5.8	7.1
2003	427	5.8	0.28	5.2	6.3	376	5.8	0.31	5.3	6.5
2004	411	5.6	0.28	5.1	6.2	366	5.9	0.31	5.3	6.5
2005	441	6.0	0.29	5.4	6.6	387	6.1	0.32	5.5	6.8
2006	435	5.8	0.28	5.2	6.3	394	6.1	0.32	5.5	6.8
2007	415	5.5	0.28	5.0	6.1	366	5.7	0.31	5.1	6.4
2008	418	5.5	0.27	5.0	6.1	368	5.7	0.31	5.1	6.3
2009	461	6.1	0.29	5.5	6.7	405	6.3	0.32	5.7	6.9
2010	437	5.6	0.28	5.1	6.2	377	5.8	0.30	5.2	6.4
2011	397	5.0	0.26	4.5	5.5	348	5.2	0.29	4.7	5.8
2012	480	6.1	0.29	5.6	6.7	407	6.2	0.31	5.6	6.8

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

*Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.*

** Statistics not shown due to fewer than 10 cases*

Invasive Cancer Mortality by Race, Illinois 1986-2012

Leukemia										
Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	1,213	5.6	0.16	5.3	5.9	136	3.0	0.27	2.4	3.5
1986	41	6.4	1.04	4.6	8.8	*	*	*	*	*
1987	39	6.3	1.03	4.4	8.6	*	*	*	*	*
1988	41	6.5	1.05	4.6	8.9	*	*	*	*	*
1989	34	4.9	0.86	3.4	6.9	*	*	*	*	*
1990	49	7.2	1.06	5.3	9.6	*	*	*	*	*
1991	47	6.6	0.99	4.8	8.8	*	*	*	*	*
1992	46	6.5	0.97	4.7	8.6	*	*	*	*	*
1993	33	4.2	0.75	2.9	6.0	*	*	*	*	*
1994	42	5.7	0.90	4.1	7.7	*	*	*	*	*
1995	42	5.5	0.86	3.9	7.4	*	*	*	*	*
1996	45	6.0	0.91	4.4	8.0	*	*	*	*	*
1997	49	6.2	0.90	4.6	8.2	*	*	*	*	*
1998	68	8.5	1.04	6.6	10.8	*	*	*	*	*
1999	46	5.5	0.82	4.0	7.3	*	*	*	*	*
2000	45	5.2	0.79	3.8	7.0	*	*	*	*	*
2001	46	5.4	0.81	4.0	7.3	*	*	*	*	*
2002	50	6.1	0.87	4.5	8.0	*	*	*	*	*
2003	48	5.7	0.83	4.2	7.5	*	*	*	*	*
2004	41	4.9	0.77	3.5	6.6	*	*	*	*	*
2005	46	5.3	0.79	3.9	7.1	*	*	*	*	*
2006	38	4.4	0.72	3.1	6.0	*	*	*	*	*
2007	40	4.4	0.71	3.2	6.0	*	*	*	*	*
2008	45	4.8	0.72	3.5	6.4	*	*	*	*	*
2009	50	5.6	0.80	4.2	7.4	*	*	*	*	*
2010	48	5.1	0.74	3.7	6.8	12	3.8	1.16	1.9	6.8
2011	38	3.8	0.63	2.7	5.2	11	4.2	1.33	2.1	7.6
2012	56	5.8	0.78	4.3	7.5	17	5.9	1.49	3.3	9.5

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Other, Ill-defined and Unknown Sites

Both Sexes

Year	All Races					White				
	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	55,998	17.3	0.07	17.2	17.4	47,269	16.9	0.08	16.7	17.0
1986	1,836	17.5	0.42	16.7	18.4	1,538	16.5	0.43	15.7	17.4
1987	1,788	16.9	0.40	16.1	17.7	1,514	16.1	0.42	15.3	16.9
1988	2,009	18.9	0.43	18.1	19.8	1,695	18.0	0.44	17.1	18.9
1989	2,043	19.2	0.43	18.3	20.0	1,715	18.2	0.44	17.3	19.1
1990	1,923	17.8	0.41	17.0	18.6	1,636	17.1	0.43	16.3	17.9
1991	1,854	16.9	0.40	16.2	17.7	1,595	16.5	0.42	15.7	17.4
1992	2,063	18.6	0.41	17.8	19.4	1,754	17.9	0.43	17.1	18.8
1993	2,108	18.7	0.41	17.9	19.5	1,782	18.0	0.43	17.2	18.9
1994	2,029	17.9	0.40	17.1	18.7	1,734	17.4	0.42	16.6	18.3
1995	2,033	17.7	0.39	16.9	18.5	1,724	17.2	0.41	16.4	18.0
1996	2,021	17.5	0.39	16.8	18.3	1,707	16.9	0.41	16.1	17.7
1997	1,993	17.1	0.38	16.3	17.8	1,688	16.6	0.40	15.8	17.4
1998	1,929	16.4	0.37	15.6	17.1	1,619	15.8	0.39	15.0	16.6
1999	2,360	19.8	0.41	19.0	20.6	1,962	18.9	0.43	18.1	19.8
2000	2,306	19.2	0.40	18.4	20.0	1,954	18.8	0.42	17.9	19.6
2001	2,216	18.3	0.39	17.5	19.1	1,853	17.7	0.41	16.9	18.5
2002	2,256	18.4	0.39	17.7	19.2	1,903	18.0	0.41	17.2	18.9
2003	2,141	17.3	0.37	16.6	18.1	1,795	16.9	0.40	16.1	17.7
2004	2,141	17.1	0.37	16.4	17.9	1,821	17.0	0.40	16.2	17.8
2005	2,191	17.3	0.37	16.6	18.1	1,847	17.1	0.40	16.3	17.9
2006	2,115	16.6	0.36	15.9	17.3	1,785	16.4	0.39	15.6	17.2
2007	2,027	15.6	0.35	15.0	16.3	1,688	15.3	0.38	14.6	16.0
2008	2,132	16.2	0.35	15.5	16.9	1,788	16.0	0.38	15.3	16.8
2009	2,133	16.0	0.35	15.3	16.7	1,826	16.2	0.38	15.4	17.0
2010	2,082	15.3	0.34	14.6	16.0	1,727	15.1	0.37	14.3	15.8
2011	2,110	15.3	0.34	14.7	16.0	1,778	15.3	0.37	14.6	16.0
2012	2,159	15.5	0.34	14.8	16.1	1,841	15.6	0.37	14.9	16.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Other, Ill-defined and Unknown Sites

Both Sexes										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	8,158	22.2	0.25	21.7	22.7	570	7.2	0.33	6.6	7.9
1986	293	27.2	1.67	24.1	30.7	*	*	*	*	*
1987	266	24.6	1.57	21.6	27.9	*	*	*	*	*
1988	306	27.2	1.61	24.1	30.5	*	*	*	*	*
1989	315	27.6	1.61	24.5	30.9	13	9.4	3.04	4.4	16.9
1990	277	24.2	1.50	21.3	27.3	10	7.2	2.41	3.3	13.3
1991	247	20.8	1.36	18.2	23.6	12	6.5	1.96	3.2	11.5
1992	296	24.8	1.48	22.0	27.9	13	8.6	2.68	4.1	15.1
1993	314	25.8	1.50	22.9	28.9	12	6.7	2.16	3.1	12.0
1994	286	23.1	1.41	20.4	26.0	*	*	*	*	*
1995	291	23.1	1.39	20.4	25.9	18	8.2	2.28	4.4	13.7
1996	305	23.6	1.38	21.0	26.4	*	*	*	*	*
1997	289	22.1	1.34	19.5	24.8	16	6.0	1.65	3.2	10.0
1998	289	21.6	1.30	19.1	24.2	21	7.7	1.89	4.4	12.2
1999	378	28.6	1.51	25.7	31.6	20	6.7	1.67	3.9	10.7
2000	330	24.1	1.36	21.6	26.9	22	8.4	1.96	5.0	13.0
2001	338	24.6	1.37	21.9	27.4	25	6.6	1.40	4.2	9.9
2002	325	23.3	1.32	20.7	26.0	28	8.0	1.68	5.1	11.9
2003	317	22.3	1.28	19.9	25.0	29	8.9	1.81	5.8	13.1
2004	295	20.1	1.20	17.8	22.6	25	7.6	1.67	4.7	11.5
2005	312	20.9	1.21	18.6	23.4	32	8.9	1.71	5.9	12.8
2006	299	19.6	1.16	17.4	22.0	31	8.4	1.66	5.4	12.1
2007	319	20.8	1.19	18.5	23.2	20	5.2	1.25	3.1	8.2
2008	312	19.7	1.14	17.5	22.0	32	6.9	1.34	4.6	10.0
2009	274	17.0	1.05	15.0	19.2	33	7.0	1.32	4.6	10.0
2010	318	18.9	1.09	16.8	21.1	37	6.9	1.23	4.8	9.7
2011	285	17.1	1.04	15.1	19.2	47	9.0	1.39	6.5	12.1
2012	282	16.5	1.01	14.6	18.6	36	6.6	1.15	4.5	9.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Other, Ill-defined and Unknown Sites

Males										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	28,681	21.5	0.13	21.3	21.8	24,056	20.9	0.14	20.7	21.2
1986	942	21.9	0.75	20.4	23.4	785	20.5	0.76	19.1	22.1
1987	906	20.9	0.73	19.5	22.4	750	19.5	0.74	18.1	21.1
1988	1,043	24.4	0.79	22.9	26.0	865	23.1	0.82	21.5	24.7
1989	1,089	24.9	0.79	23.4	26.5	891	23.3	0.81	21.7	24.9
1990	955	21.6	0.73	20.2	23.0	789	20.3	0.75	18.8	21.8
1991	957	21.6	0.72	20.2	23.0	807	20.7	0.75	19.3	22.3
1992	1,043	23.3	0.74	21.8	24.8	877	22.4	0.78	20.9	24.0
1993	1,041	23.1	0.74	21.7	24.6	876	22.2	0.77	20.7	23.8
1994	976	20.9	0.69	19.6	22.3	829	20.4	0.72	19.0	21.9
1995	1,023	21.8	0.70	20.4	23.2	861	21.1	0.73	19.7	22.6
1996	1,033	21.9	0.70	20.5	23.3	861	21.0	0.73	19.6	22.5
1997	968	20.2	0.66	18.9	21.6	821	19.7	0.70	18.4	21.1
1998	960	19.7	0.65	18.4	21.0	797	18.9	0.68	17.6	20.3
1999	1,152	23.9	0.72	22.5	25.3	956	22.8	0.75	21.4	24.4
2000	1,169	23.7	0.70	22.4	25.2	979	23.0	0.74	21.5	24.5
2001	1,137	23.0	0.69	21.6	24.4	946	22.1	0.73	20.7	23.6
2002	1,175	23.5	0.70	22.1	24.9	987	22.9	0.74	21.5	24.4
2003	1,081	21.4	0.66	20.1	22.7	905	20.8	0.70	19.4	22.2
2004	1,093	21.3	0.65	20.0	22.6	931	21.1	0.70	19.8	22.5
2005	1,109	21.3	0.65	20.1	22.6	929	20.8	0.69	19.5	22.2
2006	1,121	21.1	0.64	19.9	22.4	952	21.0	0.69	19.7	22.4
2007	1,093	20.3	0.62	19.1	21.6	905	19.7	0.66	18.5	21.1
2008	1,076	19.6	0.61	18.4	20.8	899	19.2	0.65	18.0	20.5
2009	1,116	19.9	0.61	18.7	21.1	970	20.3	0.66	19.1	21.7
2010	1,120	19.4	0.59	18.3	20.6	934	19.1	0.64	17.9	20.4
2011	1,130	19.3	0.59	18.1	20.4	947	19.0	0.63	17.8	20.3
2012	1,173	19.7	0.59	18.6	20.9	1,007	20.1	0.64	18.8	21.4

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Other, Ill-defined and Unknown Sites

Males										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	4,327	28.9	0.47	28.0	29.8	298	8.3	0.55	7.2	9.4
1986	154	34.3	3.01	28.7	40.7	*	*	*	*	*
1987	152	34.5	3.05	28.8	40.9	*	*	*	*	*
1988	173	37.0	3.00	31.3	43.3	*	*	*	*	*
1989	191	40.2	3.12	34.3	46.7	*	*	*	*	*
1990	161	34.6	2.93	29.1	40.7	*	*	*	*	*
1991	144	30.1	2.67	25.1	35.7	*	*	*	*	*
1992	160	32.2	2.72	27.1	38.0	*	*	*	*	*
1993	158	31.9	2.73	26.8	37.6	*	*	*	*	*
1994	143	27.2	2.43	22.7	32.4	*	*	*	*	*
1995	150	28.5	2.49	23.9	33.8	12	9.7	3.06	4.6	17.7
1996	165	31.2	2.60	26.4	36.7	*	*	*	*	*
1997	140	26.5	2.41	22.0	31.6	*	*	*	*	*
1998	153	26.6	2.25	22.4	31.4	10	7.7	2.71	3.3	14.8
1999	187	35.5	2.78	30.2	41.3	*	*	*	*	*
2000	177	32.6	2.60	27.7	38.1	13	11.8	3.71	5.7	20.9
2001	178	33.0	2.64	28.0	38.5	13	7.0	2.08	3.5	12.5
2002	167	28.1	2.27	23.8	32.9	21	13.3	3.37	7.6	21.4
2003	161	29.5	2.50	24.8	34.7	15	11.2	3.25	5.7	19.0
2004	152	25.9	2.28	21.7	30.7	10	6.7	2.40	2.8	12.7
2005	164	27.1	2.25	22.9	31.9	16	11.4	3.28	5.9	19.2
2006	154	24.8	2.14	20.8	29.3	15	8.0	2.46	3.9	14.0
2007	175	28.0	2.26	23.7	32.7	13	6.8	2.07	3.4	12.0
2008	161	25.8	2.16	21.7	30.3	16	7.9	2.27	4.1	13.4
2009	133	20.1	1.84	16.6	24.0	13	7.2	2.26	3.4	12.7
2010	162	24.6	2.06	20.7	28.9	24	10.1	2.27	6.1	15.4
2011	164	24.5	2.03	20.6	28.7	19	7.0	1.70	4.1	11.2
2012	148	20.4	1.77	17.1	24.1	18	7.0	1.75	4.0	11.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Other, Ill-defined and Unknown Sites

Females										
All Races						White				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	27,317	14.4	0.09	14.2	14.6	23,213	14.1	0.09	13.9	14.3
1986	894	14.5	0.49	13.6	15.5	753	13.8	0.51	12.8	14.8
1987	882	14.2	0.48	13.3	15.2	764	13.8	0.51	12.9	14.9
1988	966	15.4	0.50	14.4	16.4	830	14.9	0.52	13.9	15.9
1989	954	15.2	0.50	14.2	16.2	824	14.8	0.53	13.8	15.9
1990	968	15.2	0.49	14.3	16.2	847	15.0	0.52	14.0	16.1
1991	897	13.7	0.46	12.8	14.7	788	13.6	0.49	12.6	14.6
1992	1,020	15.5	0.49	14.6	16.5	877	15.2	0.52	14.2	16.3
1993	1,067	15.9	0.49	15.0	16.9	906	15.4	0.52	14.4	16.4
1994	1,053	15.6	0.48	14.6	16.5	905	15.2	0.51	14.2	16.2
1995	1,010	14.8	0.47	13.9	15.8	863	14.5	0.50	13.5	15.5
1996	988	14.4	0.46	13.5	15.3	846	14.0	0.49	13.1	15.0
1997	1,025	14.9	0.47	14.0	15.9	867	14.5	0.50	13.6	15.5
1998	969	13.9	0.45	13.0	14.8	822	13.5	0.48	12.6	14.5
1999	1,208	17.0	0.49	16.0	18.0	1,006	16.2	0.52	15.2	17.3
2000	1,137	16.0	0.48	15.1	17.0	975	15.8	0.51	14.8	16.8
2001	1,079	15.2	0.47	14.3	16.2	907	14.9	0.50	13.9	15.9
2002	1,081	15.0	0.46	14.1	15.9	916	14.8	0.50	13.8	15.8
2003	1,060	14.6	0.45	13.7	15.5	890	14.2	0.48	13.3	15.2
2004	1,048	14.3	0.45	13.5	15.2	890	14.2	0.48	13.2	15.1
2005	1,082	14.8	0.45	13.9	15.7	918	14.7	0.49	13.8	15.7
2006	994	13.4	0.43	12.6	14.3	833	13.1	0.46	12.2	14.1
2007	934	12.3	0.41	11.5	13.2	783	12.1	0.44	11.3	13.0
2008	1,056	13.9	0.44	13.1	14.8	889	13.8	0.47	12.9	14.8
2009	1,017	13.2	0.42	12.4	14.1	856	13.2	0.46	12.3	14.1
2010	962	12.3	0.41	11.6	13.2	793	12.1	0.44	11.2	13.0
2011	980	12.4	0.40	11.6	13.2	831	12.5	0.45	11.7	13.4
2012	986	12.3	0.40	11.5	13.1	834	12.3	0.44	11.5	13.2

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases

Invasive Cancer Mortality by Race, Illinois 1986-2012

Other, Ill-defined and Unknown Sites

Females										
Black						Asian or Other Races				
Year	Count	Rate	SE	LCI	UCI	Count	Rate	SE	LCI	UCI
1986-2012	3,831	17.7	0.29	17.1	18.3	272	6.3	0.41	5.5	7.2
1986	139	22.3	1.94	18.6	26.4	*	*	*	*	*
1987	114	18.1	1.74	14.9	21.8	*	*	*	*	*
1988	133	20.2	1.79	16.8	24.0	*	*	*	*	*
1989	124	18.8	1.72	15.6	22.4	*	*	*	*	*
1990	116	17.3	1.63	14.3	20.8	*	*	*	*	*
1991	103	14.6	1.46	11.8	17.7	*	*	*	*	*
1992	136	19.6	1.70	16.4	23.2	*	*	*	*	*
1993	156	21.6	1.76	18.3	25.3	*	*	*	*	*
1994	143	19.8	1.68	16.6	23.3	*	*	*	*	*
1995	141	19.2	1.63	16.1	22.6	*	*	*	*	*
1996	140	18.4	1.57	15.5	21.7	*	*	*	*	*
1997	149	19.1	1.59	16.1	22.5	*	*	*	*	*
1998	136	17.4	1.51	14.6	20.6	11	7.5	2.50	3.4	13.8
1999	191	24.1	1.76	20.8	27.8	11	6.8	2.23	3.2	12.5
2000	153	18.7	1.52	15.9	21.9	*	*	*	*	*
2001	160	19.3	1.54	16.4	22.6	12	6.1	1.84	3.1	10.8
2002	158	19.2	1.54	16.3	22.4	*	*	*	*	*
2003	156	18.3	1.48	15.5	21.5	14	7.3	2.10	3.8	12.5
2004	143	16.6	1.40	14.0	19.6	15	8.1	2.22	4.3	13.5
2005	148	16.9	1.40	14.2	19.8	16	7.6	2.00	4.2	12.5
2006	145	16.3	1.36	13.7	19.2	16	8.2	2.17	4.5	13.4
2007	144	16.0	1.34	13.4	18.8	*	*	*	*	*
2008	151	16.0	1.31	13.5	18.8	16	6.3	1.66	3.5	10.3
2009	141	14.9	1.27	12.5	17.6	20	7.1	1.68	4.2	11.2
2010	156	15.6	1.26	13.2	18.2	13	4.4	1.28	2.3	7.6
2011	121	12.3	1.13	10.2	14.7	28	10.1	1.98	6.6	14.7
2012	134	13.6	1.19	11.4	16.1	18	6.1	1.50	3.5	9.7

Rates are expressed per 100,000 population. Age-adjustment was by the direct method adjusting to the 2000 U.S. standard population. Standard error (SE), lower (LCI) and upper confidence intervals (UCI) computed at the 95 percent level using Tiwari method.

Source: Surveillance, Epidemiology, and End Results (SEER) Program, SEER*Stat Database: Mortality - All COD, Aggregated With County, Total U.S. (1969-2012) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015. Underlying mortality data provided by NCHS.

* Statistics not shown due to fewer than 10 cases