ILLINOIS DEPARTMENT OF PUBLIC HEALTH



# Illinois Influenza Surveillance Report

Week 48: Week Ending Saturday, December 3, 2011

Division of Infectious Diseases Immunizations Section 12/9/2011

# Week 48: November 27-December 3, 2011

# **Contents**

Summary	3
CDC Flu View	4
ILI Net Provider Surveillance	5
ILI Visits by Age Group	6
Great Lakes Naval Recruit Influenza Surveillance	7
Influenza Intensive Care Unit Admissions and Deaths	7
Influenza Related ICU Admissions by Age Group	8
Laboratory Surveillance	9
Viral Resistance	10
Influenza Outbreaks Reported in Long-Term Facilities and Nursing Homes	10
Weekly Viral Subtype	11
Resources	12

#### **Summary**

- During CDC Week 48, the proportion of outpatient visits for influenza-like illness (ILI)<sup>1</sup> was 1.2% compared with 2.0% for week 47.
- The risk of illness due to influenza viruses is low. Numerous respiratory viruses can cause ILI, and these viruses are now more prevalent than influenza viruses.
- Based on CDC criteria, influenza activity is classified as **no activity** (see CDC FLU View Section) for week 48. Last week, influenza activity was classified as sporadic.
- Febrile Respiratory Illness (FRI) surveillance<sup>2</sup> at Naval Recruit Training Command, Great Lakes for week 48 is at or below expected value.
- During week 48, none of the specimens tested by Illinois Department of Public Health Laboratory were positive for influenza.
- There were no influenza-associated Intensive Care Unit (ICU) admissions<sup>3</sup> reported for week 48.
- No influenza-associated pediatric deaths were reported for week 48.
- During week 48, no influenza outbreaks were reported.

<sup>&</sup>lt;sup>1</sup> ILI "Influenza like Illness" is defined as fever ≥ 100°F and cough and/or sore throat.

FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:


<sup>1.</sup> At or below expected value (expected value shown as dashed line)

Moderately elevated

Substantially elevated

<sup>&</sup>lt;sup>3</sup> For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (DFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season.

#### **CDC Flu View**



**No activity:** No laboratory confirmed cases of influenza and no reported increase in cases of influenza like illness (ILI).

**Sporadic:** Small numbers of laboratory confirmed influenza cases or a single laboratory confirmed influenza in a single region of the state.


**Local:** Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in a single region of the state.

**Regional:** Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state.

**Widespread:** Outbreaks of influenza or increases in ILI cases and recent laboratory confirmed influenza in at least half the regions in the state.


### **ILI Net Provider Surveillance**

#### Influenza Like Illness Outpatient Surveillance 2010-2012



## **ILI Visits by Age Group**

2010 -12 INFLUENZA SEASON PROPORTION OF ILI OFFICE VISITS BY AGE GROUP



#### **Great Lakes Naval Recruit Influenza Surveillance**

Febrile Respiriratory Illness (FRI) surveillance<sup>4</sup> at Naval Recruit Training Command, Great Lakes **at or below expected value** for week ending December 3, 2011. For more information visit <a href="http://www.med.navy.mil/sites/nhrc/geis/Pages/default.aspx">http://www.med.navy.mil/sites/nhrc/geis/Pages/default.aspx</a>

#### **Influenza Intensive Care Unit Admissions and Deaths**

There were no influenza related ICU admissions or deaths for week ending December 3, 2011.

Week No	Admissions No	Deaths
45	1	0
46	0	0
47	0	0
48	0	0


<sup>&</sup>lt;sup>4</sup> FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:

<sup>4.</sup> At or below expected value (expected value shown as dashed line)

Moderately elevated

<sup>6.</sup> Substantially elevated

#### Influenza Related ICU Admissions by Age Group



#### **Laboratory Surveillance**

During week 48, six specimens were tested by Illinois WHO/NREVSS<sup>[1]</sup> collaborating laboratories in Illinois. This represents a decrease in testing compared with week 47. Of the six specimens tested, none tested positive for influenza B during week 47.

Week	No of Labs Reporting	A (H1)	2009(A)H1N1	A (H3)	A(Unable to subtype)	A(Sub typing not performed)	В	Total # Tested	% Positive
45	1	0	0	0	0	0	0	4	0
46	1	0	0	0	0	0	3	14	21%
47	1	0	0	0	0	0	5	14	36%
48	1	0	0	0	0	0	0	6	0

<sup>[1]</sup> WHO/NREVSS Collaborating Laboratories WHO/NREVSS Collaborating Laboratories: For viral surveillance - About 80 U.S. World Health Organization (WHO) Collaborating Laboratories and 70 National Respiratory and Enteric Virus Surveillance System (NREVSS), located throughout the United States report daily or weekly the results of their testing.

#### **Viral Resistance**

# Neuraminidase Inhibitor Resistance Testing Results on Samples Collected Since October 1, 2011


	Oselt	amivir	Zanamivir		
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)	
Influenza A (H3N2)	41	0 (0.0)	41	0 (0.0)	
Influenza B	2	0 (0.0)	2	0 (0.0)	
Influenza A (2009 H1N1)	10	0 (0.0)	10	0 (0.0)	


All viruses tested for the 2011-2012 season since October 1, 2011 have been susceptible to the neuraminidase inhibitor antiviral medications oseltamivir and zanamivir as were the majority of viruses tested last season; however, rare sporadic cases of oseltamivir resistant 2009 influenza A (H1N1) and A (H3N2) viruses have been detected worldwide. Antiviral treatment with oseltamivir or zanamivir is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at <a href="http://www.cdc.gov/flu/antivirals/index.htm">http://www.cdc.gov/flu/antivirals/index.htm</a>.

#### Influenza Outbreaks Reported in Long-Term Facilities and Nursing Homes

There were no outbreaks reported for week 48.

# **Weekly Viral Subtype**





#### Resources

- Centers for Disease Control and Prevention Influenza Website: <a href="http://www.cdc.gov/flu/">http://www.cdc.gov/flu/</a>
- Immunization Action Coalition Website: <a href="http://immunize.org/">http://immunize.org/</a>
- IDPH Website: http://www.idph.state.il.us/flu/surveillance.htm
- ACL Clinical Laboratory Respiratory Panel: <a href="http://www.acllaboratories.com/">http://www.acllaboratories.com/</a>
- St Louis Children's Hospital Clinical Laboratory Respiratory Panel: http://www.stlouischildrens.org/content/ClinicalLaboratories.htm

<sup>&</sup>lt;sup>i</sup> For more information regarding circulating respiratory viruses, see

a. http://www.cdc.gov/surveillance/nrevss/rsv/state.html

b. http://www.acllaboratories.com/

c. www.stlouischildrens.org/content/ClinicalLaboratories.htm