ILLINOIS DEPARTMENT OF PUBLIC HEALTH



Illinois Influenza Surveillance Report

Week 46: Week Ending Saturday, November 17, 2012

Division of Infectious Diseases Immunization Section 11/26/2012

Contents

Summary	3
CDC Flu View	4
ILI Net Provider Surveillance	5
ILI Visits by Age Group	6
Great Lakes Naval Recruit Influenza Surveillance	7
Influenza Intensive Care Unit Admissions and Deaths	7
Influenza Related ICU Admissions by Age Group	8
Laboratory Surveillance	9
Viral Resistance	10
Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)	10
IDPH, Immunization Section Regional Map	11
Weekly Viral Subtype	12
Resources	13

Summary

- During CDC surveillance week 46, the proportion of outpatient visits for influenza-like illness (ILI)¹ • was 2.2% compared with 1.6 % for week 45.
- Based on CDC criteria, influenza activity is classified as SPORADIC (see CDC FLU View Section) for week 46.
- Febrile Respiratory Illness (FRI) surveillance² at Naval Recruit Training Command, Great Lakes was at or below expected value.
- During week 46, one specimen was tested for influenza by Illinois Department of Public Health Laboratory, Specimen tested positive for Influenza A.
- Two influenza-associated Intensive Care Unit (ICU) admissions³ were reported during week 46.
- No influenza-associated pediatric death was reported for week 46.
- During week 46, no influenza outbreak was reported in a long-term care facility.

¹ ILI "Influenza like Illness" is defined as fever ≥ 100°F and cough and/or sore throat.

² FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:


At or below expected value (expected value shown as dashed line) 1.

^{2.} Moderately elevated 3.

Substantially elevated

³ For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (DFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season.

CDC Flu View



*This map indicates geographic spread and does not measure the severity of influenza activity.

No activity: No laboratory confirmed cases of influenza and no reported increase in cases of influenza like illness (ILI).


Sporadic: Small numbers of laboratory confirmed influenza cases or a single laboratory confirmed influenza in a single region of the state.

Local: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in a single region of the state.

Regional: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state.


Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory confirmed influenza in at least half the regions in the state.

ILI Net Provider Surveillance



ILI Visits by Age Group

2011-13 INFLUENZA SEASON PROPORTION OF ILI OFFICE VISITS BY AGE GROUP



WEEK ENDING

NUMBER OF VISITS

Great Lakes Naval Recruit Influenza Surveillance

Febrile Respiratory Illness (FRI) surveillance⁴ at Naval Recruit Training Command, Great Lakes was **at or below expected value** for week ending November 3, 2012. For more information visit http://www.med.navy.mil/sites/nhrc/geis/Pages/default.aspx

Influenza Intensive Care Unit Admissions and Deaths

There were two influenza related ICU admissions and no deaths reported for week ending November 3, 2012.


Year	Week No	Admissions No	Deaths
2012	42	1	0
2012	43	1	0
2012	44	1	0
2012	45	0	0
2012	46	2	0

 ⁴ FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:
At or below expected value (expected value shown as dashed line)

^{5.} Moderately elevated

^{6.} Substantially elevated

Influenza Related ICU Admissions by Age Group



WEEK ENDING

Laboratory Surveillance

During week 44, no specimen was tested for Influenza by Illinois Department of Public Health Laboratory.

Year	Week	A (H1)	2009(A)H1N1	A (H3)	A(Unable to subtype)	A(Sub typing not performed)	В	Total # Tested	% Positive
2012	43	0	0	0	0	0	0	0	0%
2012	44	0	0	0	0	0	0	0	0%
2012	45	0	0	0	0	0	0	2	0%
2012	46	1	0	0	0	0	0	1	100%

Viral Resistance

The majority of currently circulating influenza viruses are susceptible to the neuraminidase inhibitor antiviral medications oseltamivir and zanamivir; however, rare sporadic cases of oseltamivir resistant 2009 influenza A (H1N1) and A (H3N2) viruses have been detected worldwide.

Neuraminidase Inhibitor Resistance Testing Results on Samples Collected Since October 1, 2012							
	Oselta	amivir	Zanamivir				
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)			
Influenza A (H3N2)	43	0 (0.0)	43	0 (0.0)			
Influenza B	37	0 (0.0)	37	0 (0.0)			
2009 H1N1	1	0 (0.0)	1	0 (0.0)			

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among 2009 H1N1 and A (H3N2) viruses (the adamantanes do not have activity against influenza B viruses). Antiviral treatment as early as possible with oseltamivir or zanamivir is recommended for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for influenza-related complications. Additional information treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at http://www.cdc.gov/flu/antivirals/index.htm.

Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)


No influenza outbreaks were reported in long-term care facilities within Illinois for week ending November 17, 2012.





Weekly Viral Subtype

Influenza Isolates from Illinois Reported by WHO/NREVSS Collaborating Laboratories 2011-2013



Resources

- Centers for Disease Control and Prevention Influenza Website: <u>http://www.cdc.gov/flu/</u>
- Immunization Action Coalition Website: <u>http://immunize.org/</u>
- IDPH Website: http://www.idph.state.il.us/flu/surveillance.htm
- ACL Clinical Laboratory Respiratory Panel: <u>http://www.acllaboratories.com/</u>
- St Louis Children's Hospital Clinical Laboratory Respiratory Panel: http://www.stlouischildrens.org/health-care-professionals/clinical-laboratories