

Week 52: December 22 – December 28, 2013

ILLINOIS DEPARTMENT OF PUBLIC HEALTH



Illinois Influenza Surveillance Report

Week 52: Week Ending Saturday, December 28, 2013

Division of Infectious Diseases Communicable Disease Section

1/3/2014

Contents

Summary	3
CDC FluView	4
ILINet Provider Surveillance.....	5
ILI Visits by Age Group	6
Influenza Intensive Care Unit (ICU) Admissions and Deaths	7
Influenza Related ICU Admissions by Age Group, 2012-2014.....	8
Laboratory Surveillance	9
Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)	10
Viral Resistance	10
Weekly Viral Subtype	11
IDPH Infectious Diseases Regional Map	12
Resources	13

Summary

- For this reporting week, the proportion of outpatient visits for influenza-like illness (ILI)¹ was **4.17%**, which is **above** the regional baseline of **1.60%**.
- Based on CDC criteria, Illinois influenza activity is classified as **Widespread** (see CDC FluView Section) for this reporting week.
- For this reporting week there were **506** influenza specimens tested by WHO/NREVSS collaborating Illinois laboratories (which includes all Illinois Department of Public Health Laboratories). **150** specimens tested positive for Influenza.
- **Forty-two** influenza-associated Intensive Care Unit (ICU) admissions³ were reported for this reporting week.
- **No** influenza-associated pediatric deaths were reported for this reporting week.
- For this reporting week, **one** influenza outbreak was reported in an assisted living facility in Whiteside County.

Novel Influenza A

H9N2

- China has reported two cases of H9N2.
- One case is in a 7-year-old boy and one is in an 86-year-old man.
- These are the first cases of H9N2 in 4 years
- For additional information on these cases, visit the Center for Infectious Disease Research and Policy <http://www.cidrap.umn.edu/news-perspective/2014/01/flu-scan-jan-02-2014>

H7N9

- Taiwan has confirmed its second case of H7N9 in an 86-year-old man.
- For additional information on this case, visit the Center for Infectious Disease Research and Policy <http://www.cidrap.umn.edu/news-perspective/2013/12/news-scan-dec-31-2013>


¹ ILI "Influenza like Illness" is defined as fever \geq 100°F and cough and/or sore throat.

² FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:

1. At or below expected value
2. Moderately elevated
3. Substantially elevated

³ For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (DFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season.

CDC FluView



* This map indicates geographic spread & does not measure the severity of influenza activity

No activity: No laboratory confirmed cases of influenza and no reported increase in cases of influenza like illness (ILI).

Sporadic: Small numbers of laboratory confirmed influenza cases or a single laboratory confirmed influenza in a single region of the state.


Local: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in a single region of the state.

Regional: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state.

Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory confirmed influenza in at least half the regions in the state.


ILINet Provider Surveillance

Influenza Like Illness Outpatient Surveillance 2012-2014



ILI Visits by Age Group

2012-2014 INFLUENZA SEASON PROPORTION OF ILI OFFICE VISITS BY AGE GROUP




Influenza Intensive Care Unit (ICU) Admissions and Deaths

There were **42** influenza related ICU admissions and **2** deaths for this reporting week.

Year	Week No	Admissions	Deaths ⁴
2013	40	2	0
2013	41	1	0
2013	42	1	0
2013	43	3	0
2013	44	1	0
2013	45	5	1
2013	46	2	0
2013	47	2	1
2013	48	6	1
2013	49	4	0
2013	50	23	0
2013	51	30	1
2013	52	42	2
Total (Provisional) for 2013-14 Season		122	6

Influenza Related ICU Admissions by Age Group, 2012-2014



Laboratory Surveillance

- For this reporting week there were **506** influenza specimens tested by WHO/NREVSS collaborating Illinois laboratories (which includes all Illinois Department of Public Health Laboratories). **150** specimens tested positive for Influenza.

For more information about circulating viruses visit:

- St Louis Children’s Hospital Weekly Virus/Microbiology Update:
<http://slchlabtestguide.bjc.org/Default.aspx?url=63e0653d-fe31-466f-9228-d4de90fa7424>

Year	Week	A (H1)	2009(A) H1N1	A (H3)	A (Unable to subtype)	A (Sub typing not performed)	B	Total # Tested	Total # Positive	% Positive
2013	46	0	2	0	0	0	1	185	3	2.0%
2013	47	0	2	0	0	1	0	132	3	2.0%
2013	48	0	2	0	0	5	1	162	8	5.0%
2013	49	0	7	1	0	4	0	194	12	6.0%
2013	50	0	19	0	0	5	5	208	29	14.0%
2013	51	0	21	0	0	30	0	267	51	19.0%
2013	52	0	37	1	0	109	3	506	150	30.0%
Totals		0	92	3	0	156	10	2,215	261	11.78 %

Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)

There was **one** outbreak reported in long-term care facilities for this reporting week.

Region	2013-2014 Influenza Season - Number of outbreaks (%)
Rockford (1)	1(100%)
Peoria (2),	0(0)
Edwardsville (4),	0(0)
Marion (5),	0(0)
Champaign (6),	0(0)
West Chicago (7)	0(0)
Chicago/Cook (8)	0(0)
Total	0(0)

Viral Resistance:

Antiviral Resistance: Testing of 2009 H1N1, influenza A (H3N2), and influenza B virus isolates for resistance to neuraminidase inhibitors (oseltamivir and zanamivir) is performed at CDC using a functional assay. Additional 2009 H1N1 and influenza A (H3N2) clinical samples are tested for mutations of the virus known to confer oseltamivir resistance. The data summarized below combine the results of both testing methods. These samples are routinely obtained for surveillance purposes rather than for diagnostic testing of patients suspected to be infected with antiviral-resistant virus.

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among 2009 influenza A (H1N1) and A (H3N2) viruses (the adamantanes are not effective against influenza B viruses). As a result, data from adamantane resistance testing are not presented below.

**Neuraminidase Inhibitor Resistance Testing Results
on Samples Collected Since October 1, 2013**

	Oseltamivir		Zanamivir	
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)
Influenza A (H3N2)	66	0 (0.0)	66	0 (0.0)
Influenza B	15	0 (0.0)	15	0 (0.0)
2009 H1N1	910*	10 (1.1)	482	0 (0.0)


*Includes specimens tested in national surveillance and additional specimens tested at public health laboratories in 11 states (AZ, CO, FL, GA, HI, MD, MI, NY, TX, WA, and WI) who share testing results with CDC.

The majority of currently circulating influenza viruses are susceptible to the neuraminidase inhibitor antiviral medications oseltamivir and zanamivir; however, rare sporadic cases of oseltamivir-resistant 2009 H1N1 and A (H3N2) viruses have been detected worldwide. Antiviral treatment with oseltamivir or zanamivir is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for serious influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at <http://www.cdc.gov/flu/antivirals/index.htm>.

Weekly Viral Subtype

Influenza Isolates from Illinois Reported by WHO/NREVSS Collaborating Laboratories, 2013-2014 Influenza Season

Influenza Isolates from Illinois Reported by
WHO/NREVSS Collaborating Laboratories
2013-2014 Influenza Season



IDPH Infectious Diseases Regional Map



- 1 – Rockford Region
- 2 - Peoria Region
- 4 – Edwardsville Region
- 5 – Marion Region
- 6 - Champaign Region
- 7 – West Chicago Region
- 8 – Chicago/Cook Co Region

Resources

- Centers for Disease Control and Prevention Influenza Website: <http://www.cdc.gov/flu/>
- Immunization Action Coalition Website: <http://immunize.org/>
- IDPH Seasonal Influenza Website: <http://www.idph.state.il.us/flu/surveillance.htm>
- St Louis Children's Hospital Weekly Virus/Microbiology Update:
<http://slchlabtestguide.bjc.org/Default.aspx?url=63e0653d-fe31-466f-9228-d4de90fa7424>
- Center for Infectious Disease Research and Policy
 - <http://www.cidrap.umn.edu/news-perspective/2014/01/flu-scan-jan-02-2014>
 - <http://www.cidrap.umn.edu/news-perspective/2013/12/news-scan-dec-31-2013>