ILLINOIS DEPARTMENT OF PUBLIC HEALTH



Illinois Influenza Surveillance Report

Week 16: Week Ending Saturday, April 19, 2014

Division of Infectious Diseases Communicable Disease Section 4/25/2014

Contents

Summary	
CDC FluView	
ILINet Provider Surveillance	
ILI Visits by Age Group	
Influenza Intensive Care Unit (ICU) Admissions and Deaths7	
Influenza Related ICU Admissions by Age Group, 2012-2014	
Laboratory Surveillance	
Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Nursing Homes (NH)Error! Bookmark not defined	۱.
Viral Resistance	
Weekly Viral Subtype	
IDPH Infectious Diseases Regional Map12	
Resources	

Summary

- For this reporting week, the proportion of outpatient visits for influenza-like illness (ILI)¹ was **1.23%**, which is **Below** the regional baseline of **1.60%**.
- Based on CDC criteria, Illinois influenza activity is classified as **Sporadic** (see CDC FluView Section) for this reporting week.
- For this reporting week there were 163 influenza specimens tested by WHO/NREVSS collaborating Illinois laboratories (which includes all Illinois Department of Public Health Laboratories). 10 specimens tested positive for Influenza.
- **Eight** influenza-associated Intensive Care Unit (ICU) admissions³ were reported for this reporting week.
- No influenza-associated pediatric deaths were reported for this reporting week.
- For this reporting week, **No** influenza outbreaks were reported.

Novel Influenza A

H7N9

- There are multiple cases of H7N9 being reported daily.
- For up-to-date information, please visit the WHO Global Alert and Response & Center for Infectious Disease Research and Policy
 - o http://www.flutrackers.com/forum/showthread.php?t=202713 human H7N9 Case List
 - o http://www.who.int/csr/don/en/
 - o http://www.cidrap.umn.edu/news-perspective/2014/04/news-scan-apr-18-2014
 - o http://www.cidrap.umn.edu/news-perspective/2014/04/news-scan-apr-21-2014

H5

- The USDA reports a low-pathogenic H5 avain flu being detected among Japanese quail on a commercial farm.
- North Korea has experienced another H5N1 avian flu outbreak in domestic poultry.
- For up-to-date information, please visit the Center for Infectious Disease Research and Policy:
 - o http://www.cidrap.umn.edu/news-perspective/2014/04/avian-flu-scan-apr-23-2014

¹ ILI "Influenza like Illness" is defined as fever $\geq 100^{\circ}$ F and cough and/or sore throat.


² FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:

true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season. CDC FluView

At or below expected value
 Moderately elevated

^{3.} Substantially elevated

³ For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (DFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and



No activity: No laboratory confirmed cases of influenza and no reported increase in cases of influenza like illness (ILI).


Sporadic: Small numbers of laboratory confirmed influenza cases or a single laboratory confirmed influenza in a single region of the state.

Local: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in a single region of the state.

Regional: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state.


Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory confirmed influenza in at least half the regions in the state.

ILINet Provider Surveillance



ILI Visits by Age Group

NUMBER OF VISITS



6


Influenza Intensive Care Unit (ICU) Admissions and Deaths

There were 8 influenza related ICU admissions and 1 death for this reporting week.

Year	Week No	Admissions	Deaths ⁴
2013	40-52	165	7
2013/2014	01	134	6
2014	02	133	17
2014	03	61	10
2014	04	59	10
2014	05	43	14
2014	06	33	6
2014	07	16	3
2014	08	12	5
2014	09	14	3
2014	10	16	4
2014	11	8	5
2014	12	11	0
2014	13	6	0
2014	14	9	2
2014	15	6	0
2014	16	8	1
-	isional) for Season	734	93

⁴ Deaths are reported for a) adults admitted to an intensive care unit who have a positive test for influenza and b) hospitalized and non-hospitalized children (less than 18 years of age) with a positive influenza test. The degree to which influenza infection is an immediate or underlying cause of death is not ascertained. CDC Influenza-Associated Pediatric Mortality data: <u>http://gis.cdc.gov/GRASP/Fluview/PedFluDeath.html</u>

Influenza Related ICU Admissions by Age Group, 2012-2014



Laboratory Surveillance

• For this reporting week there were **163** influenza specimens tested by WHO/NREVSS collaborating Illinois laboratories (which includes all Illinois Department of Public Health Laboratories). **10** specimens tested positive for Influenza.

For more information about circulating viruses visit:

 St Louis Children's Hospital Weekly Virus/Microbiology Update: <u>http://slchlabtestguide.bjc.org/Default.aspx?url=63e0653d-fe31-466f-9228-d4de90fa7424</u>

Year	Week	A (H1)	2009(A) H1N1	A (H3)	A (Unable to subtype)	A (Sub typing not performed)	В	Total # Tested	Total # Positive	% Positive
2014	10	0	10	0	0	6	2	224	18	8.0%
2014	11	0	5	0	0	5	0	131	10	8.0%
2014	12	0	1	0	0	3	1	115	5	4.0%
2014	13	0	6	0	0	2	8	182	16	9.0%
2014	14	0	1	0	0	6	4	204	11	5.0%
2014	15	0	1	2	0	0	14	169	17	10.0%
2014	16	0	0	0	0	2	8	163	10	6.0%
Seasor	n Totals	0	440	12	0	644	75	7162	1171	16.4%

Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Correctional Facilities

There were **no** outbreaks reported for this reporting week.

Region	2013-2014 Influenza Season - Number of outbreaks (%)
Rockford (1)	4 (11%)
Peoria (2),	7 (19%)
Edwardsville (4),	5 (14%)
Marion (5),	5 (14%)
Champaign (6),	1 (3%)
West Chicago (7)	9 (24%)
Chicago/Cook (8)	6 (16%)
Total	37

Viral Resistance:

Antiviral Resistance: Testing of 2009 H1N1, influenza A (H3N2), and influenza B virus isolates for resistance to neuraminidase inhibitors (oseltamivir and zanamivir) is performed at CDC using a functional assay. Additional 2009 H1N1 and influenza A (H3N2) clinical samples are tested for mutations of the virus known to confer oseltamivir resistance. The data summarized below combine the results of both testing methods. These samples are routinely obtained for surveillance purposes rather than for diagnostic testing of patients suspected to be infected with antiviral-resistant virus.


High levels of resistance to the adamantanes (amantadine and rimantadine) persist among 2009 influenza A (H1N1) and A (H3N2) viruses (the adamantanes are not effective against influenza B viruses). As a result, data from adamantine resistance testing are not presented below.

Neuraminidase Inhibitor Resistance Testing Results on Samples Collected Since October 1, 2013							
	Oseltamivir		Oseltamivir		Zanamivir		
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)			
Influenza A (H3N2)	489	0 (0.0)	489	0 (0.0)			
Influenza B	355	0 (0.0)	355	0 (0.0)			
2009 H1N1	4,968*	59 (1.2)	1,811	0 (0.0)			

The majority of currently circulating influenza viruses are susceptible to the neuraminidase inhibitor antiviral medications oseltamivir and zanamivir; however, rare sporadic cases of oseltamivir-resistant 2009 H1N1 and A (H3N2) viruses have been detected worldwide. Antiviral treatment with oseltamivir or zanamivir is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at greater risk for serious influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at http://www.cdc.gov/flu/antivirals/index.htm.

Weekly Viral Subtype

Influenza Isolates from Illinois Reported by WHO/NREVSS Collaborating Laboratories, 2013-2014 Influenza Season





IDPH Infectious Diseases Regional Map

* The numbers in red that are located on this map indicate the number of participating Sentinel Sites per county.

Resources

- Centers for Disease Control and Prevention Influenza Website:
 <u>http://www.cdc.gov/flu/</u>
- Flu Trackers H7N9 Forum (Human Cases Line List)

 <u>http://www.flutrackers.com/forum/showthread.php?t=202713</u>
- Immunization Action Coalition Website: <u>http://immunize.org/</u>
- IDPH Seasonal Influenza Website: <u>http://www.idph.state.il.us/flu/surveillance.htm</u>
- National Respiratory and Enteric Virus Surveillance System (NREVSS), CDC website: <u>https://wwwn.cdc.gov/nrevss/account/export.aspx</u>
- St Louis Children's Hospital Weekly Virus/Microbiology Update: <u>http://slchlabtestguide.bjc.org/Default.aspx?url=63e0653d-fe31-466f-9228-d4de90fa7424</u>
- WHO Global Alert and Response H7N9:
 http://www.who.int/csr/don/en/
- Center for Infectious Disease Research and Policy:
 - o http://www.cidrap.umn.edu/news-perspective/2014/04/news-scan-apr-18-2014
 - o http://www.cidrap.umn.edu/news-perspective/2014/04/news-scan-apr-21-2014
 - o http://www.cidrap.umn.edu/news-perspective/2014/04/avian-flu-scan-apr-23-2014