ILLINOIS DEPARTMENT OF PUBLIC HEALTH



Illinois Influenza Surveillance Report

Week 11: Week Ending Saturday, March 21

Division of Infectious Diseases, Communicable Disease Section 3/27/2015

Week 11: March 15, 2015 - March 21, 2015

Contents

Summary	2
ILINet Provider Surveillance	5
ILI Visits by Age Group	6
Influenza Intensive Care Unit (ICU) Admissions and Deaths	7
Influenza Related ICU Admissions by Age Group	8
Laboratory Surveillance	9
Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Correctional Facilities	9
Weekly Viral Subtype	11
IDPH Infectious Diseases Regional Map	12
Resources	13

Summary

- For this reporting week, the proportion of outpatient visits for influenza-like illness (ILI)¹ was **2.00%**, which is **above** the regional baseline of **1.70%**.
- Based on CDC criteria, Illinois influenza activity is classified as **Regional** (see CDC FluView Section) for this reporting week.
- For this reporting week there were 430 influenza specimens tested by WHO/NREVSS collaborating Illinois laboratories and 13 tested by Illinois Department of Public Health Laboratories for a total of 443 specimens tested. 99 specimens tested positive for Influenza.
- 23 influenza-associated Intensive Care Unit (ICU) admissions³ were reported for this reporting week.
- No influenza-associated pediatric deaths were reported for this reporting week.
- For this reporting week, **one** influenza outbreaks was reported.

Since the start of the season, influenza A (H3N2) viruses have predominated nationally and in Illinois, however in recent weeks the proportion of influenza B viruses has been increasing.

Yearly influenza vaccination is recommended for everyone six months of age and older as the first and most important step in protecting against influenza infection. People should begin getting vaccinated soon after vaccine becomes available, ideally by October, to ensure that as many people as possible are protected before influenza season begins. However, as long as influenza viruses are circulating in the community, it's not too late to get vaccinated. Information on the influenza vaccine effectiveness for the 2014-2015 season can be found on the CDC 2014-2015 Flu Season website.

Additionally, clinicians are reminded of the use of neuraminidase inhibitor antiviral medications when indicated for treatment and prevention of influenza, as an adjunct to vaccination. For more information see the <u>CDC Influenza Antiviral</u> webpage.

¹ ILI "Influenza like Illness" is defined as fever ≥ 100°F and cough and/or sore throat.

² FRI surveillance is ongoing at 8 U.S. military basic training centers, representing all service branches. FRI Rate Status is classified into one of 3 categories:


^{1.} At or below expected value

^{2.} Moderately elevated

Substantially elevated

³ For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (IFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season.

CDC FluView



^{*} This map indicates geographic spread & does not measure the severity of influenza activity

No activity: No laboratory confirmed cases of influenza and no reported increase in cases of influenza like illness (ILI).

Sporadic: Small numbers of laboratory confirmed influenza cases or a single laboratory confirmed influenza in a single region of the state.


Local: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in a single region of the state.

Regional: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state.

Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory confirmed influenza in at least half the regions in the state.


ILINet Provider Surveillance





ILI Visits by Age Group

2013-2015 INFLUENZA SEASON PROPORTION OF ILI OFFICE VISITS BY AGE GROUP




Influenza Intensive Care Unit (ICU) Admissions and Deaths

There were **23** influenza related ICU admissions and **0** pediatric death for this reporting week. Preliminary data from previous weeks during this season was updated for this report.

Year	Week No	Admissions	Pediatric Deaths	
2014	40-45	20	0	
2014	46	6	0	
2014	47	15	0	
2014	48	65	0	
2014	49	122	0	
2014	50	169	0	
2014	51	217	0	
2014	52	189	0	
2014	53	149	0	
2015	01	45	0	
2015	02	31	0	
2015	03	22	0	
2015	04	9	0	
2015	05	13	0	
2015	06	13	0	
2015	07	18	0	
2015	08	23	0	
2015	09	16	0	
2015	10	12	0	
2015	11	23	0	
Total (Provisional) for 2014-15 Season		1177	0	

Influenza Related ICU Admissions by Age Group, 2013-2015



Laboratory Surveillance

• For this reporting week there were **430** influenza specimens tested by WHO/NREVSS collaborating Illinois laboratories and **13** influenza specimens tested by Illinois Department of Public Health Laboratories for a **total of 443** specimens. **99** specimens tested positive for Influenza.

Year	Week	A (H1)	2009(A)H1 N1	А (Н3)	A (Unable to subtype)	A (Sub typing not performed)	В	Total # Tested	Total # Positive	% Positive
2014	40-49	0	1	79	0	806	18	4216	904	21.4%
2014	50	0	0	137	0	154	3	1039	294	28.3%
2014	51	0	0	119	0	277	10	1387	406	29.3%
2014	52	0	0	91	0	85	4	656	180	27.4%
2014	53	0	0	42	0	72	5	673	119	17.7%
2015	01	0	0	33	0	40	5	577	78	13.5%
2015	02	0	1	61	0	24	24 11		97	15.8%
2015	03	0	0	14	0	9	11	434	34	7.8%
2015	04	0	0	2	0	11	9	433	22	5.1%
2015	05	0	0	5	0	4	11	388	20	5.2%
2015	06	0	0	1	0	2	19	414	22	5.3%
2015	07	0	0	3	0	5	29	433	37	8.6%
2015	08	0	0	2	0	3	32	424	37	8.7%
2015	09	0	0	1	0	3	62	484	66	13.6%
2015	10	0	1	1	0	3 117		578	122	21.1%
2015	11	0	0	0	0	0	99	443	99	22.3%
Seaso	n Totals	0	3	591	0	1498	445	13192	2537	19.2%

Influenza Outbreaks Reported in Long-Term Facilities (LTC) and Correctional Facilities

There was **one** outbreak reported for this reporting week.

Region	2014-2015 Influenza Season -Number of outbreaks (%)
Rockford (1)	22 (11.5)
Peoria (2),	31 (16.2)
Edwardsville (4),	36 (18.8)
Marion (5),	15 (7.9)
Champaign (6),	13 (6.8)
West Chicago (7)	54 (28.3)
Chicago/Cook (8)	20 (10.5)
Total	191

Viral Resistance:

Antiviral Resistance: Testing of 2009 H1N1, influenza A (H3N2), and influenza B virus isolates for resistance to neuraminidase inhibitors (oseltamivir and zanamivir) is performed at CDC using a functional assay. Additional 2009 H1N1 and influenza A (H3N2) clinical samples are tested for mutations of the virus known to confer oseltamivir resistance. The data summarized below combine the results of both testing methods. These samples are routinely obtained for surveillance purposes rather than for diagnostic testing of patients suspected to be infected with antiviral-resistant virus.

High levels of resistance to the adamantanes (amantadine and rimantadine) persist among A(H1N1)pdm09 and A(H3N2) viruses (the adamantanes are not effective against influenza B viruses). Therefore, data from adamantane resistance testing are not presented below.


Neuraminidase Inhibitor Resistance Testing Results on Samples Collected Since October 1, 2014

	Ose	Itamivir	Zar	namivir	Peramivir		
	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)	Virus Samples tested (n)	Resistant Viruses, Number (%)	
Influenza A(H1N1)pmd09	38	1 (2.6)	33	0 (0.0)	38	1 (2.6)	
Influenza A (H3N2)	2,443	0 (0.0)	2,443	0 (0.0)	1,407	0 (0.0)	
Influenza B	338	0 (0.0)	338	0 (0.0)	338	0 (0.0)	


In the United States, the vast majority of recently circulating influenza viruses have been susceptible to the neuraminidase inhibitor antiviral medications, oseltamivir, zanamivir, and peramivir; rare sporadic instances of oseltamivir-resistant A(H1N1)pdm09 and A(H3N2) viruses have been detected worldwide. Antiviral treatment is recommended as early as possible for patients with confirmed or suspected influenza who have severe, complicated, or progressive illness; who require hospitalization; or who are at high risk for serious influenza-related complications. Additional information on recommendations for treatment and chemoprophylaxis of influenza virus infection with antiviral agents is available at http://www.cdc.gov/flu/antivirals/index.htm.

Weekly Viral Subtype

Influenza Isolates from Illinois Reported by WHO/NREVSS Collaborating Laboratories, 2014-2015 Influenza Season.



IDPH Infectious Diseases Regional Map



Resources

- Centers for Disease Control and Prevention Influenza Website:
 - o http://www.cdc.gov/flu/
- Immunization Action Coalition Website: http://immunize.org/
- IDPH Seasonal Influenza Website: http://www.idph.state.il.us/flu/surveillance.htm
- National Respiratory and Enteric Virus Surveillance System (NREVSS), CDC website: https://wwwn.cdc.gov/nrevss/account/export.aspx
- St Louis Children's Hospital Weekly Virus/Microbiology Update: http://slchlabtestguide.bjc.org/Default.aspx?url=63e0653d-fe31-466f-9228-d4de90fa7424