


Illinois Tuberculosis (TB) Statistics

Slide Show

TB Cases Illinois, 1980-2003




TB Cases and Rates, 1999-2003

Year	Cases		Rate*	
	U.S.	Illinois	U.S.	Illinois
1999	17,531	825	6.4	6.8
2000	16,377	743	5.8	6.0
2001	15,989	707	5.6	5.7
2002	15,075	680	5.2	5.4
2003	14,871	633	5.1	5.0

*Cases per 100,000 population

TB Rates,* 1995-2003




■ US ■ Illinois


*per 100,000 population

Source: Population data - Illinois Department of Public Health, Center for Health Statistics; and U.S. Bureau of the Census


Reported Tuberculosis Cases Illinois, 1999-2003




Number of TB Cases, U.S.-born vs. Foreign-born Persons, Illinois, 1999-2003




Percentage of Foreign-born vs. U.S.-born TB Cases, Illinois, 1995-2003




Country of Birth for Foreign-born Persons Reported with TB in Illinois, 2003



Distribution of Foreign-born TB Cases by Jurisdiction




Black, Non-Hispanic Cases* By Jurisdiction




*Includes Foreign-born


TB Cases by Ethnicity/Race Illinois, 1999-2003




TB Cases by Age Group Illinois, 1999-2003



TB Rates by Age Group Illinois, 1999-2003




Primary Anti-TB Drug Resistance Illinois, 1999-2003




*MDR, or multi-drug resistant, TB is defined as resistance to at least isoniazid and rifampin.

TB Cases Reported HIV Positive* 1999-2003



* Medical documentation or patient history


HIV Positive and Percentage of Reporting in Cases ages 25-44, Illinois, 1999-2003



■ HIV Positive **—● Percentage Valid Response***


* Valid response = positive, negative or indeterminate

HIV Valid* Reporting Ages 25-44, Illinois, 1999-2003



* Valid Response includes positive, negative and indeterminate.

Completion of TB Therapy Illinois, 1995-2002




* National Objective-Healthy People 2010 target: Completed in one year or less

** Preliminary data

Note: Persons with initial isolate resistant to rifampin and children younger than 15 years of age with meningeal, bone or joint, or military disease are excluded.

Mode of Treatment Administration, Reported Cases, Illinois, 1993-2002



Directly observed therapy (DOT); Self-administered therapy (SA)