

Compendium of Animal Rabies Prevention and Control, 2002*
National Association of State Public Health Veterinarians, Inc. (NASPHV)

The purpose of this Compendium is to provide rabies information to veterinarians, public health officials, and others concerned with rabies prevention and control. These recommendations serve as the basis for animal rabies-control programs throughout the United States and facilitate standardization of procedures among jurisdictions, thereby contributing to an effective national rabies-control program. This document is reviewed annually and revised as necessary. Vaccination procedure recommendations are contained in Part I; all animal rabies vaccines licensed by the United States Department of Agriculture (USDA) and marketed in the United States are listed in Part II; Part III details the principles of rabies control.

Part I: Recommendations for Parenteral Vaccination Procedures

- A. VACCINE ADMINISTRATION:** All animal rabies vaccines should be restricted to use by, or under the direct supervision of, a veterinarian.
- B. VACCINE SELECTION:** Part II lists all vaccines licensed by USDA and marketed in the United States at the time of publication. New vaccine approvals or changes in label specifications made subsequent to publication should be considered as part of this list. Any of the listed vaccines can be used for revaccination, even if the product is not the same brand as previously administered vaccines. Vaccines used in state and local rabies control programs should have a 3-year duration of immunity. This constitutes the most effective method of increasing the proportion of immunized dogs and cats in any population.
- C. ROUTE OF INOCULATION:** All vaccines must be administered in accordance with the specifications of the product label or package insert. Adverse reactions and vaccine failures should be reported to USDA, Animal and Plant Health Inspection Service, Center for Veterinary Biologics at (800) 752-6255 or by e-mail at CVB@usda.gov.
- D. WILDLIFE AND HYBRID ANIMAL VACCINATION:** The efficacy of parenteral rabies vaccination of wildlife and hybrids (the offspring of wild animals crossbred to domestic dogs and cats) has not been established, and no such vaccine is licensed for these animals. Zoos or research institutions may establish vaccination programs which attempt to protect valuable animals, but these should not replace appropriate public health activities that protect humans.
- E. ACCIDENTAL HUMAN EXPOSURE TO VACCINE:** Human exposure to parenteral animal rabies vaccines listed in Part II does not constitute a risk for rabies infection. However, human exposure to vaccinia-vectored oral rabies vaccines should be reported to state health officials.¹
- F. IDENTIFICATION OF VACCINATED ANIMALS:** Agencies and veterinarians may adopt the standard tag system to aid in the administration of animal rabies control procedures.

1. RABIES TAGS

CALENDAR YEAR	COLOR	SHAPE
2002	Orange	Oval
2003	Green	Bell

- 2. RABIES CERTIFICATE:** All agencies and veterinarians should use the NASPHV Form #51, "Rabies Vaccination Certificate," which can be obtained from vaccine manufacturers. Computer-generated forms containing the same information are acceptable.

THE NASPHV COMMITTEE

Suzanne R. Jenkins, VMD, MPH, Chair
Michael Auslander, DVM, MSPH
Lisa Conti, DVM, MPH
William B. Johnston, DVM
Mira J. Leslie, DVM
Faye E. Sorhage, VMD, MPH

***Address all correspondence to:**

Suzanne R. Jenkins, VMD, MPH
Virginia Department of Health
Office of Epidemiology
Post Office Box 2448, Room 113
Richmond, VA 23218

CONSULTANTS TO THE COMMITTEE

James E. Childs, ScD; Centers for Disease Control and Prevention (CDC)
Mary Currier, MD, MPH; CSTE
Nancy Frank, DVM, MPH; AVMA Council on Public Health and Regulatory Veterinary Medicine
Donna M. Gatewood, DVM, MS; Animal and Plant Health Inspection Service, USDA
Carolyn L. Schumacher, DVM, PhD; Animal Health Institute
Charles E. Rupprecht, VMD, PhD; CDC
Charles V. Trimarchi, MS; New York State Health Department

ENDORSED BY:

American Veterinary Medical Association (AVMA)
Council of State and Territorial Epidemiologists (CSTE)

Part II: Rabies Vaccines Licensed and Marketed in the U.S., 2002

Product Name	Produced by	Marketed by	For Use In	Dosage	Age at Primary Vaccination ¹	Booster Recommended	Route of Inoculation
A) MONOVALENT (Inactivated)							
TRIMUNE	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats	1 ml 1 ml	3 months ² 3 months	1 year later & triennially 1 year later & triennially	IM ³ IM
ANNUMUNE	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	IM IM
DEFENSOR 1	Pfizer, Incorporated License No. 189	Pfizer, Incorporated	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	IM or SC ⁴ SC
DEFENSOR 3	Pfizer, Incorporated License No. 189	Pfizer, Incorporated	Dogs Cats Sheep Cattle	1 ml 1 ml 2 ml 2 ml	3 months 3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually Annually	IM or SC SC IM IM
RABDOMUN	Pfizer, Incorporated License No. 189	Schering-Plough	Dogs Cats Sheep Cattle	1 ml 1 ml 2 ml 2 ml	3 months 3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually Annually	IM or SC SC IM IM
RABDOMUN 1	Pfizer, Incorporated License No. 189	Schering-Plough	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	IM or SC SC
RABVAC 1	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	IM or SC IM or SC
RABVAC 3	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats Horses	1 ml 1 ml 2 ml	3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually	IM or SC IM or SC IM
PRORAB-1	Intervet, Incorporated License No. 286	Intervet, Incorporated	Dogs Cats Sheep	1 ml 1 ml 2 ml	3 months 3 months 3 months	Annually Annually Annually	IM or SC IM or SC IM
PRORAB-3F	Intervet, Incorporated License No. 286	Intervet, Incorporated	Cats	1 ml	3 months	1 year later & triennially	IM or SC
IMRAB 3	Merial, Incorporated License No. 298	Merial, Incorporated	Dogs Cats Sheep Cattle Horses Ferrets	1 ml 1 ml 2 ml 2 ml 2 ml 1 ml	3 months 3 months 3 months 3 months 3 months 3 months	1 year later & triennially 1 year later & triennially 1 year later & triennially Annually Annually Annually	IM or SC IM or SC IM or SC IM or SC IM or SC SC
IMRAB Bovine Plus	Merial, Incorporated License No. 298	Merial, Incorporated	Cattle Horses Sheep	2 ml 2 ml 2 ml	3 months 3 months 3 months	Annually Annually 1 year later & triennially	IM or SC IM or SC IM or SC
IMRAB 1	Merial, Incorporated License No. 298	Merial, Incorporated	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	SC SC
B) MONOVALENT (Rabies glycoprotein, live canary pox vector)							
PUREVAX Feline Rabies	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
C) COMBINATION (Inactivated rabies)							
ECLIPSE 3 + FeLV/R	Fort Dodge Animal Health License No. 112	Schering-Plough	Cats	1 ml	3 months	Annually	IM or SC
ECLIPSE 4 + FeLV/R	Fort Dodge Animal Health License No. 112	Schering-Plough	Cats	1 ml	3 months	Annually	IM or SC
Fel-O-Guard 3 + FeLV/R	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Cats	1 ml	3 months	Annually	IM or SC
Fel-O-Guard 4 + FeLV/R	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Cats	1 ml	3 months	Annually	IM or SC
IMRAB 3 + Feline 3	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1 ml	3 months	1 year later & triennially	SC
IMRAB 3 + Feline 4	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1 ml	3 months	1 year later & triennially	SC
MYSTIQUE II POTOMAVAC+	Bayer Corporation License No. 52	Bayer Corporation	Horses	1 ml	3 months	Annually	IM
Equine POTOMAVAC + IMRAB	Merial, Incorporated License No. 298	Merial, Incorporated	Horses	1 ml	3 months	Annually	IM
D) COMBINATION (Rabies glycoprotein, live canary pox vector)							
PUREVAX Feline 3/Rabies	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
PUREVAX Feline 3/ Rabies + LEUCAT	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
PUREVAX Feline 4/ Rabies + LEUCAT	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC

Product Name	Produced by	Marketed by	For Use In	Dosage	Age at Primary Vaccination ¹	Booster Recommended	Route of Inoculation
D) COMBINATION (Rabies glycoprotein, live canary pox vector) continued							
PUREVAX Feline/ Rabies + LEUCAT	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
E) ORAL (Rabies glycoprotein, live vaccinia vector) - RESTRICTED TO USE IN STATE AND FEDERAL RABIES CONTROL PROGRAMS							
RABORAL V-RG	Merial, Incorporated License No. 298	Merial, Incorporated	Raccoons	N/A	N/A	As determined by local authorities	Oral

1 Minimum age (or older) and revaccinated one year later.

2 A month = 28 days

3 Intramuscularly

4 Subcutaneously

Part III: Rabies Control

A. PRINCIPLES OF RABIES CONTROL

- RABIES EXPOSURE:** Rabies is transmitted only when the virus is introduced into bite wounds, open cuts in skin, or onto mucous membranes.
- HUMAN RABIES PREVENTION:** Rabies in humans can be prevented either by eliminating exposures to rabid animals or by providing exposed persons with prompt local treatment of wounds combined with human rabies immune globulin and vaccine. The rationale for recommending preexposure and postexposure rabies prophylaxis and details of their administration can be found in the current recommendations of the Advisory Committee on Immunization Practices (ACIP)². These recommendations, along with information concerning the current local and regional status of animal rabies and the availability of human rabies biologics, are available from state health departments.
- DOMESTIC ANIMALS:** Local governments should initiate and maintain effective programs to ensure vaccination of all dogs, cats, and ferrets and to remove strays and unwanted animals. Such procedures in the United States have reduced laboratory-confirmed cases of rabies in dogs from 6,949 in 1947 to 114 in 2000. Because more rabies cases are reported annually involving cats (249 in 2000) than dogs, vaccination of cats should be required. The recommended vaccination procedures and the licensed animal vaccines are specified in Parts I and II of the Compendium.
- RABIES IN WILDLIFE:** The control of rabies among wildlife reservoirs is difficult. Vaccination of free-ranging wildlife or selective population reduction might be useful in some situations, but the success of such procedures depends on the circumstances surrounding each rabies outbreak. (See Part C. Control Methods in Wildlife.) Because of the risk of rabies in wild animals (especially raccoons, skunks, coyotes, foxes, and bats), the AVMA, the NASPHV, and the CSTE strongly recommend the enactment of state laws prohibiting their importation, distribution, and relocation.
- RABIES SEROLOGY:** Evidence of circulating rabies virus neutralizing antibodies should not be used as a substitute for current vaccination in managing rabies exposures or determining the need for booster vaccinations.

B. CONTROL METHODS IN DOMESTIC AND CONFINED ANIMALS

1. PREEXPOSURE VACCINATION AND MANAGEMENT

Parenteral animal rabies vaccines should be administered only by, or under the direct supervision of, a veterinarian. This ensures that a qualified and responsible person can be held accountable to assure the public that the animal has been properly vaccinated. Within twenty-eight (28) days after primary vaccination, a peak rabies antibody titer is reached and the animal can be considered immunized.

An animal is currently vaccinated and is considered immunized if the primary vaccination was administered at least 28 days previously and vaccinations have been administered in accordance with this Compendium. Regardless of the age of the animal at initial vaccination, a booster vaccination should be administered 1 year later. (See Parts I and II for vaccines and procedures) Because a rapid anamnestic response is expected, an animal is considered currently vaccinated immediately after a booster vaccination.

(a) DOGS, CATS, AND FERRETS

All dogs, cats, and ferrets should be vaccinated against rabies and revaccinated in accordance with Part II of this Compendium. If a previously vaccinated animal is overdue for a booster, it should be revaccinated with a single dose of vaccine. Immediately following the booster, the animal is considered currently vaccinated and should be placed on an annual or triennial schedule depending on the type of vaccine used.

(b) LIVESTOCK

Consideration should be given to vaccinating livestock that are particularly valuable or that might have frequent contact with humans. Horses traveling interstate should be currently vaccinated against rabies.

(c) CONFINED ANIMALS

(1) WILD

No parenteral rabies vaccine is licensed for use in wild animals. Wild animals or hybrids should not be kept as pets.

(2) MAINTAINED IN EXHIBITS AND IN ZOOLOGICAL PARKS

Captive animals that are not completely excluded from all contact with rabies vectors can become infected. Moreover, wild animals might be incubating rabies when initially captured; therefore, wild-caught animals susceptible to rabies should be quarantined for a minimum of 6 months before being exhibited. Employees who work with animals at such facilities should receive preexposure rabies vaccination. The use of pre- or postexposure rabies vaccinations for employees who work with animals at such facilities might reduce the need for euthanasia of captive animals. Carnivores and bats should be housed in a manner that precludes direct contact with the public.

2. STRAY ANIMALS

Stray dogs, cats, and ferrets should be removed from the community. Local health departments and animal control officials can enforce the removal of strays more effectively if owned animals are confined or kept on leash. Strays should be impounded for at least 3 days to determine if human exposure has occurred and to give owners sufficient time to reclaim animals.

3. IMPORTATION AND INTERSTATE MOVEMENT OF ANIMALS

(a) INTERNATIONAL

CDC regulates the importation of dogs and cats into the United States. Imported dogs must satisfy rabies vaccination requirements (42 CFR, Part 71.51[c], www.cdc.gov/ncidod/dq/lawsand/htm). The appropriate health official of the state of destination should be notified within 72 hours of the arrival into his or her jurisdiction of any imported dog required to be placed in confinement under the CDC regulation. Failure to comply with these requirements should be promptly reported to the Division of Quarantine, CDC, (404) 639-8107.

CDC regulations alone are insufficient to prevent the introduction of rabid animals into the country. All imported dogs and cats are subject to state and local laws governing rabies and should be currently vaccinated against rabies in accordance with the Compendium. Failure to comply with state or local requirements should be referred to the appropriate state or local official.

(b) INTERSTATE

Before interstate movement, dogs, cats, and ferrets should be currently vaccinated against rabies in accordance with the Compendium's recommendations (See Part III, B.1. Preexposure Vaccination and Management). Animals in transit should be accompanied by a currently valid NASPHV Form #51, Rabies Vaccination Certificate. When an interstate health certificate or certificate of veterinary inspection is required, it should contain the same rabies vaccination information as Form #51.

4. ADJUNCT PROCEDURES

Methods or procedures which enhance rabies control include the following:

- (a) **IDENTIFICATION.** Dogs, cats and ferrets should be identified (e.g., metal or plastic tags, microchips, etc.) to allow for verification of rabies vaccination status.
- (b) **LICENSURE.** Registration or licensure of all dogs, cats, and ferrets may be used to aid in rabies control. A fee is frequently charged for such licensure and revenues collected are used to maintain rabies- or animal-control programs. Vaccination is an essential prerequisite to licensure.
- (c) **CANVASSING OF AREA.** House-to-house canvassing by animal control personnel facilitates enforcement of vaccination and licensure requirements.
- (d) **CITATIONS.** Citations are legal summonses issued to owners for violations, including the failure to vaccinate or license their animals. The authority for officers to issue citations should be an integral part of each animal-control program.
- (e) **ANIMAL CONTROL.** All communities should incorporate stray animal control, leash laws, and training of personnel in their programs.

5. POSTEXPOSURE MANAGEMENT

ANY ANIMAL POTENTIALLY EXPOSED TO RABIES VIRUS (See Part III, A. 1. Rabies Exposure) BY A WILD, CARNIVOROUS MAMMAL OR A BAT THAT IS NOT AVAILABLE FOR TESTING SHOULD BE REGARDED AS HAVING BEEN EXPOSED TO RABIES.

(a) DOGS, CATS, AND FERRETS

Unvaccinated dogs, cats, and ferrets exposed to a rabid animal should be euthanized immediately. If the owner is unwilling to have this done, the animal should be placed in strict isolation for 6 months and vaccinated 1 month before being released. Animals with expired vaccinations need to be evaluated on a case-by-case basis. Dogs, cats, and ferrets that are currently vaccinated should be revaccinated immediately, kept under the owner's control, and observed for 45 days.

(b) LIVESTOCK

All species of livestock are susceptible to rabies; cattle and horses are among the most frequently infected. Livestock exposed to a rabid animal and currently vaccinated with a vaccine approved by USDA for that species should be revaccinated immediately and observed for 45 days. Unvaccinated livestock should be slaughtered immediately. If the owner is unwilling to have this done, the animal should be kept under close observation for 6 months.

The following are recommendations for owners of unvaccinated livestock exposed to rabid animals:

- (1) If the animal is slaughtered within 7 days of being bitten, its tissues may be eaten without risk of infection, provided that liberal portions of the exposed area are discarded. Federal meat inspectors must reject for slaughter any animal known to have been exposed to rabies within 8 months.
- (2) Neither tissues nor milk from a rabid animal should be used for human or animal consumption. Pasteurization temperatures will inactivate rabies virus, therefore, drinking pasteurized milk or eating cooked meat does not constitute a rabies exposure.
- (3) Having more than one rabid animal in a herd or having herbivore-to-herbivore transmission is uncommon; therefore, restricting the rest of the herd if a single animal has been exposed to or infected by rabies might not be necessary.

(c) **OTHER ANIMALS**

Other mammals bitten by a rabid animal should be euthanized immediately. Animals maintained in USDA licensed research facilities or accredited zoological parks should be evaluated on a case-by-case basis.

6. MANAGEMENT OF ANIMALS THAT BITE HUMANS

- (a) A healthy dog, cat, or ferret that bites a person should be confined and observed daily for 10 days; administration of rabies vaccine is not recommended during the observation period. Such animals should be evaluated by a veterinarian at the first sign of illness during confinement. Any illness in the animal should be reported immediately to the local health department. If signs suggestive of rabies develop, the animal should be euthanized and the head shipped for testing as described in (c) below. Any stray or unwanted dog, cat, or ferret that bites a person may be euthanized immediately and the head submitted for rabies examination.
- (b) Other biting animals which might have exposed a person to rabies should be reported immediately to the local health department. Prior vaccination of an animal may not preclude the necessity for euthanasia and testing if the period of virus shedding is unknown for that species. Management of animals other than dogs, cats, and ferrets depends on the species, the circumstances of the bite, the epidemiology of rabies in the area, and the biting animal's history, current health status, and potential for exposure to rabies.
- (c) Rabies testing should be done by a qualified laboratory, designated by the local or state health department. Euthanasia³ should be accomplished in such a way as to maintain the integrity of the brain so that the laboratory can recognize the anatomical parts. Except in the case of very small animals, such as bats, only the head or brain (including brain stem) should be submitted to the laboratory. Any animal or animal part being submitted for testing should be kept under refrigeration (not frozen or chemically fixed) during storage and shipping.

C. CONTROL METHODS IN WILDLIFE

The public should be warned not to handle wildlife. Wild mammals and hybrids that bite or otherwise expose persons, pets or livestock should be considered for euthanasia and rabies examination. A person bitten by any wild mammal should immediately report the incident to a physician who can evaluate the need for antirabies treatment (See current rabies prophylaxis recommendations of the ACIP²). State regulated wildlife rehabilitators may play a role in a comprehensive rabies control program. Minimum standards for persons who rehabilitate wild mammals should include rabies vaccination, appropriate training and continuing education. Translocation of infected wildlife has contributed to the spread of rabies; therefore, the translocation of known terrestrial rabies reservoir species should be prohibited.

1. TERRESTRIAL MAMMALS

The use of licensed oral vaccines for the mass vaccination of free-ranging wildlife should be considered in selected situations, with the approval of the state agency responsible for animal rabies control. The distribution of oral rabies vaccine should be based on scientific assessments of the target species and followed by timely and appropriate analysis of surveillance data; such results should be provided to all stakeholders. Continuous and persistent government-funded programs for trapping or poisoning wildlife are not cost effective in reducing wildlife rabies reservoirs on a statewide basis. However, limited control in high-contact areas (e.g., picnic grounds, camps, suburban areas) may be indicated for the removal of selected high-risk species of wildlife. State agriculture, public health and wildlife agencies should be consulted for planning, coordination and evaluation of vaccination or population-reduction programs.

2. BATS

Indigenous rabid bats have been reported from every state except Hawaii, and have caused rabies in at least 33 humans in the United States. Bats should be excluded from houses and adjacent structures to prevent direct association with humans. Such structures should then be made bat-proof by sealing entrances used by bats. Controlling rabies in bats by programs designed to reduce bat populations is neither feasible nor desirable.

REFERENCE

1. Rupprecht CE, et. al. Brief Report: Human Infection Due to Recombinant Vaccinia-Rabies Glycoprotein virus. *N Engl J Med* 2001; 345:8,582-586.
2. Centers for Disease Control and Prevention: Human rabies prevention--United States, 1999. recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1999;48(No. RR-1).
3. 2000 Report of the AVMA Panel on Euthanasia. *JAVMA* 2001; 218:5,669-696.