



Rod R. Blagojevich, Governor
 Eric E. Whitaker, M.D., M.P.H., Director

525-535 West Jefferson Street • Springfield, Illinois 62761-0001 • www.idph.state.il.us

MEMORANDUM

TO: Local Health Departments, Regional Offices of Illinois Department of Public Health, Illinois Department of Public Health Rabies Laboratories, Illinois Department of Agriculture Rabies Laboratories, Illinois Department of Agriculture Animal Welfare, Illinois Department of Natural Resources

FROM: Connie Austin, D.V.M., M.P.H., Ph.D.
 State Public Health Veterinarian

DATE: March 2, 2004

SUBJECT: **Animal Rabies Surveillance for 2003**

In 2003, a total of 3776 animals were submitted for rabies testing to the Illinois Department of Agriculture (IDA) and the Illinois Department of Public Health (IDPH) diagnostic laboratories. Fifty-six were not suitable for testing. Of the total tested, 24 were fluorescent antibody-positive for rabies and all were bats. Animals testing positive for rabies are listed in Table 1.

Table 1. Rabies test positive animals by species and year in Illinois

Species	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
Bat	16	12	23	16	16	10	22	24	31	24
Cat	0	0	1	0	0	0	0	0	0	0
Cattle	0	1	0	0	0	0	0	0	0	0
Dog	1	0	0	0	0	0	0	0	0	0
Fox	0	0	0	0	0	0	0	0	0	0
Skunk	4	2	1	4	1	0	0	0	0	0
Raccoon	0	0	0	0	0	0	0	0	0	0
Total	21	15	25	20	17	10	22	24	31	24

BATS

In 2003 bats were the only species identified with rabies in Illinois. For the past four years no terrestrial animals were diagnosed with rabies. This year, 870 bats have been tested for rabies. Of those tested, approximately 846 were determined negative for the virus, 24 were confirmed positive. An additional 24 bats were submitted for testing but results were inconclusive. The percent of bats tested which were positive for rabies in 2003 was 2.7 percent (See Table 2).


Table 2. Bat rabies testing results in Illinois for 2003.

Negative	Positive	Inconclusive	Total Tested	% Positive
846	24	24	870	2.7

Bats testing positive for rabies were found in the following counties: two each in Dupage, Jackson, Williamson and Winnebago and one each in Boone, Champaign, Cook, DeWitt, Edwards, Kankakee, Kendall, Knox, Lake, Madison, McHenry, Perry, Rock Island, Sangamon, Wayne and White. The majority (57 percent) of testing occurred from May to August when bats are most active and likely to come in contact with humans or pets, as demonstrated in Figure 1.

Two of the positive bats were speciated, one was a red bat and one was a small brown bat.

Figure 1. Number of all bats and positive bats tested in Illinois by month for 2003.



SPECIMEN SUBMISSION

Rabies specimens should be submitted under the following conditions:

1. When a domestic animal or a person is bitten by an animal.
2. When an animal exhibits the neurological symptoms of rabies.
3. Routine submission of skunks for rabies surveillance.

All brain or head specimens should be submitted immediately for testing and not batched with other specimens. If someone has been exposed to a possibly rabid animal, the animal should be sent immediately for testing. If the animal should test positive, the person would need to be started immediately on rabies treatment. A delay in obtaining results could result in a delay in starting needed treatment and result in a poor outcome for the exposed individual.

SKUNKS

In 2003 no skunks tested positive for rabies, yet the number of skunks tested continues to be low. Of the 3776 animals tested, only 73 (1.9 percent) were skunks. The following counties have submitted skunks for rabies testing: Champaign, Coles, Cook, DuPage, Jackson, Kane, Lake, Livingston, Madison, McHenry, McLean, Monroe, Morgan, Sangamon, St. Clair, Stephenson, Warren, Whiteside and Will. Forty of the 73 skunks (55 percent) were submitted from Cook County.

To maintain adequate surveillance in the state, testing of the skunk must be maintained, since it is the main terrestrial animal reservoir for the rabies virus in Illinois. Negative testing of wild mammals, especially skunks, in counties is one factor used to determine whether rabies post-exposure treatment is recommended for stray dog and cat bites.

County animal control officers are encouraged to submit skunks for testing if

- they have exposed a person or a domestic animal or
- they have signs suggestive of rabies, including
 - lack of fear of humans
 - aggressive behavior
 - salivation
 - staggering
 - paralysis and muscle tremors
 - found in places unusual for a skunk

Local animal control and local public health agencies are also asked to submit skunks which have not exposed persons or animals. This is especially important if no skunks have been tested for rabies in their jurisdiction. Local agencies are encouraged to contact nuisance wildlife trappers and others in their jurisdictions who may have freshly killed skunks that could be tested and arrange for transportation and testing by one of the IDOA or IDPH laboratories. Rabies testing is free at the IDOA and IDPH laboratory for this enhanced skunk surveillance program. For other animal rabies testing, the IDOA laboratories charge \$12 per animal. All rabies testing is free at the IDPH laboratories.

Attached is the Compendium of Animal Rabies Prevention and Control, 2004. Please share this document and the Compendium with your local animal control partners. The Compendium provides recommendations from the Compendium committee on animal control issues related to rabies and provides a list of licensed rabies vaccines for animals. When evaluating what to do with animals exposed to rabies or animals that have bitten someone in Illinois, please follow the Illinois Animal Control Act administered by the Illinois Department of Agriculture.

Compendium of Animal Rabies Prevention and Control, 2004*

National Association of State Public Health Veterinarians, Inc. (NASPHV)

Rabies is a fatal viral zoonosis and a serious public health problem¹. The purpose of this Compendium is to provide information to veterinarians, public health officials, and others concerned with rabies prevention and control. These recommendations serve as the basis for animal rabies-control programs throughout the United States and facilitate standardization of procedures among jurisdictions, thereby contributing to an effective national rabies-control program. This document is reviewed annually and revised as necessary. Parenteral vaccination procedure recommendations are contained in Part I; Part II details the principles of rabies control; all animal rabies vaccines licensed by the United States Department of Agriculture (USDA) and marketed in the United States are listed in Part III.

THE NASPHV COMMITTEE

Suzanne R. Jenkins, VMD, MPH, Chair
Michael Auslander, DVM, MSPH
Lisa Conti, DVM, MPH
Mira J. Leslie, DVM, MPH
Faye E. Sorhage, VMD, MPH
Ben Sun, DVM, MPVM

*Address all correspondence to:

Suzanne R. Jenkins, VMD, MPH
Virginia Department of Health
Office of Epidemiology
Post Office Box 2448, Room 113
Richmond, VA 23218

CONSULTANTS TO THE COMMITTEE

Mary Currier, MD, MPH; CSTE
Donna M. Gatewood, DVM, MS; Center for Veterinary
Biologics, USDA
Dan Knox, DVM; NACA
Charles E. Rupprecht, VMD, MS, PhD; CDC
John Schiltz, DVM; AVMA
Carolyn L. Schumacher, DVM, PhD; Animal Health Institute
Charles V. Trimarchi, MS; New York State Health Department

ENDORSED BY:

American Veterinary Medical Association (AVMA)
Centers for Disease Control and Prevention (CDC)
Council of State and Territorial Epidemiologists (CSTE)
National Animal Control Association (NACA)

Part I: Recommendations for Parenteral Vaccination Procedures

- A. VACCINE ADMINISTRATION:** All animal rabies vaccines should be restricted to use by, or under the direct supervision of a veterinarian², except as recommended in Part II.B.1. All vaccines must be administered in accordance with the specifications of the product label or package insert.
- B. VACCINE SELECTION:** Part III lists all vaccines licensed by USDA and marketed in the United States at the time of publication. New vaccine approvals or changes in label specifications made subsequent to publication should be considered as part of this list. Any of the listed vaccines can be used for revaccination, even if the product is not the same brand as previously administered vaccines. Vaccines used in state and local rabies control programs should have a 3-year duration of immunity. This constitutes the most effective method of increasing the proportion of immunized dogs and cats in any population³. There are no laboratory or epidemiologic data to support the annual or biennial administration of 3-year vaccines following the initial series.
- C. ADVERSE EVENTS:** Currently, there is no epidemiologic association between a particular licensed vaccine product and adverse events including vaccine failure^{4,5}. Adverse events should be reported to the vaccine manufacturer and to the USDA, Animal and Plant Health Inspection Service, Center for Veterinary Biologics (at <http://www.aphis.usda.gov/vs/cvb/ic/adverseeventreport.htm>, by telephone at 800-752-6255, or by e-mail to CVB@usda.gov).
- D. WILDLIFE AND HYBRID ANIMAL VACCINATION:** The efficacy of parenteral rabies vaccination of wildlife and hybrids (the offspring of wild animals crossbred to domestic animals) has not been established, and no such vaccine is licensed for these animals. Zoos or research institutions may establish vaccination programs, which attempt to protect valuable animals, but these should not replace appropriate public health activities that protect humans.
- E. ACCIDENTAL HUMAN EXPOSURE TO VACCINE:** Human exposure to parenteral animal rabies vaccines listed in Part III does not constitute a risk for rabies infection. However, human exposure to vaccinia-vectored oral rabies vaccines should be reported to state health officials⁶.
- F. RABIES CERTIFICATE:** All agencies and veterinarians should use the NASPHV Form #51, "Rabies Vaccination Certificate," which can be obtained from vaccine manufacturers. The form must be fully completed and signed by the administering or supervising veterinarian. This form can also be found on the CDC website (www.cdc.gov/ncidod/dvrd/rabies/professional/professi.htm). Computer-generated forms containing the same information are acceptable.

Part II: Rabies Control

A. PRINCIPLES OF RABIES CONTROL

- 1. RABIES EXPOSURE:** Rabies is transmitted only when the virus is introduced into bite wounds, open cuts in skin, or onto mucous membranes⁷.
- 2. HUMAN RABIES PREVENTION:** Rabies in humans can be prevented either by eliminating exposures to rabid animals or by providing exposed persons with prompt local treatment of wounds combined with the administration of human rabies immune globulin and vaccine. The rationale for recommending preexposure and postexposure rabies prophylaxis and details of their administration can be found in the current recommendations of the Advisory Committee on Immunization Practices (ACIP)⁷. These recommendations, along with information concerning the current local and regional status of animal rabies and the availability of human rabies biologics, are available from state health departments.
- 3. DOMESTIC ANIMALS:** Local governments should initiate and maintain effective programs to ensure vaccination of all dogs, cats, and ferrets and to remove strays and unwanted animals. Such procedures in the United States have reduced laboratory-confirmed cases of rabies in dogs from 6,949 in 1947 to 99 in 2002⁸. Because more rabies cases are reported annually involving cats (299 in 2002) than dogs, vaccination of cats should be required. Animal shelters and animal control authorities should establish policies to ensure that adopted animals are vaccinated against rabies. The recommended vaccination procedures and the licensed animal vaccines are specified in Parts I and III of the Compendium.
- 4. RABIES IN VACCINATED ANIMALS:** Rabies is rare in vaccinated animals. If such an event is suspected it should be reported to state public health officials, the vaccine manufacturer, and to the USDA, Animal and Plant Health Inspection Service, Center for Veterinary Biologics (<http://www.aphis.usda.gov/vs/cbv/ic/adverseeventreport.htm>, by telephone at 800-752-6255, or by e-mail to CVB@usda.gov). The laboratory diagnosis should be confirmed and the virus characterized by a rabies reference laboratory. A thorough epidemiologic investigation should be conducted.
- 5. RABIES IN WILDLIFE:** The control of rabies among wildlife reservoirs is difficult⁹. Vaccination of free-ranging wildlife or selective population reduction might be useful in some situations, but the success of such procedures depends on the circumstances surrounding each rabies outbreak (See Part C. Control Methods in Wildlife). Because of the risk of rabies in wild animals (especially raccoons, skunks, coyotes, foxes, and bats), the AVMA, the NASPHV, and the CSTE strongly recommend the enactment of state laws prohibiting their importation, distribution, and relocation.
- 6. RABIES SURVEILLANCE:** Laboratory-based rabies surveillance is an essential component of rabies control and prevention programs. Accurate and timely information is necessary to: guide human postexposure prophylaxis decisions; determine the management of potentially exposed animals; aid in emerging pathogen discovery; describe the epidemiology of the disease; and assess the need for and effectiveness of oral vaccination programs for wildlife.
- 7. RABIES DIAGNOSIS:** Rabies testing should be done by a qualified laboratory, designated by the local or state health department¹⁰ in accordance with the established national standardized protocol for rabies testing (http://www.cdc.gov/ncidod/dvrd/rabies/Professional/publications/DFA_diagnosis/DFA_protocol-b.htm). Euthanasia¹¹ should be accomplished in such a way as to maintain the integrity of the brain so that the laboratory can recognize the anatomical parts. Except in the case of very small animals, such as bats, only the head or brain (including brain stem) should be submitted to the laboratory. Any animal or animal part being submitted for testing should be kept under refrigeration (not frozen or chemically fixed) during storage and shipping.
- 8. RABIES SEROLOGY:** Some “rabies-free” jurisdictions may require evidence of vaccination and rabies antibodies for importation purposes. Rabies antibody titers are indicative of an animal’s response to vaccine or infection; titers are not indicators of protection. Other immunologic factors also play a role in preventing rabies and our abilities to measure and interpret those other factors are not well developed. Therefore, evidence of circulating rabies virus antibodies should **not** be used as a substitute for current vaccination in managing rabies exposures or determining the need for booster vaccinations¹².

B. CONTROL METHODS IN DOMESTIC AND CONFINED ANIMALS

- 1. PREEXPOSURE VACCINATION AND MANAGEMENT:** Parenteral animal rabies vaccines should be administered only by, or under the direct supervision of a veterinarian. Rabies vaccinations may also be administered under the supervision of a veterinarian to animals held in animal control shelters prior to release. Any veterinarian signing a rabies certificate should assure that the person administering vaccine is identified on the certificate and is appropriately trained in vaccine storage, handling, administration, management of adverse events, etc. This practice ensures that a qualified and responsible person can be held accountable to assure that the animal has been properly vaccinated.

Within twenty-eight (28) days after primary vaccination, a peak rabies antibody titer is reached and the animal can be considered immunized. An animal is currently vaccinated and is considered immunized if the primary vaccination was administered at least 28 days previously and vaccinations have been administered in accordance with this Compendium.

Regardless of the age of the animal at initial vaccination, a booster vaccination should be administered 1 year later (See Parts I and III for vaccines and procedures). There are no laboratory or epidemiologic data to support the annual or biennial administration of 3-year vaccines following the initial series. Because a rapid anamnestic response is expected, an animal is considered currently vaccinated immediately after a booster vaccination.

(a) **DOGS, CATS, AND FERRETS**

All dogs, cats, and ferrets should be vaccinated against rabies and revaccinated in accordance with Part III of this Compendium. If a previously vaccinated animal is overdue for a booster, it should be revaccinated with a single dose of vaccine. Immediately following the booster, the animal is considered currently vaccinated and should be placed on an annual or triennial schedule depending on the type of vaccine used.

(b) **LIVESTOCK**

Consideration should be given to vaccinating livestock that are particularly valuable or that might have frequent contact with humans (e.g., in petting zoos, fairs, and other public exhibitions; see www.avma.org/pubhlth/comp_animals_public_settings.asp). Horses traveling interstate should be currently vaccinated against rabies.

(c) **CONFINED ANIMALS**

(1) **WILD**

No parenteral rabies vaccines are licensed for use in wild animals. Wild animals or hybrids should not be kept as pets¹³⁻¹⁶.

(2) **MAINTAINED IN EXHIBITS AND IN ZOOLOGICAL PARKS**

Captive mammals that are not completely excluded from all contact with rabies vectors can become infected. Moreover, wild animals might be incubating rabies when initially captured; therefore, wild-caught animals susceptible to rabies should be quarantined for a minimum of 6 months before being exhibited. Employees who work with animals at such facilities should receive preexposure rabies vaccination. The use of pre- or postexposure rabies vaccinations for employees who work with animals at such facilities might reduce the need for euthanasia of captive animals. Carnivores and bats should be housed in a manner that precludes direct contact with the public.

2. **STRAY ANIMALS:** Stray dogs, cats, and ferrets should be removed from the community. Local health departments and animal control officials can enforce the removal of strays more effectively if owned animals are confined or kept on leash. Strays should be impounded for at least 3 days to determine if human exposure has occurred and to give owners sufficient time to reclaim animals.

3. **IMPORTATION AND INTERSTATE MOVEMENT OF ANIMALS**

(a) **INTERNATIONAL.** CDC regulates the importation of dogs and cats into the United States. Importers of dogs must comply with rabies vaccination requirements (42 CFR, Part 71.51[c], <http://www.cdc.gov/ncidod/dq/lawsand/htm>) and complete the CDC form 75.37. The appropriate health official of the state of destination should be notified within 72 hours of the arrival into his or her jurisdiction of any imported dog required to be placed in confinement under the CDC regulation. Failure to comply with these requirements should be promptly reported to the Division of Global Migration and Quarantine, CDC, (404) 498-1670.

Federal regulations alone are insufficient to prevent the introduction of rabid animals into the country^{17,18}. All imported dogs and cats are subject to state and local laws governing rabies and should be currently vaccinated against rabies in accordance with this Compendium. Failure to comply with state or local requirements should be referred to the appropriate state or local official.

(b) **INTERSTATE.** Before interstate movement, dogs, cats, ferrets, and horses should be currently vaccinated against rabies in accordance with the Compendium's recommendations (See Part II, B.1. Preexposure Vaccination and Management). Animals in transit should be accompanied by a currently valid NASPHV Form #51, Rabies Vaccination Certificate. When an interstate health certificate or certificate of veterinary inspection is required, it should contain the same rabies vaccination information as Form #51.

- 4. ADJUNCT PROCEDURES:** Methods or procedures which enhance rabies control include the following:
- (a) **IDENTIFICATION.** Dogs, cats, and ferrets should be identified (e.g., metal or plastic tags, microchips, etc.) to allow for verification of rabies vaccination status.
 - (b) **LICENSURE.** Registration or licensure of all dogs, cats, and ferrets may be used to aid in rabies control. A fee is frequently charged for such licensure and revenues collected are used to maintain rabies- or animal-control programs. Vaccination is an essential prerequisite to licensure.
 - (c) **CANVASSING OF AREA.** House-to-house canvassing by animal control officials facilitates enforcement of vaccination and licensure requirements.
 - (d) **CITATIONS.** Citations are legal summonses issued to owners for violations, including the failure to vaccinate or license their animals. The authority for officers to issue citations should be an integral part of each animal-control program.
 - (e) **ANIMAL CONTROL.** All communities should incorporate stray animal control, leash laws, and training of personnel in their programs.
- 5. POSTEXPOSURE MANAGEMENT:** Any animal potentially exposed to rabies virus (See Part II, A. 1. Rabies Exposure) by a wild, carnivorous mammal or a bat that is not available for testing should be regarded as having been exposed to rabies.
- (a) **DOGS, CATS, AND FERRETS.** Unvaccinated dogs, cats, and ferrets exposed to a rabid animal should be euthanized immediately. If the owner is unwilling to have this done, the animal should be placed in strict isolation for 6 months and vaccinated 1 month before being released. Animals with expired vaccinations need to be evaluated on a case-by-case basis. Protocols for the postexposure vaccination of previously unvaccinated domestic animals have not been validated, and there is evidence that the use of vaccine alone will not prevent the disease¹⁹. Dogs, cats, and ferrets that are currently vaccinated should be revaccinated immediately, kept under the owner's control, and observed for 45 days.
 - (b) **LIVESTOCK.** All species of livestock are susceptible to rabies; cattle and horses are among the most frequently infected. Livestock exposed to a rabid animal and currently vaccinated with a vaccine approved by USDA for that species should be revaccinated immediately and observed for 45 days. Unvaccinated livestock should be slaughtered immediately. If the owner is unwilling to have this done, the animal should be kept under close observation for 6 months.

The following are recommendations for owners of unvaccinated livestock exposed to rabid animals:

- (1) If the animal is slaughtered within 7 days of being bitten, its tissues may be eaten without risk of infection, provided that liberal portions of the exposed area are discarded. Federal guidelines for meat inspectors require that any animal known to have been exposed to rabies within 8 months be rejected for slaughter.
 - (2) Neither tissues nor milk from a rabid animal should be used for human or animal consumption²⁰. Pasteurization temperatures will inactivate rabies virus, therefore, drinking pasteurized milk or eating cooked meat does not constitute a rabies exposure.
 - (3) Having more than one rabid animal in a herd or having herbivore-to-herbivore transmission is uncommon; therefore, restricting the rest of the herd if a single animal has been exposed to or infected by rabies might not be necessary.
- (c) **OTHER ANIMALS.** Other mammals bitten by a rabid animal should be euthanized immediately. Animals maintained in USDA licensed research facilities or accredited zoological parks should be evaluated on a case-by-case basis.
- 6. MANAGEMENT OF ANIMALS THAT BITE HUMANS:**
- (a) **DOGS, CATS, AND FERRETS.** Rabies virus may be excreted in the saliva of infected dogs, cats, and ferrets during illness and/or for only a few days prior to illness or death²¹⁻²³. A healthy dog, cat, or ferret that bites a person should be confined and observed daily for 10 days; administration of rabies vaccine is not recommended during the observation period. Such animals should be evaluated by a veterinarian at the first sign of illness during confinement. Any illness in the animal should be reported immediately to the local health department. If signs suggestive of rabies develop, the animal should be euthanized and the head shipped for testing as described in Part II.A.7. Any stray or unwanted dog, cat, or ferret that bites a person may be euthanized immediately and the head submitted for rabies examination.
 - (b) **OTHER BITING ANIMALS.** Other biting animals which might have exposed a person to rabies should be reported immediately to the local health department. Prior vaccination of an animal may not preclude the necessity for euthanasia

and testing if the period of virus shedding is unknown for that species. Management of animals other than dogs, cats, and ferrets depends on the species, the circumstances of the bite, the epidemiology of rabies in the area, the biting animal's history, current health status, and potential for exposure to rabies.

C. PREVENTION AND CONTROL METHODS RELATED TO WILDLIFE: The public should be warned not to handle or feed wild mammals. Wild mammals and hybrids that bite or otherwise expose persons, pets, or livestock should be considered for euthanasia and rabies examination. A person bitten by any wild mammal should immediately report the incident to a physician who can evaluate the need for antirabies treatment (See current rabies prophylaxis recommendations of the ACIP⁷). State regulated wildlife rehabilitators may play a role in a comprehensive rabies control program. Minimum standards for persons who rehabilitate wild mammals should include rabies vaccination, appropriate training and continuing education. Translocation of infected wildlife has contributed to the spread of rabies^{24,25}; therefore, the translocation of known terrestrial rabies reservoir species should be prohibited.

1. TERRESTRIAL MAMMALS: The use of licensed oral vaccines for the mass vaccination of free-ranging wildlife should be considered in selected situations, with the approval of the state agency responsible for animal rabies control⁹. The distribution of oral rabies vaccine should be based on scientific assessments of the target species and followed by timely and appropriate analysis of surveillance data; such results should be provided to all stakeholders. Continuous and persistent programs for trapping or poisoning wildlife are not effective in reducing wildlife rabies reservoirs on a statewide basis. However, limited control in high-contact areas (e.g., picnic grounds, camps, suburban areas) may be indicated for the removal of selected high-risk species of wildlife⁹. State agriculture, public health, and wildlife agencies should be consulted for planning, coordination, and evaluation of vaccination or population-reduction programs.

2. BATS: Indigenous rabid bats have been reported from every state except Hawaii, and have caused rabies in at least 40 humans in the United States²⁶⁻²⁹. Bats should be excluded from houses, public buildings, and adjacent structures to prevent direct association with humans³⁰⁻³¹. Such structures should then be made bat-proof by sealing entrances used by bats. Controlling rabies in bats through programs designed to reduce bat populations is neither feasible nor desirable.

Part III: Rabies Vaccines Licensed and Marketed in the U.S., 2004

Product Name	Produced by	Marketed by	For Use In	Dosage	Age at Primary Vaccination ^a	Booster Recommended	Route of Inoculation
A) MONOVALENT (Inactivated) continued							
DEFENSOR 1	Pfizer, Incorporated License No. 189	Pfizer, Incorporated	Dogs Cats	1 ml 1 ml	3 months ^b 3 months	Annually Annually	IM ^c or SC ^d SC
DEFENSOR 3	Pfizer, Incorporated License No. 189	Pfizer, Incorporated	Dogs Cats Sheep Cattle	1 ml 1 ml 2 ml 2 ml	3 months 3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually Annually	IM or SC SC IM IM
RABDOMUN 1	Pfizer, Incorporated License No. 189	Schering-Plough	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	IM or SC SC
RABDOMUN	Pfizer, Incorporated License No. 189	Schering-Plough	Dogs Cats Sheep Cattle	1 ml 1 ml 2 ml 2 ml	3 months 3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually Annually	IM or SC SC IM IM
RABVAC 1	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	IM or SC IM or SC
RABVAC 3	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats Horses	1 ml 1 ml 2 ml	3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually	IM or SC IM or SC IM
RABVAC 3 TF	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Dogs Cats Horses	1 ml 1 ml 2 ml	3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually	IM or SC IM or SC IM
PRORAB-1	Intervet, Incorporated License No. 286	Intervet, Incorporated	Dogs Cats Ferrets Sheep	1 ml 1 ml 1 ml 2 ml	3 months 3 months 3 months 3 months	Annually Annually Annually Annually	IM or SC IM or SC IM or SC IM
PRORAB-3F	Intervet, Incorporated License No. 286	Intervet, Incorporated	Cats	1 ml	3 months	1 year later & triennially	IM or SC
IMRAB 3	Merial, Incorporated License No. 298	Merial, Incorporated	Dogs Cats Sheep Cattle Horses Ferrets	1 ml 1 ml 2 ml 2 ml 2 ml 1 ml	3 months 3 months 3 months 3 months 3 months 3 months	1 year later & triennially 1 year later & triennially 1 year later & triennially Annually Annually Annually	IM or SC IM or SC IM or SC IM or SC IM or SC SC
IMRAB 3 TF	Merial, Incorporated License No. 298	Merial, Incorporated	Dogs Cats Ferrets	1 ml 1 ml 1 ml	3 months 3 months 3 months	1 year later & triennially 1 year later & triennially Annually	IM or SC IM or SC SC

Product Name	Produced by	Marketed by	For Use In	Dosage	Age at Primary Vaccination ^a	Booster Recommended	Route of Inoculation
A) MONOVALENT (Inactivated) continued							
IMRAB Large Animal	Merial, Incorporated License No. 298	Merial, Incorporated	Cattle Horses Sheep	2 ml 2 ml 2 ml	3 months 3 months 3 months	Annually Annually 1 year later & triennially	IM or SC IM or SC IM or SC
IMRAB 1	Merial, Incorporated License No. 298	Merial, Incorporated	Dogs Cats	1 ml 1 ml	3 months 3 months	Annually Annually	SC SC
B) MONOVALENT (Rabies glycoprotein, live canary pox vector)							
PUREVAX Feline Rabies	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	2 months	Annually	SC
C) COMBINATION (Inactivated rabies)							
ECLIPSE 3 + FeLV/R	Fort Dodge Animal Health License No. 112	Schering-Plough	Cats	1 ml	3 months	Annually	IM or SC
ECLIPSE 4 + FeLV/R	Fort Dodge Animal Health License No. 112	Schering-Plough	Cats	1 ml	3 months	Annually	IM or SC
Fel-O-Guard 3 + FeLV/R	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Cats	1 ml	3 months	Annually	IM or SC
Fel-O-Guard 4 + FeLV/R	Fort Dodge Animal Health License No. 112	Fort Dodge Animal Health	Cats	1 ml	3 months	Annually	IM or SC
IMRAB 3 + Feline 3	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1 ml	3 months	1 year later & triennially	SC
IMRAB 3 + Feline 4	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1 ml	3 months	1 year later & triennially	SC
Equine POTOMAVAC + IMRAB	Merial, Incorporated License No. 298	Merial, Incorporated	Horses	1 ml	3 months	Annually	IM
MYSTIQUE II POTOMAVAC +	Intervet, Incorporated License No. 286	Intervet, Incorporated	Horses	1 ml	3 months	Annually	IM
D) COMBINATION (Rabies glycoprotein, live canary pox vector)							
PUREVAX Feline 3/ Rabies	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
PUREVAX Feline 4/ Rabies	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
PUREVAX Feline 3/ Rabies + LEUCAT	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
PUREVAX Feline 4/ Rabies + LEUCAT	Merial, Incorporated License No. 298	Merial, Incorporated	Cats	1ml	8 weeks	Annually	SC
E) ORAL (Rabies glycoprotein, live vaccinia vector) - RESTRICTED TO USE IN STATE AND FEDERAL RABIES CONTROL PROGRAMS							
RABORAL V-RG	Merial, Incorporated License No. 298	Merial, Incorporated	Raccoons Coyotes	N/A	N/A	As determined by local authorities	Oral

- a. Minimum age (or older) and revaccinated one year later.
- b. A month = 28 days
- c. Intramuscularly
- d. Subcutaneously

REFERENCES

1. Rabies. In: Chin J, ed. *Control of Communicable Diseases Manual, 17th ed.* Washington, DC: American Public Health Association, 2000:411-419.
2. Model rabies control ordinance. In: Directory and resource manual. Schaumburg, IL: American Veterinary Medical Association, 2002:114-116.
3. Bunn TO. Canine and feline vaccines, past and present. In Baer GM, ed. *The Natural History of Rabies, 2nd ed.* Boca Raton, FL: CRC Press, 1991:415-425.
4. Gobar GM, Kass PH. World Wide Web-based survey of vaccination practices, postvaccinal reactions, and vaccine site-associated sarcomas in cats, *J Am Vet Med Assoc* 2002; 220:1477-1482
5. Macy DW, Hendrick MJ. The potential role of inflammation in the development of postvaccinal sarcomas in cats. *Vet Clin North Am Small Anim Pract* 1996; 26:103-109.
6. Rupprecht CE, Blass L, Smith K et al. Human infection due to recombinant vaccinia-rabies glycoprotein virus. *N Engl J Med* 2001; 345:582-586.
7. CDC. Human rabies prevention-United States, 1999. Recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1999; 48:(No. RR-1).
8. Krebs JW, et al. Rabies Surveillance in the United States during 2002. *J Am Vet Med Assoc* 2003.
9. Hanlon CA, Childs JE, Nettles VF, et al. Recommendations of the Working Group on Rabies, Article III: Rabies in wildlife. *J Am Vet Med Assoc* 1999; 215:1612-1618.
10. Hanlon CA, Smith, JS, Anderson, GR, et al. Recommendations of the Working Group on Rabies, Article II: Laboratory diagnosis of rabies. *J Am Vet Med Assoc* 1999; 215:1444-1446.
11. 2000 Report of the AVMA Panel on Euthanasia. *J Am Vet Med Assoc* 2001; 218:5,669-696.
12. Tizard I, Ni Y. Use of serologic testing to assess immune status of companion animals. *J Am Vet Med Assoc* 1998; 213:54-60.
13. Wild Animals as Pets. In: Directory and resource manual. Schaumburg, IL: American Veterinary Medical Association 2002:126.
14. Position on Canine Hybrids. In: Directory and resource manual. Schaumburg, IL: American Veterinary Medical Association 2002:88-89.
15. Siino BS. Crossing the line. American Society for the Prevention of Cruelty to Animals Animal Watch 2000; Winter:22-29.
16. Jay MT, Reilly KF, DeBess EE, Haynes EH, Bader DR, Barrett LR. Rabies in a vaccinated wolf-dog hybrid. *J Am Vet Med Assoc* 1994; 205:1729-1732.
17. CDC. An imported case of rabies in an immunized dog. *MMWR* 1987, 36:94-96,101.
18. CDC. Imported dog and cat rabies – New Hampshire, California. *MMWR* 1988, 37:559-560.
19. Hanlon CA, Niezgodna MN, Rupprecht CE. Postexposure prophylaxis for prevention of rabies in dogs. *Am J Vet Res* 2002; 63:1096-1100.
20. CDC. Mass treatment of humans who drank unpasteurized milk from rabid cows-Massachusetts, 1996-1998. *MMWR* 1999; 48:228-229.
21. Vaughn JB, Gerhardt P, Paterson J. Excretion of street rabies virus in saliva of cats. *J Am Vet Med Assoc* 1963, 184:705.
22. Vaughn JB, Gerhardt P, Newell KW. Excretion of street rabies virus in saliva of dogs. *J Am Vet Med Assoc* 1965, 193:363-368.
23. Niezgodna M, Briggs DJ, Shaddock J, Rupprecht CE. Viral excretion in domestic ferrets (*Mustela putorius furo*) inoculated with a raccoon rabies isolate. *Am J Vet Res* 1998, 59:1629-1632.
24. Jenkins SR, Perry BD, Winkler WG. Ecology and epidemiology of raccoon rabies. *Rev Inf Dis* 1988; 10:Suppl 4:S620-S625.
25. CDC. Translocation of Coyote Rabies – Florida, 1994. *MMWR* 44:580-587.
26. Messenger SL, Smith JS, Rupprecht CE. Emerging epidemiology of bat-associated cryptic cases of rabies in humans in the United States. *Clin Inf Dis* 2002; 35:738-747.
27. CDC. Human Rabies – California, 2002. *MMWR* 51:686-688.
28. CDC. Human Rabies – Tennessee, 2002. *MMWR* 51:828-829.
29. CDC. Human Rabies – Iowa, 2002. *MMWR* 52:47-48.
30. Frantz SC, Trimarchi CV. Bats in human dwellings: health concerns and management In: Decker DF, ed. Proceedings of the first eastern wildlife damage control conference. Ithaca, NY: Cornell University, 1983: 299-308.
31. Greenhall AM. House bat management. US Fish and Wildlife Service, Resource Publication 143, 1982.