


Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using a most or moderately effective* form of birth control

Among women who were not trying to get pregnant and who were not currently pregnant

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes to a most or moderately effective method



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	45.0	41.9	48.0	1,220	57,799
Maternal Age	<20 yrs	41.8	27.7	57.3	49	2,132
	20 to 24 yrs	49.4	41.4	57.4	187	10,037
	25 to 29 yrs	45.4	39.8	51.1	345	17,055
	30 to 34 yrs	43.5	38.3	48.9	386	17,285
	=>35 yrs	43.8	37.3	50.5	253	11,290
Race/Ethnicity	NH White	43.4	39.4	47.6	646	29,713
	NH Black	41.1	34.2	48.5	237	9,147
	Hispanic	46.6	39.8	53.6	238	12,840
	Other	61.5	50.3	71.6	90	5,735.2
Education	<12 yrs	46.9	37.5	56.5	134	6,418
	12 yrs	44.2	37.9	50.8	275	13,527
	>12 yrs	45.1	41.4	48.9	790	36,775
Birth Weight	Low	50.8	45.2	56.4	310	4,816
	Normal	44.5	41.3	47.8	910	52,983
Marital Status	Married	46.1	42.4	50.0	750	36,026
	Other	43.2	38.2	48.3	470	21,773
Insurer Now	Medicaid	42.3	37.3	47.5	448	19,963
	Private	46.6	42.5	50.7	639	31,285
	Uninsured	43.0	33.1	53.5	112	5,024

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents or numerator <6 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Most Moderately Effective Method includes male/female sterilization, birth control pills, implant (e.g., Nexplanon®, Implanon®), IUD (e.g., Mirena®, ParaGard®, Liletta®, Skyla®), contraceptive patch (e.g. OrthoEvra®) or shot (Depo-Provera®), or vaginal ring (e.g., NuvaRing®).