

Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using the following birth control methods

Among women reporting using birth control now

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Condoms	29.9	27.0	33.0	1,029	32,786
Birth control pills	24.9	22.1	27.9	1,029	27,301
IUD (including Mirena®, ParaGard®, Liletta®, or Skyla®)	16.1	13.8	18.7	1,029	17,606
Withdrawal (pulling out)	15.1	12.9	17.6	1,029	16,529
Not having sex (abstinence)	8.3	6.7	10.3	1,029	9,134
Tubes tied or blocked (female sterilization or Essure®)	7.6	6.1	9.6	1,029	8,385
Shots or injections (Depo-Provera®)	7.0	5.5	8.9	1,029	7,673
Natural family planning (including rhythm method)	5.5	4.2	7.2	1,029	6,039
Contraceptive implant in the arm (Nexplanon® or Implanon®)	4.0	2.9	5.6	1,029	4,437
Vasectomy (male sterilization)	3.8	2.7	5.2	1,029	4,119
Contraceptive patch (OrthoEvra®) or vaginal ring (NuvaRing®)	2.0	1.2	3.2	1,029	2,170

¹Percentage = Weighted percentage²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents or numerator <6 respondents

³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition