


Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using a most or moderately effective* form of birth control

Among women who were not trying to get pregnant and who were not currently pregnant

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes to a most or moderately effective method



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	53.2	50.0	56.3	1,166	66,424
Maternal Age	<20 yrs	57.2	40.5	72.3	43	2,574
	20 to 24 yrs	64.6	56.2	72.2	179	15,568
	25 to 29 yrs	52.6	46.6	58.5	315	17,122
	30 to 34 yrs	51.1	45.6	56.6	367	19,599
	=>35 yrs	45.5	38.9	52.2	262	11,561
Race/Ethnicity	NH White	51.8	47.5	56.2	606	33,757
	NH Black	56.2	47.9	64.1	194	11,429
	Hispanic	58.7	52.0	65.1	259	16,946
	Other	38.9	28.5	50.4	96	3585.2
Education	<12 yrs	55.8	45.5	65.6	121	7,035
	12 yrs	57.2	50.2	63.9	254	16,553
	>12 yrs	51.5	47.6	55.3	780	42,222
Birth Weight	Low	56.6	51.1	62.0	314	5,250
	Normal	52.9	49.5	56.3	852	61,174
Marital Status	Married	46.9	42.9	50.9	708	34,561
	Other	62.3	57.1	67.2	458	31,862
Insurer Now	Medicaid	60.8	55.2	66.2	402	25,880
	Private	48.6	44.3	52.8	632	32,668
	Uninsured	54.9	45.0	64.5	113	7,334

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents or numerator <6 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Most Moderately Effective Method includes male/female sterilization, birth control pills, implant (e.g., Nexplanon®, Implanon®), IUD (e.g., Mirena®, ParaGard®, Liletta®, Skyla®), contraceptive patch (e.g. OrthoEvra®) or shot (Depo-Provera®), or vaginal ring (e.g., NuvaRing®).